



OPEN ACCESS

The problem with root cause analysis

Mohammad Farhad Peerally,¹ Susan Carr,² Justin Waring,³ Mary Dixon-Woods¹

¹SAPPHIRE, Department of Health Sciences, University of Leicester, Leicester, UK

²John Walls Renal Unit, University Hospitals of Leicester, Leicester, UK

³CHILL, Nottingham University Business School, University of Nottingham, Nottingham, UK

Correspondence to

Dr Mohammad Farhad Peerally, SAPPHIRE, Department of Health Sciences, University of Leicester, Centre for Medicine University Road Leicester, LE1 7RH UK; mfp6@le.ac.uk

Accepted 20 May 2016

INTRODUCTION

Attempts to learn from high-risk industries such as aviation and nuclear power have been a prominent feature of the patient safety movement since the late 1990s. One noteworthy practice adopted from such industries, endorsed by health-care systems worldwide for the investigation of serious incidents,^{1–3} is root cause analysis (RCA). Broadly understood as a method of structured risk identification and management in the aftermath of adverse events,¹ RCA is not a single technique. Rather, it describes a range of approaches and tools drawn from fields including human factors and safety science^{4,5} that are used to establish how and why an incident occurred in an attempt to identify how it, and similar problems, might be prevented from happening again.⁶ In this article, we propose that RCA does have potential value in healthcare, but it has been widely applied without sufficient attention paid to what makes it work in its contexts of origin, and without adequate customisation for the specifics of healthcare.^{7,8} As a result, its potential has remained under-realised⁷ and the phenomenon of organisational forgetting⁹ remains widespread (box 1). Here, we identify eight challenges facing the usage of RCA in healthcare and offer some proposals on how to improve learning from incidents.

The unhealthy quest for ‘the’ root cause

The first problem with RCA is its name. By implying—even inadvertently—that a single root cause (or a small number of causes) can be found, the term ‘root cause analysis’ promotes a flawed reductionist view.¹⁰ Incident investigation in the aftermath of an adverse event is intended to identify the latent and active factors contributing to the genesis of a particular adverse event,⁴ but too often results in a simple linear narrative that

Box 1 Lessons not learnt

This example provides a summary of a real case that occurred in a hospital and the failure to learn from the incident in spite of a root cause analysis.

In a large acute hospital, a patient underwent a routine cataract surgery—an operation with a minimal risk profile—led by an experienced ophthalmologist. The wrong lens was inserted during the operation. The error was promptly recognised postoperatively; the patient was returned to the operating room and the procedure was safely redone.

A subsequent root cause analysis identified that two lenses were in the operating room, one (the wrong one) brought in by an operating department assistant and the other by the surgeon. The investigation report identified that having more than one lens in the operating room and a failure in the double-checking process had caused the incident. The action plan included the development of a new protocol emphasising the individual responsibility of the surgeon to select the appropriate lens, a training programme, improved documentation and a poster emphasising the importance of double checks.

One year later, in the same hospital, a different patient with a different surgeon had the same procedure. Once again, the wrong lens was implanted. This time, the staff member who chose the wrong lens was the surgeon.

displaces more complex, and potentially fruitful, accounts of multiple and interacting contributions to how events really unfold.^{7,10–12} This is a tendency exacerbated by use of some RCA techniques (such as timelines or the ‘five whys’) that tend to favour a temporal narrative rather than a wider systems view.

To cite: Peerally MF, Carr S, Waring J, et al. *BMJ Qual Saf* Published Online First: [please include Day Month Year] doi:10.1136/bmjqs-2016-005511

The problem with...

Questionable quality of RCA investigations

Once an adverse event is classified as meeting the definition of a serious incident, an RCA is supposed to involve the convening of a skilled multidisciplinary investigation team, preferably with representations from risk management personnel and clinical teams.¹³ Over a predefined timeframe, which is mandated in some countries (60 days in the UK, 45 days in the USA),^{3 14} this team collects and analyses data and formulates an action plan.¹³ However, challenges pertaining to the quality of this process abound. The task facing the investigation team is far from straightforward: the events underlying an incident have to be reconstructed from many different sources of varying degrees of reliability, usefulness and accessibility, ranging from hospital records, staff interviews and statements, to records of workforce rotas.¹⁵ The information obtained directly from healthcare workers is influenced by their willingness and ability to provide relevant data^{16 17} and by nature of the relationships and conversations between investigators and other stakeholders.¹⁸ The involvement of patients and families affected by the incident is wildly variable, with only limited evidence-based guidance on how it can best be done.¹⁹ Yet, despite the complexities, sensitivities and challenges of this work, RCAs in healthcare are typically conducted by local teams, not the expert accident investigators who are proficient in systems thinking and human factors, cognitive interviewing, staff engagement and data analysis that are characteristic of other high-risk industries.^{20–22} Further, inconsistent use is made of the various investigative tools that are available.^{15 23} As a result, exemplary practice in the analysis of healthcare incidents is rare.^{24 25}

Political hijack

Constrained by strict timelines, and skewed by hindsight bias²⁶ and lack of independence from the organisation where the event took place, RCAs in healthcare often end up a compromise between ‘depth of data and accuracy of the investigation’.¹⁶ The quest to complete an investigation on time and produce a report risks goal displacement, where the report is seen as the end product rather than the beginning of a learning cycle. Reports themselves, influenced by the need to preserve interpersonal relationships and by hierarchical tensions and partisan interests,^{8 16 27–29} may not always reflect the content of discussions during investigations nor the realities of what happened.^{15 16} Investigating teams may end their analysis once they have reached a cause of mutual convenience, perhaps one that edits out causes (and thus solutions) deemed to be beyond the remit or capacities of the organisation¹⁶ and that occludes deeper organisational and socio-political dynamics.^{7 15 16}

Poorly designed or implemented risk controls

The key goal of RCA is to prevent similar events from recurring.^{7 10 16} But few studies have investigated the nature and effectiveness of risk control strategies stemming from RCA investigations in healthcare. The available evidence points to the endemic tendency of investigators to settle for administrative and perhaps ‘weaker’ solutions (such as reminders) rather than those that address the latent causes, such as poorly designed technology or defective operational systems.^{8 16 30–32} Again, some of the reasons for this lie in the limited expertise of local investigation teams in selecting and designing appropriate risk controls.^{7 30 33} Only limited guidance is available^{21 33–37} and what is available may not be sufficiently attentive to the specifics of the healthcare context. Yet poorly designed or ineffectual corrective actions may do harm.^{31 32} Among other unintended consequences, risk migration, where attempts to mitigate a risk create new risks, may easily occur.^{38 39} Recommended actions may also, of course, result in little change,^{7 30 40} especially (but not only) when senior managers are not involved in the generation of action plans and do not support their implementation.^{41 42} Despite the time and effort invested in RCAs,^{7 40 43} few incentives exist to follow-up formally on action plans:^{8 44} estimates of implementation rates vary between 45% and 70%.^{31 45}

Poorly functioning feedback loops

For learning to occur, several conditions must be satisfied. Among the most important of these are the sharing of the outcomes of incident analysis with those involved, those who reported, and those likely to be affected in the future, especially in implementing recommendations. Evidence in other fields suggests that learning from events does not happen by itself:⁴⁶ purposeful intent is needed both to disseminate the findings⁴⁷ and ensure that the recommended actions made salient and actionable.⁴⁶ Yet, as currently practised, feedback mechanisms in healthcare RCAs function poorly, contributing to the disenchantment of staff^{48 49} and frustrating the kind of double-loop learning⁵⁰ needed to secure change.

Disaggregated analysis focused on single organisations and incidents

The current RCA approach favours analysis of individual incidents in isolation and within bounded organisations. The consequent tendency to generate localised action plans that are not shared more widely may result in failure to disseminate painfully acquired learning and to address deeper, institutionally engrained patient safety concerns.^{21 51} Single incident analysis also frustrates the organisation’s ability to assess its vulnerability to recurring events.⁵² Organisations’ inability to effectively prioritise actions may lead to an unwarranted commitment of resources to averting specific very rare events rather than addressing the conditions that

allowed the event to occur. Though mechanisms for aggregating learning from incidents and creating alerts do exist in some countries, their impact to date has been limited: similar events often recur in the same or similar organisations (box 1), suggesting failure to learn both within and across organisations.²⁴

Confusion about blame

Though healthcare is often exhorted to embrace a ‘no-blame’ culture, the extent to which this urging is based on a correct understanding of what happens in other high-risk industries is questionable.^{15 53} Investigators in other industries do not set out with a remit to assign blame,²⁰ but that does not mean that individual or organisational culpability is forever sequestered. The vast majority of mistakes and other errors are the result of systems defects that need to be corrected, but when blatant transgressions, neglect or unacceptable behaviour is found, it is clearly wrong to write accountability out of the picture.⁵⁴ Nor is no-blame the reality in practice, since disciplinary, institutional and legal (civil and criminal) processes continue to operate and are highly visible to healthcare practitioners and managers, yet may appear arbitrary and unsatisfactory both to them and to patients and families. A ‘just culture’ is increasingly promoted in many organisations to balance the disparity between individual blame and organisational accountability.⁵⁵ This approach, however, comes with problems of its own when applied to incident investigation in healthcare. For instance, some of the more visible features of the just culture philosophy in incident investigation is the use of prescriptive algorithms and decision tools (such as culpability tree) to objectify culpability. Such ‘calculus-like logic’⁵⁶ may imply that actions committed by staff are binary (either acceptable or unacceptable) without appropriate appreciation of the messiness of the system in which the action occurred.^{56 57}

The problem of many hands

RCA is further challenged by the problem of many hands, which describes the problem that many actors and their actions may contribute to an outcome, yet no individual is responsible either for that outcome or for fixing the problems that caused it.⁵⁸ This problem, which is endemic in healthcare, makes it difficult to address hazards that arise at the level of the system, since many of the actors that are implicated in hazards—including, for example, drug and equipment suppliers—are outside the direct control of individual care organisations. RCA investigations may fail to assign responsibility to such actors, instead reabsorbing responsibility into the organisation where the incident occurred. These organisations typically lack the legal mandate, resources and structural authority necessary to make the changes required.

DISCUSSION

RCA is a promising approach with considerable face validity as a way of producing learning from things that have gone wrong. But it has consistently failed to deliver benefits on the scale or quality needed. The eight problems we have discussed here mean that, too often, RCA results in the tombstone effect: though its purpose is to guard against a similar incident in the future, it may instead function primarily as a procedural ritual, leaving behind a memorial that does little more than allow a claim that something has been done.^{59 60} Incident investigation clearly will continue to play an important role in making healthcare safer, but it must first get better at doing what it does.

The first step in securing improvement is likely to involve the professionalisation of incident investigation: those conducting it need specialist expertise in underlying theories, ergonomics, human factors and hands-on experience of analytical methods.²⁰ For these reasons, the establishment of professional investigatory bodies, such as the one shortly to be launched in the UK, are welcome—though the scope, reach and impact of such bodies will need careful monitoring. Second, the role of patients and relatives in the investigative process needs to be recognised and valued. Such engagement has the potential to generate a unique perspective of the service provided from the end-user’s perspective and may foster dialogue that is informative to both causal analysis and design of risk controls.⁶¹ The psychological and emotional readiness of patients and families involved in the investigative process needs to be considered, along with the maturity and ability of the organisation to facilitate such a process within the appropriate legal framework. Transparency on the agreed level of involvement is paramount from the start and the outcomes of investigations should be available to patients and relatives, though clarity on how this should best be done is not yet available.¹⁹ Third, better understanding of the role of blame is needed. The dissonance caused by claims of no-blame or even just culture and the reality is a source of confusion and distress in relation to RCAs. To address current confusions, clarity is needed on the distribution of responsibility between bodies investigating incidents (whose prime mandate would be to promote learning) and other bodies (including professional regulators and the law courts), and in what instances the investigative body needs to make referrals.²²

Fourth, healthcare must focus increasingly on aggregated analysis of incidents.^{31 32 45 62 63} Such a bird’s eye view of incidents may facilitate prioritisation of interventions, based on the harm associated with incidents and also on the associated risks. Aggregated analyses can be performed at numerous levels of the organisational hierarchy, for example, the micro (within one department) and at the meso level (organisational).⁴¹ At the national level, aggregated analyses offer a way of identifying common themes across

The problem with...

similar and apparently more disparate incidents^{31 32 45} and may also serve as a means of generating actions that require collaborative efforts between healthcare organisations or indeed between industry and healthcare. Such an example could be for instance product redesigning—a solution that may not be identified through the analysis of a single incident within one department but may reveal itself as a recurring theme when analysing multiple incidents across many organisations. Linked to this, healthcare urgently needs to develop and evaluate much better methods for designing risk controls and other improvement actions. One possibility that could be evaluated, for example, is that of a hierarchy of risk controls.^{33 34 36 37 64} More broadly, the use of active surveillance of issues that have already been detected and monitoring of effectiveness of risk controls need to become a routine part of the risk management process following RCAs. Healthcare also needs to markedly improve its capacity to evaluate, curate and share these risk controls. Such an approach would help to address the problem that organisations tend to constantly reinvent risk controls, resulting in waste and the creation of new risks.⁵⁸ An easily accessible database with descriptions of risk controls and contexts would enable lessons learnt from one RCA to be shared widely and support a participatory approach⁶⁵ to organisational learning.

Finally, healthcare needs to do more to detect hazards and assess risks proactively. RCA is essentially retrospective, and depends crucially on an incident being recognised as such, but that may not happen for a variety of reasons: healthcare personnel may have become habituated to particular practices or outcomes, or fear and other negative emotions discourage reporting. Though RCAs were imported from other high-risk industries, the other tools and techniques commonly used in those industries to assess systems and assure their safety before an incident has occurred—such as failure modes and effects analysis (FMEA), hierarchical task analysis and so on—have had far less attention in healthcare⁶⁶ FMEA, in particular, may be especially useful for the rigorous proactive risk assessment of a select few but high-priority hazards.⁶⁷ For healthcare truly to become a learning system, action is needed on multiple levels. RCAs have dominated for too long as the principal means of generating learning. The time has come to recognise both their opportunities and their limits.

- ▶ RCA is a promising incident investigation technique borrowed from other high-risk industries, but has failed to live up to its potential in healthcare.
- ▶ A key problem with RCA is its name, which implies a singular, linear cause.
- ▶ Other problems include the questionable quality of many RCAs, their susceptibility to political hijack, their tendency to produce poor risk controls, poorly functioning feedback loops, failure to aggregate learning across incidents and confusion about blame and responsibility.

- ▶ Implementation and evaluation of risk controls to eliminate or minimise identified hazards need to become a more visible feature of the RCA process.
- ▶ To maximise learning, lessons learnt from incidents, descriptions of implemented risk controls and their effectiveness need to be shared within and across organisations.

Twitter Follow Mohammad Peerally at @FP_Farhad

Contributors MFP wrote the first draft of the manuscript, which was subsequently revised critically and edited by SC, JW and MD-W. MD-W edited the final version of the manuscript. All authors approved the final manuscript version being submitted for publication.

Funding Wellcome Trust Senior Investigator Award (Mary Dixon-Woods (WT097899)) and Health Foundation (Improvement Science Doctoral Awards).

Competing interests MD-W is the Deputy Editor-in-Chief of *BMJ Quality and Safety*.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Bagian JP, Gosbee J, Lee CZ, *et al*. The Veterans Affairs root cause analysis system in action. *Jt Comm J Qual Patient Saf* 2002;28:531–45.
- 2 Braithwaite J, Westbrook MT, Mallock NA, *et al*. Experiences of health professionals who conducted root cause analyses after undergoing a safety improvement programme. *Qual Saf Health Care* 2006;15:393–9.
- 3 NHS England. Serious incident framework. Supporting Learning to Prevent Recurrence. 2015. <http://www.england.nhs.uk/wp-content/uploads/2015/04/serious-incident-framework-upd.pdf> (accessed 31 Dec 2015).
- 4 Reason J. *Human error*. Cambridge University Press, 1990.
- 5 Vincent C, Taylor-Adams S, Stanhope N. Framework for analysing risk and safety in clinical medicine. *BMJ* 1998;316:1154–7.
- 6 US Department of Veterans Affairs. VA National Center for Patient Safety: Root Cause Analysis. 2015. <http://www.patientsafety.va.gov/professionals/onthejob/rca.asp> (accessed 31 Dec 2015).
- 7 Wu AW, Lipshutz AK, Pronovost PJ. Effectiveness and efficiency of root cause analysis in medicine. *JAMA* 2008;299:685–7.
- 8 Percarpio KB, Watts BV, Weeks WB. The effectiveness of root cause analysis: what does the literature tell us? *Jt Comm J Qual Patient Saf* 2008;34:391–8.
- 9 Haunschild PR, Polidoro F Jr, Chandler D. Organizational oscillation between learning and forgetting: the dual role of serious errors. *Organ Sci* 2015;26:1682–701.
- 10 Vincent CA. Analysis of clinical incidents: a window on the system not a search for root causes. *Qual Saf Health Care* 2004;13:242–3.
- 11 National Patient Safety Foundation (NPSF). RCA²: improving root cause analyses and actions to prevent harm. 2016. <http://c>

- ymcdn.com/sites/www.npsf.org/resource/resmgr/PDF/RCA2_v2-online-pub_010816.pdf (accessed 2 Mar 2016).
- 12 Iedema RAM, Jorm C, Long D, *et al*. Turning the medical gaze in upon itself: root cause analysis and the investigation of clinical error. *Soc Sci Med* 2006;62:1605–15.
 - 13 Woloshynowych M, Rogers S, Taylor-Adams S, *et al*. The investigation and analysis of critical incidents and adverse events in healthcare. *Health Technol Assess* 2005;9:i–143.
 - 14 The Joint Commission. Comprehensive accreditation manual for hospitals. Sentinel Event Policy and Procedures. 2016. http://www.jointcommission.org/assets/1/6/CAMH_24_SE_all_CURRENT.pdf (accessed 25 Jan 2016).
 - 15 Nicolini D, Waring J, Mengis J. The challenges of undertaking root cause analysis in health care: a qualitative study. *J Health Serv Res Policy* 2011;16(Suppl 1):34–41.
 - 16 Nicolini D, Waring J, Mengis J. Policy and practice in the use of root cause analysis to investigate clinical adverse events: mind the gap. *Soc Sci Med* 2011;73:217–25.
 - 17 Kewell B, Beck M. NHS inquiries: a time series analysis. *Public Money Manag* 2008;28:375–82.
 - 18 Carroll JS, Rudolph JW, Hatakenaka S. Lessons learned from non-medical industries: root cause analysis as culture change at a chemical plant. *Qual Saf Health Care* 2002;11:266–9.
 - 19 Zimmerman TM, Amori G. Including patients in root cause and system failure analysis: legal and psychological implications. *J Healthc Risk Manag* 2007;27:27–34.
 - 20 Macrae C. *Close calls: managing risk and resilience in airline flight safety*. Basingstoke: Palgrave Macmillan, 2014.
 - 21 Pham JC, Kim GR, Natterman JB, *et al*. ReCASTing the RCA: an improved model for performing root cause analyses. *Am J Med Qual* 2010;25:186–91.
 - 22 House of Commons' Public Administration Select Committee (PASC). *Investigating clinical incidents in the NHS*. Sixth Report of Session 2014–2015. 2015.
 - 23 Diller T, Helmrich G, Dunning S, *et al*. The Human Factors Analysis Classification System (HFACS) applied to health care. *Am J Med Qual* 2014;29:181–90.
 - 24 Keogh B. *Review into the quality of care and treatment provided by 14 hospital trusts in England: overview report*. NHS, 2013.
 - 25 Wallace LM. From root causes to safer systems: international comparisons of nationally sponsored healthcare staff training programmes. *Qual Saf Health Care* 2006;15:388–9.
 - 26 Henriksen K, Kaplan H. Hindsight bias, outcome knowledge and adaptive learning. *Qual Saf Health Care* 2003;12(Suppl 2):ii46–50.
 - 27 Wald H, Shojania KG. *Root Cause Analysis. Making health care safer: a critical analysis of patient safety practices*. Rockville, MD: Agency for Healthcare Research and Quality, 2001:51–8.
 - 28 Motschman TL, Moore SB. Corrective and preventive action. *Transfus Sci* 1999;21:163–78.
 - 29 Sweitzer SC, Silver MP. Learning from unexpected events: a root cause analysis training program. *J Healthcare Qual* 2005;27:11–19.
 - 30 Card AJ, Ward J, Clarkson PJ. Successful risk assessment may not always lead to successful risk control: a systematic literature review of risk control after root cause analysis. *J Healthcare Risk Manag* 2012;31:6–12.
 - 31 Mills PD, Neily J, Kinney LM, *et al*. Effective interventions and implementation strategies to reduce adverse drug events in the Veterans Affairs (VA) system. *Qual Saf Health Care* 2008;17:37–46.
 - 32 Mills PD, Neily J, Luan D, *et al*. Using aggregate root cause analysis to reduce falls and related injuries. *Jt Comm J Qual Patient Saf* 2005;31:21–31.
 - 33 Card AJ, Ward JR, Clarkson PJ. Generating Options for Active Risk Control (GO-ARC): introducing a novel technique. *J Healthc Qual* 2014;36:32–41.
 - 34 The National Institute for Occupational Safety and Health (NIOSH). Hierarchy of controls. 2015. <http://www.cdc.gov/niosh/topics/hierarchy/> (accessed 26 Jan 2016).
 - 35 Vacher A, d'Hollander A, El Mhamdi S, *et al*. Effectiveness of a tool for structuring action plan after analysis of adverse event. *Proceedings of the Human Factors and Ergonomics Society Annual Meeting*. Sage Publications, 2011;55:1631–4.
 - 36 Card AJ, Ward JR, Clarkson PJ. Rebalancing risk management—Part 2: The Active Risk Control (ARC) Toolkit. *J Healthc Risk Manag* 2015;34:4–17.
 - 37 Card AJ, Ward JR, Clarkson PJ. Rebalancing risk management—part 1: the process for active risk control (PARC). *J Healthc Risk Manag* 2014;34:21–30.
 - 38 Kirwan B. Incident reduction and risk migration. *Saf Sci* 2011;49:11–20.
 - 39 Deming WE. *The new economics: for industry, government, education*. MIT Press, 2000.
 - 40 Wrigstad J, Bergström J, Gustafson P. Mind the gap between recommendation and implementation—principles and lessons in the aftermath of incident investigations: a semi-quantitative and qualitative study of factors leading to the successful implementation of recommendations. *BMJ Open* 2014;4:e005326.
 - 41 Morse RB, Pollack MM. Root cause analyses performed in a children's hospital: events, action plan strength, and implementation rates. *J Healthc Qual* 2012;34:55–61.
 - 42 Iedema R, Jorm C, Braithwaite J. Managing the scope and impact of root cause analysis recommendations. *J Health Organ Manag* 2008;22:569–85.
 - 43 Rollenhagen C, Westerlund J, Lundberg J, *et al*. The context and habits of accident investigation practices: a study of 108 Swedish investigators. *Saf Sci* 2010;48:859–67.
 - 44 Hetteringer AZ, Fairbanks RJ, Hegde S, *et al*. An evidence-based toolkit for the development of effective and sustainable root cause analysis system safety solutions. *J Healthc Risk Manag* 2013;33:11–20.
 - 45 Mills PD, Neily J, Luan D, *et al*. Actions and implementation strategies to reduce suicidal events in the Veterans Health Administration. *Jt Comm J Qual Patient Saf* 2006;32:130–41.
 - 46 Ramanujam R, Goodman PS. The challenge of collective learning from event analysis. *Saf Sci* 2011;49:83–9.
 - 47 Peerally M, Carr S. Longitudinal patient safety education as a means to promote organizational learning in different resource settings. *Med Teach* 2016;38:532.
 - 48 Evans SM, Berry JG, Smith BJ, *et al*. Attitudes and barriers to incident reporting: a collaborative hospital study. *Qual Saf Health Care* 2006;15:39–43.
 - 49 Kingston M, Evans S, Smith B, *et al*. Attitudes of doctors and nurses towards incident reporting: a qualitative analysis. *Med J Aust* 2004;181:36–9.
 - 50 Argyris C, Schön DA. *Organizational learning: a theory of action perspective*. MA: Addison-Wesley Reading, 1978.
 - 51 Lundberg J, Rollenhagen C, Hollnagel E. What you find is not always what you fix—how other aspects than causes of accidents decide recommendations for remedial actions. *Accid Anal Prev* 2010;42:2132–9.
 - 52 Card AJ, Ward JR, Clarkson PJ. Trust-level risk evaluation and risk control guidance in the NHS east of England. *Risk Anal* 2014;34:1469–81.

The problem with...

- 53 Wachter RM, Pronovost PJ. Balancing “no blame” with accountability in patient safety. *N Engl J Med* 2009;361:1401–6.
- 54 Reason J. Engineering a just culture. In: Ashgate, ed. *Managing the risks of organizational accidents*, 1997:205–12.
- 55 Marx D. Patient safety and the “just culture:” a primer for health care executives. 2001. <http://www.safer.healthcare.ucla.edu/safer/archive/ahrq/FinalPrimerDoc.pdf> (accessed 5 Mar 2016).
- 56 Aveling E, Parker M, Dixon-Woods M. What is the role of individual accountability in patient safety? A multi-site ethnographic study. *Sociol Health Illn* 2016;38:216–32.
- 57 Dekker SWA, Hugh TB. A just culture after Mid Staffordshire. *BMJ Qual Saf* 2014;23:356–8.
- 58 Dixon-Woods M, Pronovost PJ. Patient safety and the problem of many hands. *BMJ Qual Saf* 2016; ■■■.
- 59 Dixon-Woods M, Yeung K, Bosk CL. Why is UK medicine no longer a self-regulating profession? The role of scandals involving “bad apple” doctors. *Soc Sci Med* 2011;73:1452–9.
- 60 Hood C, Rothstein H, Baldwin R. *The government of risk: understanding risk regulation regimes*. USA: Oxford University Press, 2001.
- 61 Iedema R, Allen S. Anatomy of an incident disclosure: the importance of dialogue. *Jt Comm J Qual Patient Saf* 2012;38:435–42.
- 62 National Patient Safety Agency. Root Cause Analysis Investigation Tools. A guide to aggregated and multi-incident RCA investigations 2008;0769.
- 63 Taitz J, Genn K, Brooks V, *et al*. System-wide learning from root cause analysis: a report from the New South Wales Root Cause Analysis Review Committee. *Qual Saf Health Care* 2010;19:e63.
- 64 Card AJ, Simsekler MC, Clark M, *et al*. Use of the Generating Options for Active Risk Control (GO-ARC) Technique can lead to more robust risk control options. *Int J Risk Saf Med* 2014;26:199–211.
- 65 Littlejohn A, Margaryan A, Lukic D. How organisations learn from safety incidents: a multifaceted problem. *J Workplace Learning* 2010;22:428–50.
- 66 Dixon-Woods M, Martin G, Tarrant C, *et al*. *Safer clinical systems: evaluation findings*. London: The Health Foundation, 2014.
- 67 Ward J, Clarkson J, Buckle P, *et al*. Prospective hazard analysis: tailoring prospective methods to a healthcare context. 2010; <http://www.webcitation.org/6KZ0Y4R8E> (accessed 27 Apr 2016).

The problem with root cause analysis

Mohammad Farhad Peerally, Susan Carr, Justin Waring and Mary Dixon-Woods

BMJ Qual Saf published online June 23, 2016

Updated information and services can be found at:

<http://qualitysafety.bmj.com/content/early/2016/06/23/bmjqs-2016-005511>

These include:

References

This article cites 47 articles, 11 of which you can access for free at: <http://qualitysafety.bmj.com/content/early/2016/06/23/bmjqs-2016-005511#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (266)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>