

Low-value care: an intractable global problem with no quick fix

John N Mafi,^{1,2} Michael Parchman³

¹Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

²RAND Health, RAND Corporation, Santa Monica, California, USA

³MacColl Center for Health Care Innovation, Kaiser Permanente Washington Health Research Institute, Seattle, Washington, USA

Correspondence to

Dr John N Mafi, Division of General Medicine and Health Services Research, Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, CA 90024, USA; jmafi@mednet.ucla.edu

Accepted 19 December 2017

Low-value care, or patient care that provides no net benefit in specific clinical scenarios, remains one of the most pressing problems in healthcare across the world—namely because it raises costs, causes iatrogenic patient harm, and often interferes with the delivery of high-value care. Many have argued that above all else the primary cause of low-value care lies in an unchecked fee-for-service payment system, which creates a pervasive culture that rewards providers for delivering more care, not necessarily the right care. Results reported by McAlister *et al* in this issue of *BMJ Quality & Safety* seem to up-end this belief.¹ In their analysis of 3.4 million beneficiaries in the globally-budgeted health system of Alberta, Canada, they found that low-value care commonly occurred—at a rate of approximately 5% of beneficiaries seeking care, and as high as 30% among those aged >75 years. Notably, these rates are comparable to rates in America's largely unrestrained fee-for-service system for both commercially insured (~8%) and older Medicare beneficiaries (~25-42%) seeking care, even while McAlister and colleagues used fewer low-value care measures (10) than the latter two American studies (28 and 26 respectively).^{2,3} Moreover, similar to the USA, the extent of the problem also varied substantially across frequently presumed examples of overuse. For instance, carotid artery imaging in adults without symptoms of cerebrovascular disease occurred in only 0.3% of patients, whereas 55.5% of men 75 years or older without a history of prostate cancer underwent prostate-specific antigen testing.

Although both Canadian and US physicians operate in fee-for-service payment models, Canadian physicians practice within a broader system of strict global budgets for hospitals and regional health authorities.⁴ Such financial restrictions

may reduce the overall volume of certain services: for instance, researchers found higher overall rates of CT utilisation in the USA compared with Canada.⁵ While global budgets may broadly reduce the overall volume of some (though not all) services, they provide too blunt an instrument to selectively reduce low-value care. In other words, all care may go down, not just low-value care.

This phenomenon of reducing both appropriate and inappropriate care has a long literature dating back to the RAND Health Insurance Experiment, which found similar rates of low-value care (but lower overall volume of services) in low-cost-sharing versus high-cost-sharing benefit plans, global-budgeted Health Maintenance Organizations (HMOs) compared with unrestricted fee-for-service providers.^{6,7} Similarly, other work found high rates of overuse among US safety-net physicians (usually practising within global budgets), as well as in globally-budgeted England's National Health Service (NHS), and equivalent rates of inappropriate coronary angiography in Canada and the USA.⁸⁻¹⁰ McAlister's study therefore reinforces an important lesson in health services research: while global budgets might be able to bluntly reduce the overall volume of some services, they are by themselves insufficient in changing a broader culture of medical practice that results in the delivery of low-value care, a theme we will return to later in this editorial. Low-value care is more complex than a simple financial incentive problem alone—and it remains globally pervasive and stubbornly intractable. Very few interventions have been shown to durably reduce it—and clearly as the authors argue, it is time to transition into a new era of experimentation and discovery of scalable interventions that reduce low-value care and recent research is beginning to point the way.



► <http://dx.doi.org/10.1136/bmjqs-2017-006778>
► <http://dx.doi.org/10.1136/bmjqs-2017-006699>



To cite: Mafi JN, Parchman M. *BMJ Qual Saf* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bmjqs-2017-007477

While technology such as information-based computerized clinical decision support has demonstrated limited results in reducing low-value care,¹¹ behavioural economics, or the field that posits that human beings (including physicians) predictably make irrational decisions due to known cognitive biases, has been heralded as an important new field of psychology and economics to apply to quality improvement. Low-cost and light-touch behavioural economic interventions (such as ‘nudges’)¹² hold obvious appeal. Early results show promise, with Patel and colleagues demonstrating the power of the default in computerised order entry and Meeker and colleagues publishing two rigorous, well-designed cluster-randomised controlled trials leveraging physicians’ intrinsic motivation to maintain a professional reputation and conform with peers in 2014 and 2016.^{13–16} While no panacea for fixing all aspects of healthcare delivery, these approaches showed how elegant and low-cost interventions such as displaying poster-sized commitment letters in physician exam rooms or prompting physicians to public accountable justification of low-value decisions can substantially reduce low-value antibiotic prescribing.

In this issue of *BMJ Quality & Safety*, Kullgren and colleagues continue this line of inquiry in an impressive stepwise wedge cluster randomised control trial evaluating a ‘light-touch’ behavioural economic intervention across several primary care practices in Michigan. They employed pre-commitment letters in an attempt to appeal to professionalism as an intrinsic motivator of clinicians’ behaviour.¹⁷ They trained medical assistants to prompt physicians with paper-based decision support and Choosing Wisely™ materials, sent weekly resources and Choosing Wisely™ materials to physicians, and used injunctive norms (a moral norm from an authoritative source strongly indicating how someone *ought* to behave)¹⁶ by appealing to professionalism. Despite these efforts, the intervention produced modest (at best) and unsustainable reduction in low-value care, along with an unintended increase in specialty referrals.

Why did they see no substantial reductions in low-value care while the study by Meeker and colleagues did? Not all interventions labelled with ‘behavioural economics’ are the same. For example, commitments used by Meeker *et al* were *public* and poster-sized, in the exam room, while Kullgren and colleagues used *private* signed letters. Public pre-commitments in public policy seem to have a richer literature and intuitively may be more likely to influence behaviour.¹⁵ Another important consideration is that Kullgren *et al* did not provide clinicians with suggested alternatives to providing low-value care. Meeker and colleagues offered decongestants as alternatives to unnecessary antibiotics, and the intervention by Patel and colleagues offered generic medications instead of brand name prescriptions. Such alternatives hold

intuitive merit because patients may be quite averse to losing something they expect to gain (eg, a prescription) from the doctor.

While offering an alternative may not always be necessary,¹⁵ it may aid clinicians in convincing patients to forgo low-value care—and importantly, lacking explicit alternatives in the study by Kullgren *et al* might have unintentionally led to rises in specialty care. That said, other possible explanations for the limited impact of this intervention include the attempt to reduce multiple services (Meeker and colleagues targeted just one service), and a far lower baseline rate of low-value care than did the study by Meeker *et al* (10% vs 43% of visits, and the lower the baseline, the harder it is to reduce it). Moreover, Kullgren and colleagues studied a ‘light touch’ pilot compared with more expensive initiatives such as computerized clinical decision support, which have also been unable to make large reductions in low-value care.¹¹ We look forward to future work building on their pilot study, and we congratulate them for identifying an important unintended consequence of increased use of specialty referrals, which may increase unnecessary spending.^{18 19}

So where do these studies leave us? Whether applying rational or behavioural economic models to behaviour, one stubborn yet often overlooked item that often comprise the magic sauce or the hidden cause for a behavioural intervention failing to go to scale is *culture*. Culture, that measurable yet unmeasurable ingredient, is one of major reasons we still have not (and almost certainly never will) discovered a fundamental law for human behaviour that can be applied in any context.²⁰ Yet, as it turns out, medical practice culture seems to matter. A recent mixed-methods quasi-experimental study found that prompting hospitals to foster a culture of diversity of engagement from all levels in quality improvement is associated with lower patient mortality.²¹ Other work by Gupta *et al* has found that cultural characteristics such as leadership messaging, data transparency and a blame-free environment are associated with higher-value care.²² While these are non-randomised studies, they underscore the importance of incorporating efforts to change the local culture of how medicine is practised within the context of where interventions are tested in order to move the needle on multiple fronts. Some insights about how to create conditions for change when tackling low-value care services can be found in the Taking Action on Overuse Framework.²³ More importantly, this framework suggests that the key to culture change may be ongoing sense-making conversations between clinician peers and team members about the potential for harm from overused services and deidentified current rates of use of these services. Howard Beckman’s work on engaging clinicians to reduce low-value care has been pioneering in this arena.²⁴ And while conversing with front-line clinicians, focusing on reducing iatrogenic

patient harm from low-value care seems to be a particularly favourable way to engage physicians.²⁵

Low-value care remains an intractable problem for a wide array of interrelated reasons, including clinician factors (eg, training, fear of lawsuit, time pressures, intolerance of uncertainty), patient factors (lack of knowledge or financial consequences) and healthcare system factors (institutional culture, pricing, fee-for-service payment models).^{26–39} Rather than implementing myopically top-down interventions (eg, mandating overly specific pay-for-performance policies), which may be virtually impossible to execute safely and effectively in complex, non-linear systems,^{40–42} we might instead propose an alternative strategy. It likely will require a combination of ‘light-touch’ top-down policies (eg, capitated payment arrangements that preserve clinician autonomy and access to care)^{43 44} as well as encouraging simultaneous bottom-up, pragmatic/trial-and-error-type local pilot initiatives that addresses multiple drivers of low-value care.²⁴ Starting with rigorous measurement of clinician performance,^{45 46} these interventions (ideally randomised) should simultaneously pull multiple levers of intrinsic motivation while also monitoring and adapting to unintended consequences⁴⁷ as Kullgren and colleagues astutely did. Not surprisingly, a recent meta-analysis found that multicomponent interventions are more effective than single-component interventions to reduce low-value care.³⁰ For example, Vivian Lee and colleagues demonstrated that top-level leadership on value improvement, fostering a culture of continuous improvement and providing clinicians with education and electronic cost transparency and patient outcome data, led to improvements in costs and quality and reductions in low-value daily inpatient labs.⁴⁸ While such pragmatic and multicomponent approaches may reduce national reproducibility, they may enhance local effectiveness and sustainability, particularly if they are shown to be cost-effective, seamlessly integrated into clinician workflow and free of iatrogenic harm (eg, unintentionally reducing necessary care). In an ideal world—one united in reducing harmful and unnecessary care—bottom-up, multicomponent initiatives are adaptively combined with education, ‘light-touch’ financial alignment, careful surveillance of unintended consequences and softer yet equally powerful cultural levers—all harmonising to finally tackle the problem of low-value care.

Correction notice This article has been updated since publication to correct a minor spelling mistake in the opening sentence.

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- McAlister FA, Lin M, Bakal J, *et al*. Frequency of low-value care in Alberta, Canada: a retrospective cohort study. *BMJ Qual Saf* 2017. doi: 10.1136/bmjqs-2017-006778. [Epub ahead of print 16 Sep 2017].
- Schwartz AL, Landon BE, Elshaug AG, *et al*. Measuring low-value care in Medicare. *JAMA Intern Med* 2014;174:1067–76.
- Reid RO, Rabideau B, Sood N. Low-value health care services in a commercially insured population. *JAMA Intern Med* 2016;176:1567–71.
- Commonwealth Fund Report. The canadian health care system. 2017 <http://international.commonwealthfund.org/countries/canada/> (accessed 19 Nov 2017).
- Berdahl CT, Vermeulen MJ, Larson DB, *et al*. Emergency department computed tomography utilization in the United States and Canada. *Ann Emerg Med* 2013;62:486–94.
- Newhouse JP, Manning WG, Morris CN, *et al*. Some interim results from a controlled trial of cost sharing in health insurance. *N Engl J Med* 1981;305:1501–7.
- Siu AL, Sonnenberg FA, Manning WG, *et al*. Inappropriate use of hospitals in a randomized trial of health insurance plans. *N Engl J Med* 1986;315:1259–66.
- McGlynn EA, Naylor CD, Anderson GM, *et al*. Comparison of the appropriateness of coronary angiography and coronary artery bypass graft surgery between Canada and New York State. *JAMA* 1994;272:934–40.
- Coronini-Cronberg S, Bixby H, Laverty AA, *et al*. English National Health Service's savings plan may have helped reduce the use of three 'low-value' procedures. *Health Aff* 2015;34:381–9.
- Barnett ML, Linder JA, Clark CR, *et al*. Low-value medical services in the safety-net population. *JAMA Intern Med* 2017;177:829–37.
- Goldzweig CL, Orshansky G, Paige NM, *et al*. Electronic health record-based interventions for improving appropriate diagnostic imaging: a systematic review and meta-analysis. *Ann Intern Med* 2015;162:557–65.
- Bourdeaux CP, Davies KJ, Thomas MJ, *et al*. Using 'nudge' principles for order set design: a before and after evaluation of an electronic prescribing template in critical care. *BMJ Qual Saf* 2014;23:382–8.
- Patel MS, Day S, Small DS, *et al*. Using default options within the electronic health record to increase the prescribing of generic-equivalent medications: a quasi-experimental study. *Ann Intern Med* 2014;161:S44–52.
- Patel MS, Day SC, Halpern SD, *et al*. Generic medication prescription rates after health system-wide redesign of default options within the electronic health record. *JAMA Intern Med* 2016;176:847–8.
- Meeker D, Knight TK, Friedberg MW, *et al*. Nudging guideline-concordant antibiotic prescribing: a randomized clinical trial. *JAMA Intern Med* 2014;174:425–31.
- Meeker D, Linder JA, Fox CR, *et al*. Effect of behavioral interventions on inappropriate antibiotic prescribing among primary care practices: a randomized clinical trial. *JAMA* 2016;315:562–70.
- Kullgren JT, Krupka E, Schachter A, *et al*. Precommitting to choose wisely about low-value services: a stepped wedge cluster randomised trial. *BMJ Qual Saf* 2017. doi: 10.1136/bmjqs-2017-006699. [Epub ahead of print 24 Oct 2017].
- Mafi JN, Edwards ST. How can we improve the efficiency of specialty care? *J Gen Intern Med* 2016;31:1267–9.

- 19 Mafi JN, Edwards ST, Pedersen NP, *et al.* Trends in the ambulatory management of headache: analysis of NAMCS and NHAMCS data 1999-2010. *J Gen Intern Med* 2015;30:548-55.
- 20 Morson GS, Schapiro M. *Cents and sensibility what economics can learn from the humanities*. Princeton, New Jersey: Princeton University Press, 2017.
- 21 Bradley EH, Brewster AL, McNatt Z, *et al.* How guiding coalitions promote positive culture change in hospitals: a longitudinal mixed methods interventional study. *BMJ Qual Saf* 2017. doi: 10.1136/bmjqs-2017-006574. [Epub ahead of print 3 Nov 2017].
- 22 Gupta R, Moriates C, Harrison JD, *et al.* Development of a high-value care culture survey: a modified Delphi process and psychometric evaluation. *BMJ Qual Saf* 2017;26:475-83.
- 23 Parchman ML, Henrikson NB, Blasi PR, *et al.* Taking action on overuse: creating the culture for change. *Health Care* 2017;5:199-203.
- 24 Cammisa C, Partridge G, Ardans C, *et al.* Engaging physicians in change: results of a safety net quality improvement program to reduce overuse. *Am J Med Qual* 2011;26:26-33.
- 25 Liao JM, Schapira MS, Navathe AS, *et al.* The effect of emphasizing patient, societal, and institutional harms of inappropriate antibiotic prescribing on physician support of financial penalties: a randomized trial. *Ann Intern Med* 2017;167:215-6.
- 26 Chen C, Petterson S, Phillips R, *et al.* Spending patterns in region of residency training and subsequent expenditures for care provided by practicing physicians for Medicare beneficiaries. *JAMA* 2014;312:2385-93.
- 27 Morgan DJ, Leppin AL, Smith CD, *et al.* A practical framework for understanding and reducing medical overuse: conceptualizing overuse through the patient-clinician interaction. *J Hosp Med* 2017;12:346-51.
- 28 Smith CD, Korenstein D. Harnessing the power of peer pressure to reduce health care waste and improve clinical outcomes. *Mayo Clin Proc* 2015;90:311-2.
- 29 Keyhani S, Falk R, Howell EA, *et al.* Overuse and systems of care: a systematic review. *Med Care* 2013;51:503-8.
- 30 Colla CH, Mainor AJ, Hargreaves C, *et al.* Interventions Aimed at Reducing Use of Low-Value Health Services: A Systematic Review. *Med Care Res Rev* 2017;74:507-50.
- 31 Colla CH, Morden NE, Sequist TD, *et al.* Choosing wisely: prevalence and correlates of low-value health care services in the United States. *J Gen Intern Med* 2015;30:221-8.
- 32 Colla CH. Swimming against the current--what might work to reduce low-value care? *N Engl J Med* 2014;371:1280-3.
- 33 Jena AB, Schoemaker L, Bhattacharya J, *et al.* Physician spending and subsequent risk of malpractice claims: observational study. *BMJ* 2015;351:h5516.
- 34 Gidengil CA, Mehrotra A, Beach S, *et al.* What drives variation in antibiotic prescribing for acute respiratory infections? *J Gen Intern Med* 2016;31:918-24.
- 35 Allison JJ, Kiefe CI, Cook EF, *et al.* The association of physician attitudes about uncertainty and risk taking with resource use in a Medicare HMO. *Med Decis Making* 1998;18:320-9.
- 36 Jha S. Uncertainty and the diagnostic leviathan. *JAMA Intern Med* 2015;175:1085-6.
- 37 Shah BR, Cowper PA, O'Brien SM, *et al.* Association between physician billing and cardiac stress testing patterns following coronary revascularization. *JAMA* 2011;306:1993-2000.
- 38 Mafi JN, Wee CC, Davis RB, *et al.* Association of primary care practice location and ownership with the provision of low-value care in the United States. *JAMA Intern Med* 2017;177:838.
- 39 Mafi JN, Russell K, Bortz BA, *et al.* Low-cost, high-volume health services contribute the most to unnecessary health spending. *Health Aff* 2017;36:1701-4.
- 40 Taleb NN. *The Black Swan: the impact of the highly improbable* New York. New York: Random House Books, 2007.
- 41 Mendelson A, Kondo K, Damberg C, *et al.* The effects of pay-for-performance programs on health, health care use, and processes of care: a systematic review. *Ann Intern Med* 2017;166:341-53.
- 42 Kontopantelis E, Springate DA, Ashworth M, *et al.* Investigating the relationship between quality of primary care and premature mortality in England: a spatial whole-population study. *BMJ* 2015;350:h904.
- 43 Schwartz AL, Chernew ME, Landon BE, *et al.* Changes in low-value services in year 1 of the medicare pioneer accountable care organization program. *JAMA Intern Med* 2015;175:1815-25.
- 44 Song Z, Rose S, Safran DG, *et al.* Changes in health care spending and quality 4 years into global payment. *N Engl J Med* 2014;371:1704-14.
- 45 Friedberg MW, Damberg CL. A five-point checklist to help performance reports incentivize improvement and effectively guide patients. *Health Aff* 2012;31:612-8.
- 46 Saini SD, Powell AA, Dominitz JA, *et al.* Developing and testing an electronic measure of screening colonoscopy overuse in a large integrated healthcare system. *J Gen Intern Med* 2016;31(Suppl 1):53-60.
- 47 Kerr EA, Kullgren JT, Saini SD. Choosing wisely: how to fulfill the promise in the next 5 years. *Health Aff* 2017;36:2012-8.
- 48 Lee VS, Kawamoto K, Hess R, *et al.* Implementation of a value-driven outcomes program to identify high variability in clinical costs and outcomes and association with reduced cost and improved quality. *JAMA* 2016;316:1061-72.

**BMJ Quality
& Safety**

Low-value care: an intractable global problem with no quick fix

John N Mafi and Michael Parchman

BMJ Qual Saf published online January 13, 2018

Updated information and services can be found at:

<http://qualitysafety.bmj.com/content/early/2018/01/19/bmjqs-2017-007477>

These include:

References

This article cites 42 articles, 7 of which you can access for free at:
<http://qualitysafety.bmj.com/content/early/2018/01/19/bmjqs-2017-007477#ref-list-1>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>