Cardiac rehabilitation

Jacqueline Dinnes, Jos Kleijnen, Maria Leitner, David Thompson

Cardiac rehabilitation services aim to facilitate physical, psychological, and emotional recovery in patients with heart disease (table 1) and enable them to achieve and maintain better health through exercise, education, help with psychological sequelae, or any combination of these elements.

Drug treatment is an integral part of the secondary prevention and treatment of coronary heart disease; existing systematic reviews have already shown the effectiveness of drugs such as the statins, aspirin, angiotensin converting enzyme (ACE) inhibitors, and β blockers.¹⁻⁴ This review is therefore focused specifically on rehabilitation interventions as opposed to drug treatments.

Current provision in the United Kingdom

The overall level of provision in the United Kingdom has increased rapidly over the past 10-15 years⁵⁻¹⁰ and there are currently almost 300 cardiac rehabilitation programmes in existence.¹¹ The total cost of cardiac rehabilitation is estimated to be up to £34m each year.¹² There is wide variation in practice and in the organisation and management of services; evidence exists that current service provision fails to meet the standards set in national guidelines for cardiac rehabilitation,^{13 14} and that secondary prevention measures are under applied.^{15 16} Disparities in provision are reflected in the resulting costs incurred. Annual staffing costs across 16 centres have been shown to range from £10 000 to £62 000; with a median cost per patient enrolled of £223.¹⁷

Most programmes are outpatient, hospital based,^{8 9 13} concentrating on low risk patients with myocardial infarction (MI), although many also include patients who have had coronary artery bypass surgery or angioplasty.^{8 13} Women are less likely to receive cardiac rehabilitation than men; a recent survey of 244 cardiac rehabilitation programmes in the UK found that only 15% of enrollees were

women,¹³ although they account for over one third of patients with coronary heart disease.

A recent survey of patients with coronary heart disease found that the majority were not receiving appropriate medication for their condition; were not undertaking regular exercise; were overweight; and were not eating an appropriate diet.¹⁶ Up to 90% of these patients would have benefited from further changes in lifestyle, and only 7% were receiving optimal medical management for prevention of heart disease.¹⁶

Recovery after an acute cardiac event

The needs of people recovering from an acute cardiac event vary. Some have psychological problems or misconceptions about their condition which may make it difficult for them to return to a normal life. Some require help in modifying pre-existing risk factors such as smoking, poor diet, or lack of exercise. Most are likely to benefit from lifestyle changes such as increasing physical activity.

Although 12 weeks after MI up to 30% of patients report that their quality of life has returned to previous levels,¹⁸ symptoms of anxiety and depression are common and have been shown to be associated with prolonged disability, re-infarction, and death.¹⁹⁻²³

Perceived health status, level of misconception about the heart condition, and anxiety and depression are the major predictors of return to normal activity.²⁴ Many patients who have suffered a MI fear and avoid activity²⁵⁻²⁹; up to 50% report reduced social and leisure activities four years later.²⁹ Return to work rates are fairly high, however a substantial number of patients retire early or become unemployed.²⁹⁻³¹

Methods

This paper is based on existing systematic reviews of acceptable quality, supplemented with reference to key randomised controlled trials (RCTs). In those areas where there were few RCTs, other study designs were also included. Studies were identified by a search of computerised databases, including Medline, Embase, CINAHL, and PsycLit, using keywords such as cardiac rehabilitation, rehabilitation, MI, angioplasty, and coronary artery bypass (a copy of the full search strategy is available on request). The computerised searches were supplemented with a review of the reference lists of retrieved papers and contact with experts. All retrieved studies were assessed for inclusion independently by two reviewers and any discrepancies discussed.

NHS Centre for Reviews and Dissemination, University of York, UK

Jacqueline Dinnes, research fellow
Jos Kleijnen, director
Maria Leitner, senior research fellow

Department of Health Studies, University of York, UK

David R Thompson, professor

Correspondence to:
Jacqueline Dinnes, NHS Centre for Reviews and Dissemination, University of York, York, YO10 5DD, UK.

Accepted 28 January 1999

Table 1 Cardiac rehabilitation services for patients with heart disease

Categories of patients who might benefit from cardiac rehabilitation	Estimated number each year in the UK ¹
Myocardial infarction	150 000
Coronary artery bypass graft	22 056
Coronary angioplasty	13 822
Angina	1 400 000
Heart failure	500 000
Heart transplantation	
Heart only	258
Heart and lung	46
Total	2 086 182

Table 2 Systematic reviews of cardiac rehabilitation

Authors	Objective	Search strategy	Inclusion criteria for trials	Results	Notes
Bobbio, 1989 ³³	To quantitatively evaluate the effect of post myocardial infarction rehabilitation on total mortality, cardiac mortality, and recurrence of non-fatal MI	Electronic search of Medline 1980–86 (keywords given); bibliographies of text books; trials analysed in 5 review articles; papers presented at previous 2 conferences of the World Congress on Cardiac Rehabilitation; reference sections of all articles retrieved	Required: controlled or randomised controlled trials with at least 2 years follow up; reporting of total death, cardiac death, non-fatal MIs; intention to treat analysis; publication in peer reviewed journals	8 RCTs (2260 patients) Pooled relative risk Total mortality: 0.68 (95% CI, 0.53 to 0.86, p=0.02) Cardiac mortality: 0.62 (95% CI, 0.48 to 0.82 p=0.001) Non-fatal MI: 1.12 (95% CI, 0.84 to 1.49 p=0.45)	Meta-analysis, limited search strategy, independent, blinded data extraction, relatively detailed description of methodology used but method used to pool data not completely clear, non-significant test for heterogeneity, unclear why relative risk used rather than odds ratio
Bucher, 1994 ¹⁰⁵	To estimate the effect of social support on prognosis after a first MI	Electronic search of Medline (keywords provided)	Required: an inception cohort with complete report of follow up; objective outcomes (for example, re-infarction, mortality); adequate adjustment for confounding factors; intervention studies required adequate randomisation and reporting of all “relevant clinical outcomes”	7/9 cohort studies found association between lack of social support and increased mortality 2 RCTs of increased support from health professionals appear to show reduced mortality/morbidity, but methodologically flawed	Narrative review with limited search and no stated validity assessment. Only 4/7 cohort studies used detailed measures of social support, other three used “living alone v living with someone”
Duryee, 1992 ⁴⁰	To establish whether inpatient education increases patients with MIs’ knowledge and produces lifestyle change	Electronic search of Medline 1975–89 (no keywords provided); search of “reference lists”	Required: post MI and/or cardiac surgery patients; clear description of research design; data collection using an “objective instrument”; presentation of data	10/13 studies (9 controlled trials) showed increased knowledge 6/8 studies (7 controlled trials) showed some lifestyle change especially for increasing activity and smoking cessation	Narrative review, limited search, limited study detail provided and little comment on quality of studies, precise measure of knowledge of behaviour change varies between studies
Hill <i>et al</i> , 1992 ³⁹	To determine whether psychosocial interventions can minimise psychological distress and psychiatric morbidity in coronary heart disease and cancer patients	Electronic searches of Medline and PsycLit (keywords provided); searches of “individual issues of relevant journals” in the fields of psychiatry, nursing, psychology and social work “for the past five years”; citations from experts, from reviews and from government documents	Required: replicable global psycho-social intervention; standardised mental health outcome; control or comparison group Excluded: non-cardiac, non-cancer patients; focus on Type A behaviour in cardiac patients; hypnosis based or pharmacological interventions	Myocardial infarction: 6/7 studies showed general improvements in distress (mainly anxiety or depression). 2 were reports of same study (male patients and spouses), 5/6 were RCTs CABS**: 4/5 studies suggest reductions in self-reported anxiety. 3/5 were RCTs	Narrative review with some methodological detail given, however sample sizes in all studies were small, and not enough study details were provided to judge their quality. No detail of author’s quality assessment provided.
Linden <i>et al</i> , 1996 ³⁷	To quantitatively review the evidence for the efficacy of psychosocial interventions for patients with coronary artery disease	Electronic search of Medline (no keywords or years given); references lists of retrieved papers and review articles	Required: documented coronary artery disease; at least one control group; the evaluation of the additional impact of a psychosocial intervention over usual care; randomisation	23 RCTs (3180 patients) Pooled odds ratios at ≤2 years follow up Total mortality (10 RCTs): 1.70 (95% CI, 1.09 to 2.64, p=.02) Non-fatal MI (8 RCTs): 1.84 (95% CI, 1.12 to 2.99, p=.02) Pooled odds ratios at >2 years follow-up Total mortality (3 RCTs): 1.35 (95% CI, 0.83 to 1.53, p=.13) Non-fatal MI (3 RCTs) 1.64 (95% CI, 1.06 to 2.54, p=.02) Psychosocial interventions also produced greater reductions in psychosocial distress, systolic blood pressure, heart rate, cholesterol concentrations)	Meta-analysis, very limited literature search, no detail regarding validity or quality assessment, limited study details provided, duplicate publications included in the analysis
Mullen <i>et al</i> , 1992 ³⁸	To quantitatively assess the effects of patient education on cardiac morbidity and mortality	Electronic search of several databases (years and terms provided); bibliographies of retrieved studies; database of the National Heart, Lung and Blood Institute and the Veterans Administration Health Services Research and Development Section	Required: English language; published and unpublished reports; evaluation of a psychosocial or educational intervention with adult patients diagnosed with coronary artery disease; sample size of ≥10 per arm; randomised, quasi-experimental comparison group design or a one group pre-test post-test design	28 CTs* (4512 patients) Weighted average effect sizes: Exercise (12 CTs); 0.18 (95% CI, 0.07 to 0.29) Diet (9 CTs); 0.19 (95% CI, 0.05 to 0.34) Smoking (9 CTs); 0.07 (95% CI, -0.08 to 0.22) Stress (0 CTs) Drug adherence (3 CTs); -0.09 (95% CI, -0.39 to 0.22) Morbidity (9 CTs); 0.05 (95% CI, -0.04 to 0.13) Return to work (6 CTs); 0.08 (95% CI, -0.11 to 0.27) Death (7 CTs); 0.24 (95% CI, 0.14 to 0.33) Blood pressure (5 CTs) 0.51 (95% CI, 0.24 to 0.77)	Meta-analysis—substantial literature search, good description of methodology; blinded validity assessment using Sackett and Haynes coding scheme; considerable diversity between studies Data pooling: WAESs calculated for homogenous groups of trials—that is, with any outliers removed. Descriptions of outliers and reasons for removal given in text, but could introduce significant bias Effects difficult to interpret as the basis on which the average effect sizes were calculated for each outcome measure is unknown

Table 2 Continued

Authors	Objective	Search strategy	Inclusion criteria for trials	Results	Notes
O'Connor <i>et al</i> , 1989 ³²	To quantitatively determine whether cardiac rehabilitation has a significant effect on total and cardiovascular mortality, sudden death, and fatal or non-fatal re-infarction in post-MI patients	Formal computer aided searches; informal search for studies known to the research group	Required: patients to be randomised individually; post-MI only; follow up over 1 year; intervention to include a structured exercise component; published data	22 RCTs (4554 patients) Pooled odds ratios at >36 months FU Total mortality: 0.80 (95% CI, 0.66 to 0.96) Cardiovascular mortality: 0.78 (95% CI, 0.63 to 0.96) Sudden death: 0.92 (95% CI, 0.69 to 1.23) Fatal MI: 0.75 (95% CI, 0.59 to 0.95) Non-fatal MI: 1.09 (95% CI, 0.88 to 1.34)	Meta-analysis, difficult to evaluate due to limited description of methodology, although data pooling appears to have been adequate NB 14/22 RCTs are part of one large WHO trial, not referenced individually in other meta-analyses
Oldridge <i>et al</i> , 1988 ³¹	To quantitatively examine the effect of cardiac rehabilitation after MI on total mortality, cardiac mortality, and recurrent MI	Electronic search of Medline search (keywords provided); review of relevant English language publications; suggestions from colleagues for published and unpublished data	Required: RCTs only; adequate randomisation; MI only; rehabilitation with exercise for at least 6 weeks; follow up at least 24 months; outcome had to include at least two of all causes of death, cardiovascular mortality or non-fatal MI	10 RCTs (4347 patients) Pooled odds ratios All cause death: 0.76 (95% CI, 0.63 to 0.92 p=.004) Cardiovascular mortality 0.75 (95% CI, 0.62 to 0.93 p=.006) Non-fatal MI 1.15 (95% CI, 0.93 to 1.42)	Meta-analysis, limited search, blinded quality assessment of trials, good description of data pooling methods, probably used intention-to-treat analysis of individual studies
Wenger <i>et al</i> , 1995 ³³	To evaluate the scientific evidence pertaining to the various components of cardiac rehabilitation	Electronic search of Medlars (Grateful Med, Psych, ERIC, CINAHL) and NLM database; searches in the psychosocial and professional nursing literature; citations from experts	Required: any study design; any manifestations of coronary heart disease; any aspect of cardiac rehabilitation including exercise training, education and counselling, psychosocial and behavioural interventions, medical therapies, surveillance; and transition planning or adherence	Exercise alone: improves exercise tolerance and pathophysiological outcomes; no significant effect on morbidity or mortality; effect on risk factors, psychosocial outcomes, and return to work is unclear, but may help improve symptoms. Did not evaluate individual effect of psychosocial interventions Multifactorial programmes are required in order to achieve beneficial effects on risk factors, psychosocial outcomes, morbidity and mortality	Narrative review of vast amount of literature, methodology unclear especially regarding quality assessment and data extraction; conclusions don't always accord with the evidence presented. Combination of single modality and multifactorial rehabilitation studies used to evaluate outcomes in each section, therefore no true picture of effects of different interventions is given. Study details provided in technical report

* CT=controlled trials; **CABS=coronary artery bypass surgery.

Nine systematic reviews (table 2) and 28 separate trials met the inclusion criteria. Subsequently, two reviewers extracted data on the results following the same procedure.

Results

Overall the quality of the literature in this field is poor and its evaluation difficult due to the variability of interventions and patient populations studied, methodological problems, and poor quality reporting. There is little reporting of randomisation procedures, little detail of the interventions provided, and study sample sizes are often small. In addition, the use of care "packages" complicates the evaluation of individual interventions as it is difficult to identify the impact of the specific components. The majority of studies include only low risk, male, white, middle aged patients with MI, and exclude or enrol only a small number of women, the elderly, ethnic minorities, and other cardiac patient groups such as those having had cardiac surgery, heart failure, or heart transplantation, thereby limiting the generalisability of the results. Table 2 summarises the systematic reviews on which some of the key conclusions of this paper are based, and further details are available elsewhere.³²

Exercise

Exercise alone has a positive impact on patients' physical ability to exercise, and on physiological measures of cardiac disease. It

does not have any effect on smoking cessation or reduction of cholesterol concentrations. Not enough evidence exists to evaluate its effect on body weight or blood pressure.³³

Exercise alone had no significant effect on morbidity (usually evaluated by non-fatal re-infarctions) or overall mortality rates, however a trend towards a beneficial effect on symptoms (angina) has been shown. It should be emphasised that exercise has not been found to do any harm to patients. Not enough evidence was found to evaluate the effect of exercise alone on psychological or social outcomes, or return to work.³³ These conclusions have largely been supported by additional identified trials.³⁴⁻³⁶

Psychological and educational interventions

Psychosocial interventions include patient education, counselling, and behavioural interventions. These have been shown to affect risk factors including blood pressure and cholesterol concentrations,³⁷⁻³⁸ and to produce significant improvements in psychosocial wellbeing³⁷⁻³⁹ and in patient knowledge.⁴⁰

Psychosocial interventions may significantly reduce morbidity and mortality in patients with coronary artery disease.³⁷⁻³⁸ It has been estimated that psychosocial interventions could produce a 46% reduction in non-fatal cardiac events and a 41% reduction in mortality at two

years follow up. The effect on mortality was not found to be significant after two years (although patient numbers were small).³⁷ However, this study was methodologically flawed (table 2), and its results should therefore be treated with some caution.

Additional RCTs largely support these conclusions,^{41–46} although two studies with particularly large sample sizes found no effects for mortality or morbidity, nor indeed for measures of psychological morbidity.^{42–45} However, the interventions used were mainly targeted at reducing patients' stress levels. In one trial this was done by monthly telephone contact with further interventions where necessary, which would not meet any generally accepted definition of cardiac rehabilitation.⁴² The other trial, aiming at psychological improvements, had seven weekly sessions with a psychologist or health visitor and did not include interventions aimed at risk factor modification.⁴⁵

Increases in knowledge may not necessarily be sufficient to produce changes in behaviour or lifestyle, but inpatient education has been shown to produce significant improvements in smoking behaviour, activity levels, and overall compliance with action to improve health.⁴⁰ In one RCT over half the patients were shown to be following advice one year after an acute event.⁴⁷ There is a widespread misconception among the general public and some health professionals that people should considerably limit their activity after MI to avoid a recurrence.^{48–49}

Combined exercise and psychological or educational interventions

When exercise is combined with patient education and counselling, it may improve cardiac risk factors, particularly reduced lipids and blood pressure. An intensive approach with specific anti-smoking advice may also help to improve smoking cessation rates.³³ Multifactorial rehabilitation can have an impact on exercise levels after the rehabilitation programme, at least in the short term.³³ Its effect on psychological wellbeing is not clear, but it has not been shown to have an effect on return to work or on angina.³³

Only one RCT of multifactorial rehabilitation has shown a clear reduction in mortality among patients who have had a MI.⁵⁰ Other studies are too small to show any significant impact on morbidity or mortality.³³ Data from three published meta-analyses,^{51–53} two of which were of high quality,^{51–52} suggest a reduction in cardiac mortality of about 20–25% (table 2). No significant effect on non-fatal re-infarctions was found.^{51–53} However, since the studies in these meta-analyses focused on patients at low risk of recurrence, the scope for demonstrating benefit is small. It is likely that the benefits of appropriate cardiac rehabilitation would be greater in patients with more severe cardiac disease, but caution should be used when generalising these results to other populations. These conclusions have been largely supported by additional identified trials.^{54–56}

TARGET POPULATION

Current provision of the service tends to concentrate on low risk, white, male, post-MI patients. No evidence exists of lack of benefit for other groups, such as women, the elderly, ethnic minorities, and patients with other types of heart disease.

There is some evidence that when women or the elderly attend cardiac rehabilitation programmes, the outcomes are as good as or better than for men^{57–61} and for younger groups of patients.^{62–65} Ethnic minority groups may have a greater risk of re-infarction or recurrence due to lower levels of activity⁶⁶ and higher levels of morbidity.⁶⁷ Observational studies in the United States have shown no significant racial differences in response to cardiac rehabilitation.^{68–69} The involvement of partners and other close family members in the rehabilitation process may improve patient outcomes.^{6–70–71}

A few small RCTs of patients with heart failure suggest improvements in physical aspects of the disease similar to those seen in patients after MI.³³ It is possible that increasing fitness and physical ability in patients with heart failure^{72–75} or angina^{76–78} may provide worthwhile benefits including reduced symptoms, reduced disability, and improved quality of life. No RCTs have been identified which evaluated patients after heart transplantation.³³

Organisation of services

FREQUENCY AND DURATION

Little research has evaluated the frequency and duration of cardiac rehabilitation programmes. Most studies provided three to five supervised exercise sessions each week, in combination with a patient education or psychosocial intervention, for approximately 12 weeks.³³

Significant improvements in lifestyle, symptoms, health status, and hospital readmission rates over two years have been achieved by interventions designed to initiate and maintain lifestyle change in patients with established coronary heart disease or angina. Two RCTs have shown that personal health education or visits to a secondary prevention clinic every two to six months can be more effective than routine care from general practitioners.^{79–80}

LOCATION

Although most cardiac rehabilitation programmes are conducted in a hospital outpatient setting, several studies have examined home based programmes. Studies comparing home exercise programmes with hospital based ones have shown improved cardiovascular fitness in both settings, with no increased risk of cardiac arrest in the home based programme.^{81–82}

RCTs comparing the effects of supervised home exercise with a no exercise control group^{83–87} have shown greater improvement in the rehabilitation group, in either risk factors,⁸⁵ anxiety and quality of life,^{83–84} or ability to exercise.^{85–87} Not enough studies are available to allow unsupervised home based exercise programmes to be evaluated.^{88–90} Telephone

based educational interventions with no prescribed rehabilitation programme are not effective.^{42 47 91}

Access, uptake, and adherence

Reported rates of uptake of cardiac rehabilitation range from 15%⁹² to 59%.⁹³ Approximately 20–25% of patients drop out of exercise programmes within the first three months and about 40–50% at between six and 12 months.⁹⁴

Poor uptake rates relate mainly to either service factors, such as the invitation to participate, or availability of services or patient factors. Elderly^{57 93} and female^{57 93 95} patients are significantly less likely to be invited to attend cardiac rehabilitation programmes although it is not clear from these studies why this should be the case. Patients receiving acute treatment at specific hospitals or from cardiologists are more likely to be invited to cardiac rehabilitation programmes than patients treated at other, similar hospitals or by general physicians.⁹³ The patients' perception of the strength of a physician's recommendation to attend^{57 96} and the availability and accessibility of the rehabilitation programme⁹⁷ are among the strongest predictors of whether patients attend once invited.

Uptake following invitation is much lower in women^{58 95} and in the elderly.^{58 98} This may be connected to personal beliefs either that cardiac rehabilitation is inappropriate for them or fears that they will feel out of place.^{95 99} Other sociodemographic characteristics including deprivation,⁹³ level of education,⁹⁸ and spouse involvement¹⁰⁰ are also significant predictors of uptake. Other reasons for not participating have included feelings that the wrong information or inadequate information is given or lack of motivation.^{95 96 101}

Dropouts from exercise programmes may occur more frequently in high intensity and poorly organised programmes, and among smokers and patients who have had more than one MI.^{35 94 102} Convenience of access to facilities also influences participation.^{97 98} Women are more likely than men to drop out.^{95 103}

Cost effectiveness

An American economic evaluation¹⁰⁴ has shown that under the assumption that cardiac rehabilitation produces other savings to the health service, the cost for each life year gained over three years from cardiac rehabilitation was \$21 800, or \$35 900 with no allowance for savings. The cost for each quality adjusted life year (QALY) was estimated to be \$6800.¹⁰⁴ In 1997, these results were recalculated to better reflect the situation in the United Kingdom. The results suggest a cost for each QALY of £6900, and a cost for each life year gained at three years of £15 700.¹²

Although it has been concluded that cardiac rehabilitation is cost effective,¹⁰⁴ it is clearly not a homogenous service and there are a range of factors that influence the costs and cost effectiveness of the process including: scale of the programme, location, components and intensity of the process, the patient population, and compliance. Cardiac rehabilitation may

not be cost effective in all formats for all patients. It may be more effective and therefore cost effective to provide specific interventions only to those patients who have a need for them, but this would require more adequate assessment of individual patients' needs.

Conclusions

Current service provision concentrates on low risk, white, male, middle aged post MI patients, however no evidence exists that other groups such as women, the elderly, ethnic minorities, or high risk cardiac patients do not benefit. There appears to be no basis for the exclusion of these groups from cardiac rehabilitation programmes and more research is required to identify reasons for, and strategies to improve, the current low levels of uptake and to design safe and effective programmes to meet the needs of different patient groups.

Although some benefits from single modality interventions have been shown, a combined approach of exercise and psychological and educational interventions appears to be more beneficial. Exercise has a positive impact on physical aspects of recovery at no additional risk to the patient, but no effect on risk factors, morbidity, or mortality. The effect of exercise on the psychosocial aspects of recovery are unclear. Trials have established that psychological and educational interventions can reduce risk factors, improve psychosocial well-being, and patient knowledge and may reduce morbidity and mortality. In practice, however, the information provided is often inadequate, inconsistent, and inaccurate and is frequently misunderstood by patients. The public and health professionals need to understand that activity should not be significantly reduced after a cardiac event.

Important questions remain to be answered as to the optimal mix of components and the frequency and duration of the programme. Home based rehabilitation may be as effective and safe as hospital based programmes when a prescribed rehabilitation programme and some form of supervision is provided, but further evaluation is required. Given the need to maintain improvements in lifestyle and the problems with continuing compliance, there is an existing need for long term maintenance programmes.

Further research is required to identify the optimal method of delivering the service. In particular, it is important to compare the clinical and cost benefits of menu driven systems, and home based and community based services with current hospital based programmes.

This paper is based on *Effective Health Care* volume 4, no 4 which is a systematic review of the evidence on the effectiveness of cardiac rehabilitation.

- 1 British Heart Foundation. *Coronary heart disease statistics*. London: British Heart Foundation, 1997.
- 2 NHS Centre for Reviews and Dissemination. Management of stable angina. *Effective Health Care* 1997;3.
- 3 NHS Centre for Reviews and Dissemination. Cholesterol and coronary heart disease: screening and treatment. *Effective Health Care* 1998;4.
- 4 Antman EM, Lau J, Kupelnick B, et al. A comparison of results of meta-analyses of randomised control trials and recommendations of clinical experts: treatments for myocardial infarction. *JAMA* 1992;268:240–8.

- 5 Green V, Stansfield BJ, Davidson C. Cardiac rehabilitation in the United Kingdom 1985/86: a questionnaire survey. *Physiotherapy* 1988;74:363-5.
- 6 Horgan J, Bethell H, Carson P, et al. Working party report on cardiac rehabilitation. *Br Heart J* 1992;67:412-8.
- 7 Bradley JM, Wallace ES, McCoy PM, et al. A survey of exercise based cardiac rehabilitation services in Northern Ireland. *Ulster Medical Journal* 1997;66:100-6.
- 8 Campbell NC, Grimshaw J, Rawles JM, et al. Cardiac rehabilitation in Scotland: is current provision satisfactory? *Journal of Public Health Medicine* 1996;18:478-80.
- 9 Davidson C, Reval K, Chamberlain DA, et al. A report of a working group of the British Cardiac Society: cardiac rehabilitation services in the United Kingdom 1992. *Br Heart J* 1995;73:201-2.
- 10 Lewin RJ, Ingleton R, Newens AJ, et al. Adherence to cardiac rehabilitation guidelines: a survey of cardiac rehabilitation programmes in the United Kingdom. *BMJ* 1998;316:1354-5.
- 11 Turner S. Personal communication: British Association for Cardiac Rehabilitation, 1998.
- 12 Taylor R, Kirby B. The evidence base for the cost-effectiveness of cardiac rehabilitation. *Heart* 1997;78:5-6.
- 13 Thompson DR, Bowman GS, Kitson AL. Cardiac rehabilitation services in England and Wales: a national survey. *International Journal of Cardiology* 1997;59:299-304.
- 14 Thompson DR, Bowman GS, De Bono DP, et al. *Cardiac rehabilitation: guidelines and audit standards*. London: Royal College of Physicians, 1997.
- 15 ASPIRE Steering Group. A British Cardiac Society survey of the potential for the secondary prevention of coronary heart disease: ASPIRE (action on secondary prevention through intervention to reduce events) principal results. *Heart* 1996;75:334-42.
- 16 Campbell NC, Thain J, Deans HG, et al. Secondary prevention in coronary heart disease: baseline survey of provision in general practice. *BMJ* 1998;316:1430-4.
- 17 Gray AM, Bowman GS, Thompson DR. The cost of cardiac rehabilitation services in England and Wales. *Journal of the Royal College of Physicians of London* 1997;31:57-61.
- 18 Laerum E, Johnsen N, Smith P, et al. Myocardial infarction may induce positive changes in lifestyle and in the quality of life. *Scand J Prim Health Care* 1988;6:67-71.
- 19 Barefoot JC, Schroll M. Symptoms of depression, acute myocardial infarction, and total mortality in a community sample. *Circulation* 1996;93:1976-80.
- 20 Frasure-Smith N, Lesperance F, Talajic M. Depression following myocardial infarction. *JAMA* 1993;27:1819-25.
- 21 Frasure-Smith N, Lesperance F, Talajic M. Depression and 18-month prognosis after myocardial infarction. *Circulation* 1995;91:999-1005.
- 22 Ladwig KH, Roll G, Breithardt G, et al. Post-infarction depression and incomplete recovery 6 months after acute myocardial infarction. *Lancet* 1994;343:20-3.
- 23 Ruberman W, Weinblatt E, Goldberg JD, et al. Psychosocial influences on mortality and myocardial infarction. *New Engl J Med* 1984;311:552-9.
- 24 Maeland JG, Havik OE. After the myocardial infarction. A medical and psychological study with special emphasis on perceived illness. *Scandinavian Journal of Rehabilitation Medicine: supplement* 1989;22:1-87.
- 25 Croog SH. Recovery and rehabilitation of heart patients: psychosocial aspects. In: Krantz DS, Singer JS, editors. *Handbook of psychology and health*. Hillsdale: Lawrence Erlbaum Associates, 1983:295-334.
- 26 Mallaghan M, Pemberton J. Some behavioural changes in 493 patients after an acute myocardial infarction. *Br J Prev Soc Med* 1977;163:285-92.
- 27 Wiklund I, Sanne H, Vedin A, et al. Psychosocial outcome one year after a first myocardial infarction. *J Psychosom Res* 1984;28:309-21.
- 28 Wiklund I, Sanne H, Elmfeldt D, et al. Emotional reaction, health preoccupation and sexual activity two months after a myocardial infarction. *Scand J Rehabil Med* 1984;16:47-56.
- 29 Finlayson A, McEwen J. *Coronary heart disease and patterns of living*. London: Croom Helm, 1977.
- 30 Cay EL, Vetter N, Philip A, et al. Psychological status during recovery from an acute heart attack. *J Psychosom Res* 1972;16:425-35.
- 31 Mayou R. The course and determinants of reactions to myocardial infarction. *Br J Psychiatry* 1979;134:588-94.
- 32 NHS Centre for Reviews and Dissemination. Cardiac rehabilitation. *Effective Health Care* 1998;4.
- 33 Wenger NK, Froelicher ES, Smith LK, et al. *Cardiac rehabilitation as secondary prevention*. Rockville, Maryland: Agency for Health Care Policy and Research and National Heart, Lung, and Blood Institute, 1995.
- 34 Bethell HJ, Mullee MA. A controlled trial of community based coronary rehabilitation. *Br Heart J* 1990;64:370-5.
- 35 Lee JY, Jensen BE, Oberman A, et al. Adherence in the training levels comparison trial. *Medicine and Science in Sports and Exercise* 1996;28:47-52.
- 36 Specchia G, De Servi S, Scire A, et al. Interaction between exercise training and ejection fraction in predicting prognosis after a first myocardial infarction. *Circulation* 1996;94:978-82.
- 37 Linden W, Stossel C, Maurice J. Psychosocial interventions for patients with coronary artery disease: a meta-analysis. *Arch Intern Med* 1996;156:745-52.
- 38 Mullen PD, Mains DA, Velez R. A meta-analysis of controlled trials of cardiac patient education. *Patient Education and Counseling* 1992;19:143-62.
- 39 Hill DR, Kelleher K, Shumaker SA. Psychosocial interventions in adult patients with coronary heart disease and cancer. A literature review. *General Hospital Psychiatry* 1992;14:28-42S.
- 40 Duryee R. The efficacy of inpatient education after myocardial infarction. *Heart and Lung* 1992;21:217-25.
- 41 Burell G. Behaviour modification after coronary artery bypass graft surgery: effects on cardiac morbidity and mortality. *J Rehabil Sci* 1995;8:39-40.
- 42 Frasure-Smith N, Lesperance F, Prince RH, et al. Randomised trial of home-based psychosocial nursing intervention for patients recovering from myocardial infarction. *Lancet* 1997;350:473-9.
- 43 Heller RF, Knapp JC, Valenti LA, et al. Secondary prevention after acute myocardial infarction. *Am J Cardiol* 1993;72:759-62.
- 44 Taylor CB, Houston-Miller N, Killen JD, et al. Smoking cessation after acute myocardial infarction: effects of a nurse-managed intervention. *Ann Intern Med* 1990;113:118-23.
- 45 Jones DA, West RR. Psychological rehabilitation after myocardial infarction: multicentre randomised controlled trial. *BMJ* 1996;313:1517-21.
- 46 Friedman M, Thoresen CE, Gill JJ, et al. Alteration of type A behaviour and its effect on cardiac recurrences in post myocardial infarction patients: summary results of the recurrent coronary prevention project. *Am Heart J* 1986;112:653-65.
- 47 van Elderen-van Kemenade T, Maes S, van den Broek Y. Effects of a health education programme with telephone follow-up during cardiac rehabilitation. *Br J Clin Psychol* 1994;33:367-78.
- 48 Monteiro LA. *Cardiac patient rehabilitation: social aspects of recovery*. New York: Springer, 1982.
- 49 Newens AJ, McColl E, Lewin R, et al. Cardiac misconceptions and knowledge in nurses caring for myocardial infarction patients. *Coronary Health Care* 1997;1:83-9.
- 50 Kallio V, Hamalainen H, Hakila J, et al. Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979;2:1091-4.
- 51 Oldridge NB, Guyatt GH, Fischer ME, et al. Cardiac rehabilitation after myocardial infarction. Combined experience of randomised clinical trials. *JAMA* 1988;260:945-50.
- 52 O'Connor GT, Buring JE, Yusuf S, et al. An overview of randomised trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989;80:234-44.
- 53 Bobbio M. Does post myocardial infarction rehabilitation prolong survival? A meta-analytic survey. *Giornale Italiano di Cardiologia* 1989;19:1059-67.
- 54 Carlsson R, Lindberg G, Westin L, et al. Influence of coronary nursing management follow up on lifestyle after acute myocardial infarction. *Heart* 1997;77:256-9.
- 55 Engblom E, Korpilahti K, Hamalainen H, et al. Quality of life and return to work 5 years after coronary artery bypass surgery: long-term results of cardiac rehabilitation. *J Cardpulm Rehab* 1997;17:29-36.
- 56 Picard MH, Dennis C, Schwartz RG, et al. Cost-benefit analysis of early return to work after uncomplicated acute myocardial infarction. *Am J Cardiol* 1989;63:1308-14.
- 57 Ades PA, Waldmann ML, Polk DM, et al. Referral patterns and exercise response in the rehabilitation of female coronary patients aged greater than or equal to 62 years. *Am J Cardiol* 1992;69:1422-5.
- 58 McGee HM, Horgan JH. Cardiac rehabilitation programmes: are women less likely to attend? *BMJ* 1992;305:283-4.
- 59 Cannistra LB, Balady GJ, O'Malley CJ, et al. Comparison of the clinical profile and outcome of women and men in cardiac rehabilitation. *Am J Cardiol* 1992;69:1274-9.
- 60 Lavie CJ, Milani RV. Benefits of cardiac rehabilitation and exercise training in elderly women. *Am J Cardiol* 1997;79:664-6.
- 61 Hellman EA, Williams MA. Outpatient cardiac rehabilitation in elderly patients. *Heart and Lung* 1994;23:506-12.
- 62 Ades PA, Hanson JS, Gunther PG, et al. Exercise conditioning in the elderly coronary patient. *J Am Geriatr Soc* 1987;35:121-4.
- 63 Williams MA, Maresh CM, Esterbrooks DJ, et al. Early exercise training in patients older than age 65 years compared with that in younger patients after acute myocardial infarction or coronary artery bypass grafting. *Am J Cardiol* 1985;55:263-6.
- 64 Lavie CJ, Milani RV, Littman AB. Benefits of cardiac rehabilitation and exercise training in secondary coronary prevention in the elderly. *J Am Coll Cardiol* 1993;22:678-83.
- 65 Bondestam E, Breikks A, Hartford M. Effects of early rehabilitation on consumption of medical care during the first year after acute myocardial infarction in patients greater than or equal to 65 years of age. *Am J Cardiol* 1995;75:767-71.
- 66 Ferguson JA, Tierney WM, Westmoreland GR, et al. Examination of racial differences in management of cardiovascular disease. *J Am Coll Cardiol* 1997;30:1707-13.
- 67 Young RF, Waller JB, Jr, Kahana E. Racial and socioeconomic aspects of myocardial infarction recovery: studying confounds. *Am J Prev Med* 1991;7:438-44.
- 68 Cannistra LB, O'Malley CJ, Balady GJ. Comparison of outcome of cardiac rehabilitation in black women and white women. *Am J Cardiol* 1995;75:890-3.
- 69 Romeo KC, Saccucci MS. Ethnic and gender differences in cardiac rehabilitation. *J Cardiopulm Rehabil* 1991;11:214-20.
- 70 Pell J. Cardiac rehabilitation: a review of its effectiveness. *Coronary Health Care* 1997;1:8-17.

- 71 WHO Joint Consultation Group. *Cardiac rehabilitation and secondary prevention: long term care for patients with ischaemic heart disease*. Geneva: WHO Regional Office for Europe, 1993.
- 72 Coats AJ. Exercise rehabilitation in chronic heart failure. *J Am Coll Cardiol* 1993;22:172a-7a.
- 73 Uren NG, Lipkin DP. Exercise training as therapy for chronic heart failure. *Br Heart J* 1992;67:430-3.
- 74 Kavanagh T, Myers MG, Baigrie RS, et al. Quality of life and cardiorespiratory function in chronic heart failure: effects of 12 months' aerobic training. *Heart* 1996;76:42-9.
- 75 Hedback B, Perk J. Can high-risk patients after myocardial infarction participate in comprehensive cardiac rehabilitation? *Scandinavian Journal of Rehabilitation Medicine* 1990;22:15-20.
- 76 Lewin B, Cay EL, Todd IC, et al. The angina management programme: a rehabilitation treatment for chronic angina pectoris. *Br J Cardiol* 1995;2:221-6.
- 77 Lewin B. The psychological and behavioural management of angina. *J Psychosom Res* 1997;5:452-62.
- 78 Todd IC, Ballantyne D. Antianginal efficacy of exercise training: a comparison with β blockade. *Br Heart J* 1990;64:14-9.
- 79 Cupples ME, McKnight A. Randomised controlled trial of health promotion in general practice for patients at high cardiovascular risk. *BMJ* 1994;309:993-6.
- 80 Campbell NC, Thain J, Deans HG, et al. Secondary prevention clinics for coronary heart disease: randomised trial of effect on health. *BMJ* 1998;316:1434-7.
- 81 Sparks KE, Shaw DK, Eddy D, et al. Alternatives for cardiac rehabilitation patients unable to return to a hospital-based program. *Heart and Lung* 1993;22:298-303.
- 82 DeBusk RF, Haskell WL, Miller NH, et al. Medically directed at-home rehabilitation soon after clinically uncomplicated acute myocardial infarction: a new model for patient care. *Am J Cardiol* 1985;55:251-7.
- 83 Lewin B, Robertson IH, Cay EL, et al. Effects of self-help post-myocardial-infarction rehabilitation on psychological adjustment and use of health services. *Lancet* 1992;339:1036-40.
- 84 Linden B. Evaluation of a home-based rehabilitation programme for patients recovering from acute myocardial infarction. *Intensive and Critical Care Nursing* 1995;11:10-9.
- 85 DeBusk RF, Miller NH, Superko HR, et al. A case-management system for coronary risk factor modification after acute myocardial infarction. *Ann Intern Med* 1994;120:721-9.
- 86 Taylor CB, Miller NH, Smith PM, et al. The effect of a home-based, case-managed, multifactorial risk-reduction program on reducing psychological distress in patients with cardiovascular disease. *J Cardiopulm Rehabil* 1997;17:157-62.
- 87 Haskell WL, Alderman EL, Fair JM, et al. Effects on intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease: the Stanford coronary risk intervention project (SCRIP). *Circulation* 1994;89:975-90.
- 88 Hung J, Gordon EP, Houston N, et al. Changes in rest and exercise myocardial perfusion and left ventricular function 3 to 26 weeks after clinically uncomplicated acute myocardial infarction: effects of exercise training. *Am J Cardiol* 1984;54:943-50.
- 89 Ueshima K, Saito M, Shimohara A, et al. Management and evaluation of non-supervised home exercise program in a convalescent phase of acute myocardial infarction. *Japanese Circulation Journal* 1990;54:1437-42.
- 90 Brosseau R, Juneau M, Sirard A, et al. Safety and feasibility of a self-monitored, home-based phase II exercise program for high risk patients after cardiac surgery. *Can J Cardiol* 1995;11:675-85.
- 91 Barnason S, Zimmerman L. A comparison of patient teaching outcomes among postoperative coronary artery bypass graft (CABG) patients. *Prog Cardiovasc Nurs* 1995;10:11-20.
- 92 Gattiker H, Goins P, Dennis C. Cardiac rehabilitation. Current status and future directions. *West J Med* 1992;156:183-8.
- 93 Pell J, Pell A, Morrison C, et al. Retrospective study of influence of deprivation on uptake of cardiac rehabilitation. *BMJ* 1996;313:267-8.
- 94 Oldridge NB. Compliance and exercise in primary and secondary prevention of coronary heart disease: a review. *Prev Med* 1982;11:56-70.
- 95 Radley A, Grove AM, Thurston H, et al. *WE 8: A study of the rehabilitation needs of women cardiac patients. Report to the NHS research and development directorate CVD and stroke programme*. Loughborough: Department of Social Sciences, Loughborough University, 1996.
- 96 Gori P, Pivotti F, Mase N, et al. Compliance with cardiac rehabilitation in the elderly. *Eur Heart J* 1984;5(suppl E):109-11.
- 97 Andrew GM, Oldridge NB, Parker JO, et al. Reasons for dropout from exercise programs in post-coronary patients. *Medicine and Science in Sports and Exercise* 1981;13:164-8.
- 98 Ades PA, Waldmann ML, McCann WJ, et al. Predictors of cardiac rehabilitation participation in older coronary patients. *Arch Intern Med* 1992;152:1033-5.
- 99 Campbell N, Ritchie L, Rawles J, et al. Cardiac rehabilitation: the agenda set by post-myocardial infarction patients. *Health Education Journal* 1994;53:409-20.
- 100 Dracup K, Meleis AI, Clark S, et al. Group counselling in cardiac rehabilitation: effect on patient compliance. *Patient Education and Counseling* 1984;6:169-77.
- 101 Levy JK. Standard and alternative adjunctive treatments in cardiac rehabilitation. *Texas Heart Institute Journal* 1993;20:198-212.
- 102 Fletcher GF, Balady G, Blair SN, et al. Statement on exercise: benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the committee on exercise and cardiac rehabilitation of the council on clinical cardiology, American Heart Association. *Circulation* 1996;94:857-62.
- 103 O'Callaghan WG, Teo KK, O'Riordan J, et al. Comparative response of male and female patients with coronary artery disease to exercise rehabilitation. *Eur Heart J* 1984;5:649-51.
- 104 Oldridge NB, Furlong W, Feeny D, et al. Economic evaluation of cardiac rehabilitation soon after acute myocardial infarction. *Am J Cardiol* 1993;72:154-61.
- 105 Bucher HC. Social support and prognosis following first myocardial infarction. *J Gen Intern Med* 1994;9:409-17.