

ORIGINAL ARTICLE

Case record review of adverse events: a new approach

M Woloshynowych, G Neale, C Vincent

See editorial commentary, pp 402–3

Qual Saf Health Care 2003;12:411–415

See end of article for authors' affiliations

Correspondence to:
Dr M Woloshynowych,
Clinical Safety Research
Unit, Imperial College,
Department of Surgical
Oncology and Technology,
St Mary's Hospital, London
W2 1NY, UK;
m.woloshynowych@
imperial.ac.uk

Accepted for publication
7 August 2003

Objectives: To redesign the existing clinical review form (RF2) used in previous retrospective case record review studies in order to clarify the review process and provide a more powerful analysis of adverse events; and then to ask clinicians to pilot and evaluate the new modular review form (MRF2). The review form is divided into five sections, each with a defined purpose, providing a modular structure.

Design: Design and testing of the MRF2 on a sample of medical and nursing records, and evaluation of the reviewers' responses regarding the new review form.

Setting: Hospital based teams from eight countries.

Results: The modular review form was reported to be comprehensive, well structured, and clear. Most of the reviewers agreed with the positive statements regarding the review form. Overall, the modular structure was thought to be helpful. Several modifications have been made to the final version to take account of criticisms and suggestions.

Conclusions: The full potential of case record review has yet to be explored. The benefits of this review form include a modular format which enables reviewers or project leaders to select the focus of their review based on resources and the purpose of the review, and to identify contributory factors which indicate areas for improvement and prevention. The training of reviewers is of vital importance for record review. Record review remains one of the primary methods for assessing the incidence of adverse events and the new format is suitable for both prospective and retrospective review.

The use of case record review to identify adverse events has been important in the drive to improve patient safety.^{1,2} The Harvard Medical Practice study³ carried out in New York in the mid 1980s provided powerful evidence of the scale of harm to patients in hospital. These initial findings were substantiated in further studies in the United States,⁴ in Australia,⁵ the UK,⁶ Denmark,⁷ and New Zealand,⁸ and there are ongoing studies in other countries.

The basic methodology was developed in the early 1970s for the Californian Insurance Feasibility Study.⁹ Retrospective case record review studies are carried out in two stages. First, using Review Form 1 (RF1), medical records are screened according to 18 predefined criteria (for example, unexpected death, hospital acquired infection/sepsis, unplanned return to the operating theatre) to identify records of patients more likely to have suffered an adverse event. Records meeting one or more of the screening criteria are forwarded for clinical review using Review Form 2 (RF2). In this second stage trained clinicians examine each case record in detail to determine whether or not an adverse event has occurred and to extract information about the nature and causes of adverse events. Each research group^{4–8} made minor modifications to the RF2, adding or subtracting specific questions, but the basic format used in the Harvard study³ was maintained.

The methodology of case record review has considerable strengths. It has provided a more complete indication of the incidence of adverse events or critical incidents than reporting systems, even when they are backed by additional monitoring by a dedicated risk manager.¹⁰ Familiarity and other biases are reduced when external independent assessors are used to conduct the review. The review forms provide a standardised method of recording and data collection which is robust when used on a random sample of case records. The epidemiological data obtained are potentially useful for comparative studies, although any comparisons need to take account of variations in methodology, particularly with the definition and inclusion criteria.

Readers are invited to respond to the three questions listed at the end of the accompanying commentary by Ross Wilson on page 402 by using the rapid response function <http://qhc.bmjournals.com/cgi/eletter-submit/12/6/402>.

Case record review is, however, wholly dependent on the accuracy, completeness and legibility of patient records. Some information, such as the effects of the adverse event on the patient, is not generally recorded and often the adverse event itself is not explicitly stated in the record and may not be recognised until the patient is readmitted. Low to moderate inter-rater reliability has been reported.^{5,11} Finally, retrospective case record review can be time consuming and expensive. Despite these limitations, however, case record review has yielded important epidemiological data that have had a major effect on governmental policies and action by healthcare providers.²

RATIONALE FOR A REVISED REVIEW FORM

Earlier studies^{3–5} focused on the incidence and type of adverse events with some attempt to identify causes and methods of prevention.¹² The original review form uses a mixture of taxonomies which are not always clearly distinguished. Reviewers in our study found that the structure of the review form meant that results did not always reflect the underlying clinical reality. For instance, with a standard RF2 a postoperative infection is classified as an operative event (because it occurred within 30 days of surgery) and non-technical (because it was not directly related to the operation itself). In a subsequent examination of the data we analysed in detail the narratives provided by the reviewers.¹³ This enabled us to define when in the process of care the adverse event occurred and the nature of the underlying problem

(box 1, C1–C5). As a result, a postoperative chest infection would be classified as a ward based event and the problem as one related perhaps to drug administration (oversedation), to failure to monitor (such as over a weekend), or to failure to provide physiotherapy.

Adverse events frequently involve a chain of events and a series of care management or care delivery problems leading up to the incident itself.¹⁴ In previous versions of the form multiple problems were sometimes identified, but were not prioritised and were not sufficiently distinguished in the final analyses. However, we found that there is usually one critical period in which the primary problem occurred. We therefore introduced the concept of a “principal problem”, being the most important problem in the delivery of care. This enabled a precise identification of a particular period in the process of care, now incorporated in the review process, and a defined principal problem which greatly clarified the review process. To provide additional clarity we divided the review form into five sections each with a defined purpose, providing a modular structure to both the form and the review process.

A final important change has been a much stronger emphasis on the broader organisational, environmental, and other factors that contribute to adverse events. While earlier versions paid some attention to these factors, the approach was not sufficiently systematic. The present approach is based on a previously devised framework¹⁵ and method of individual case analysis.¹⁶ Identifying contributory factors is best done from observation and interview, although this is quite labour and resource intensive. We believe that expert reviewers or those familiar with the working environment can often comment on some of the major contributory factors. This is particularly important as the identification of such factors offers a route to devising methods of prevention.

Thus, we have attempted to address three areas in which the earlier review forms have limitations:

- We have divided hospital stay into distinct periods of care and then identified the clinical aspects associated with each period—to address the former mixed categories of the type of adverse event.
- We have introduced the term “principal problem” to help clarify the review process in order to identify the healthcare management or treatment, or delay or lack of treatment, that led to the adverse event.
- We have provided a comprehensive list of contributory factors to include organisational, environmental, and other contextual factors that influence the provision of health care.

THE MODULAR REVIEW FORM (MRF2)

The MRF2 comprises five stages or modules:

- (A) Patient information and background to the adverse event
- (B) Disability caused by the adverse event
- (C) Period of hospitalisation during which the adverse event occurred
- (D) Principal problems in the process of care
- (E) Causative/contributory factors and preventability of the adverse event

Once an adverse event has been identified following the first stage review (RF1), reviewers are required to complete stages A and B in full. When completing stage A they are required to identify one or more of five periods of care during which adverse event(s) occurred (C1–C5). Next they

Box 1 Structure of the modular review form (MRF2)

(A) Patient information and background to adverse event (AE)

- Patient data
- Primary illness; prognosis
- Co-morbidity; speciality
- Main features of AE
- Identification of principal problem
→ identify period(s) of care (C1–C5)
- Adequacy of records

(B) The injury and its effects

- Disability (including death) caused by AE
- Effect of AE on hospital resources (e.g. additional bed days)
- Additional treatment as a result of AE

(C) Period of hospitalisation during which AE occurred (as identified in stage A)

- C1: Care on admission ward including preoperative care →
- C2: During procedure including surgery, anaesthesia →
- C3: Immediate postoperative and ITU/HDC care →
- C4: General ward care →
- C5: Care/assessment during discharge
→ principal problem (D1–D7)

(D) Principal problems in the process of care (as identified in relevant subsection(s) in C)

- D1: Diagnostic/assessment error
- D2: In relation to patient's overall condition
- D3: Medical management/monitoring including nursing care
- D4: In relation to infection
- D5: Procedure (including anaesthesia/surgery)
- D6: In relation to drug/IV fluid/blood transfusion
- D7: In relation to a resuscitation procedure

(E) Causative/contributory factors and preventability of AE

- Causative factors: patient characteristics, task factors, individual factors, team factors, work environment, organisational/management factors
- Preventability of AE
- Expertise of reviewer

complete the relevant part(s) of stage C. In so doing, they are required to identify the nature of the underlying cause(s) of the adverse event ranging from errors in diagnosis to mismanagement at the time of discharge. This leads to the completion of one or more of subsections D1–D7. Finally, stage E provides the opportunity for identifying contributory factors and potential preventability. Details of each stage are shown in box 1 and fig 1. The modular structure allows reviewers to select which stages to complete, depending on the focus of the review. The MRF2 is available on the QSHC website (<http://www.qshc.com/supplemental>).

The objectives of this study were to pilot and evaluate the new review form (MRF2) using an evaluation questionnaire.

Table 1 Responses to questionnaire statements

Statement	Mean	Median	Scores*				
			1	2	3	4	5
The review form is comprehensive	3.92	4	0	0	2	9	1
The review form is well structured	3.58	4	1	1	2	6	2
The review terminology is clear	3.42	4	0	4	0	7	1
The modular structure is useful	3.92	4	0	1	2	6	3
The amount of explanation and instruction on the form itself is about right	3.42	4	0	3	2	6	1
The accompanying separate instructions are helpful	3.67	4	0	1	3	7	1
The range of options for each section is sufficient	3.97	4	0	3	0	7	2
The review form records sufficient patient information	3.83	4	0	2	1	6	3
The review form records sufficient detail about problems in process of care	3.67	4	0	3	1	5	3
The review form records sufficient detail about the impact of AE	3.75	4	0	3	0	6	3
The review form records sufficient detail about the causes of AE	3.58	4	0	4	0	5	3
The information on the review form has the potential to lead to meaningful clinical improvements	3.58	4	0	2	2	7	1

AE = adverse event.

*The MRF2 was rated on a 5 point scale: 1 = strongly disagree; 2 = disagree; 3 = no opinion; 4 = agree; 5 = strongly agree.

Box 2 Comments from reviewers

General comments

Strengths of the review form

- Layout comprehensive and easy to follow.
- Comprehensive but lengthy—relatively easy to complete.
- The modular structure is definitely helpful in identifying the particular period of care in which problems arose.
- Clear separation of different components allows detailed review.
- Good basis; good mix of open ended questions and categorisation of events.
- Useful for clinicians for risk management.
- Strong emphasis on identifying systemic factors and contributory causes.
- Easy to complete.
- Problems can be identified with greater precision by probing with follow up questions.
- Specific and potential to obtain a lot of information.

Limitations of the review form

- Layout initially appears confusing.
- Too complex—too specific and too exhaustive—need specialist experience to complete and/or considerable dedication to unearth circumstances surrounding an adverse event.
- Not enough information for epidemiological analysis but too complicated for clinical use.
- Not suitable for root cause analysis.
- Some questions are too subjective—will get different answers from different reviewers.
- Too time consuming.
- Sometimes impossible to answer questions—need more “don’t know” or “data not available” options.
- Repetition in some areas.

Devising a review form that is capable of encompassing almost all aspects of hospital care inevitably results in a long document. We have, where possible, tried to reduce the level of detail in the interest of making the whole process more manageable. We purposely kept the guidance notes short and

inserted definitions within the review form itself, as some reviewers in our previous study⁶ found it irksome to consult the manual while completing the form. While the complete form may still look daunting, reviewing a single record only involves completing a relatively small number of specific subsections. The process could be made more efficient with an electronic version of the form. Reviewers may have to adapt the forms to their particular country and circumstances, and perhaps add additional guidance notes appropriate to the local setting. However, results from this pilot study suggest that the underlying structure should prove robust. Reviewers with a particular interest in a certain type of adverse event may wish to include much more detailed questions on some aspects of care.

Our experience of studying adverse events and piloting these forms indicates that training of reviewers remains of vital importance. Clinical ability and acumen, although essential, do not guarantee that the review process will be either understood or adhered to. Preparation by review of trial cases followed by discussion is essential. Without training in identifying the key issues quickly and efficiently, the process can become unduly time consuming. In our experience^{6,13} most disagreements usually revolve around definitions and terminology rather than differences of opinion on clinical matters.

We acknowledge that the present study is only a preliminary test of the modified review form and that it needs to be compared with previous versions and tried out in larger studies in a variety of settings. As no formal comparison with the old form has been made, we are unable to comment on whether the MRF2 is an improvement on previous versions. Furthermore, we were unable to control for variation in the specific instructions and information that was given to the reviewers by team leaders or the variability of specialties and experience of each review team. However, it is worth acknowledging the practical setting in which these forms may be used. We consider that the positive nature of the present findings suggests that the modified form should be considered for use in future case record reviews.

In our view the full potential of case record review has yet to be explored. Most adverse event reviews conducted so far have been labour intensive, large scale studies. However, we believe that there is considerable potential for small scale studies which may be either local clinical review or formal research. Case record review could be especially valuable for routine review at the local level, particularly when specific types of cases are targeted and with a full exploration of causes and methods of prevention. The modular structure of the present form allows those with limited resources to cut

Table 2 Specific comments from reviewers to MRF2

Stage	Suggestions/difficulties	Authors' comments
A	Need to elucidate meaning of principal problem Add to co-morbidity list Include time of event	This has been clarified by re-structuring the stage and giving examples The co-morbidity list has been modified Space to record the time of the event is included
B	Answers to the following will be subjective: <ul style="list-style-type: none"> ● Additional bed days caused by AE ● Degree of disability ● Emotional trauma 	Agree—but approximate data are useful
C	May be difficult to determine: <ul style="list-style-type: none"> ● Person responsible at that period of care ● Date and time of event 	Agree—but may be able to determine whether or not there was lack of input from experienced staff
D	May be difficult to determine: <ul style="list-style-type: none"> ● How management contributed to the AE and respects in which it was unsatisfactory ● Quality of hand over ● Person responsible at the time of AE and whether or not it was appropriate ● Whether there were avoidable delays or inappropriate procedures ● The cause of drug related injury 	Agree—but experienced reviewers will be able to make “intelligent estimates”
E	Impossible to relate the relative importance of contributory factors Difficult to determine reasons for failure to prevent AE Specialist reviewers would be needed	It is important that reviewers are adequately trained and that they are aware of their own limitations As the form is modular, stage E may be omitted. Specialists should be consulted on specific points

Key messages

- For 25 years a basic form has been used for recording the results of case record review to identify adverse events in hospital practice.
- The method has provided epidemiological data but has been of limited value in defining targets for improvement in care
- We present a revised form which has a modular structure. Key components define (1) the period of care during which action or inaction led to the adverse event and (2) the type of clinical action or inaction.
- A further module offers the opportunity of recording factors that may have contributed to the adverse event.
- This revised form received mostly favourable comments when assessed by researchers in 12 other units.
- The method needs to be tested in practice; it may be of particular value in institutions that wish to identify areas of improvement based on contributing factors.
- As with previous versions of the form, training in completing the form is essential.

down the review process to its essentials and adapt it to their particular purpose. There is also considerable scope for using record review prospectively in combination with staff interviews. Record review, whether retrospective or prospective, remains one of the primary methods for assessing the incidence of adverse events.^{10 11}

ACKNOWLEDGEMENTS

The authors thank the Dunhill Medical Trust and the BUPA Foundation for funding this pilot study; and J Minton (UK), C Pearson (UK), J Smit (Australia), M Silver (USA), J M Aranaz (Spain), R Cinotti (Italy), P Michel (France), A Gillian (Australia), M Kelly (USA), D Kunac (New Zealand), F Briganti (Italy), T Ohkubo (Japan), and their colleagues for their time and comments regarding the MRF2.



The Modular Review Form 2 (MRF2) is available on the QSHC website at www.qshc.com/supplemental.

Authors' affiliations

M Woloshynowych, G Neale, C Vincent, Clinical Safety Research Unit, Imperial College, Department of Surgical Oncology and Technology, St Mary's Hospital, London W2 1NY, UK

REFERENCES

- 1 **Vincent C**. Risk, safety and the dark side of quality. *BMJ* 1997;**314**:1775–6.
- 2 **Neale G, Woloshynowych M**. Retrospective case record review: a blunt instrument that needs sharpening. *Qual Saf Health Care* 2003;**12**:2–3.
- 3 **Brennan TA, Leape LL, Laird NM, et al**. Incidence of adverse events and negligence in hospitalised patients. Results of the Harvard Medical Practice Study I. *N Engl J Med* 1991;**324**:370–6.
- 4 **Thomas EJ, Studdert DM, Burstin HR, et al**. Incidence and types of adverse events and negligent care in Utah and Colorado. *Med Care* 1999;**38**:261–71.
- 5 **Wilson RM, Runciman WB, Gibberd RW, et al**. The Quality in Australian Health Care Study. *Med J Aust* 1995;**163**:458–71.
- 6 **Vincent C, Neale G, Woloshynowych M**. Adverse events in British hospitalized patients: preliminary retrospective record review. *BMJ* 2000;**322**:517–9.
- 7 **Schiolar T, Lipczak H, Pedersen BL, et al**. Forekomsten af utilsigtede haendesler pa sygehuse. En retrospektiv gennemgang af journaler. *Ugeskr Laeger* 2001;**163**:5370–8.
- 8 **Davis P, Lay-Yee R, Schug S, et al**. Adverse events regional feasibility study: indicative findings. *NZ Med J* 2001;**114**:203–5.
- 9 **Mills DH**. Medical insurance feasibility study. A technical study. *West J Med* 1978;**128**:360–5.
- 10 **Stanhope N, Crowley-Murphy M, Vincent C, et al**. An evaluation of adverse incident reporting. *J Eval Clin Pract* 1999;**5**:5–12.
- 11 **Thomas EJ, Peterson LA**. Measuring errors and adverse events in health care. *J Gen Intern Med* 2002;**18**:61–7.
- 12 **Wilson RM, Harrison BT, Gibberd RW, et al**. An analysis of the causes of adverse events from the Quality in Australian Health Care Study. *Med J Aust* 1999, **170**:411–5.
- 13 **Neale G, Woloshynowych M, Vincent C**. Exploring the causes of adverse events in British NHS hospital practice. *J R Soc Med* 2001;**94**:322–30.
- 14 **Vincent C, Taylor-Adams S, Chapman EJ, et al**. How to investigate and analyse clinical incidents: clinical risk unit and association of litigation and risk management protocol. *BMJ* 2000;**320**:777–81.
- 15 **Vincent CA, Taylor-Adams S, Stanhope N**. A framework for the analysis of risk and safety in medicine. *BMJ* 1998;**316**:1154–7.
- 16 **Vincent CA, Taylor-Adams S, Chapman EJ, et al**. *A protocol for the investigation and analysis of clinical incidents*. London: University College London/Association of Litigation and Risk Management, 1999.