

ERROR MANAGEMENT

Patients use an internet technology to report when things go wrong

John H Wasson, Todd A MacKenzie, Michael Hall

Qual Saf Health Care 2007;16:213–215. doi: 10.1136/qshc.2006.019810

See end of article for authors' affiliations

Correspondence to:
Dr J H Wasson, Dartmouth Medical School, Hanover, New Hampshire 03755, USA; john.h.wasson@dartmouth.edu

Accepted 30 January 2007

Background: As patients directly experience harm from adverse events, investigators have proposed patient-report to complement professional reporting of adverse events.

Objective: To investigate how an automated health assessment system can be used to identify adverse events.

Design and setting: Internet survey responses from April 2003 to April 2005 involving communities and clinical practices across the USA.

Patients: 44 860 adults aged 19–69 years.

Outcome: Patient perceptions of adverse events experienced during the previous year. Independent legal review was also used to estimate how many patient-reports were serious enough to be potentially compensable.

Results: Although patient reports of possible adverse events was low (1.4%), the percentage of adverse events was eight times higher for patients with the greatest burden of illness than for those with the least (3.4% vs 0.4%). Two expert malpractice attorneys agreed that 9% of the adverse events seemed to be serious.

Conclusions: Patients will use internet technology to report their perceptions of health-related adverse events. Some of the patient-reported events reported will be serious.

A highly influential report from the Institute of Medicine, Washington DC, USA, "To err is human: building a safer health care system", encouraged better documentation of errors and adverse events, so that causes for problems might be identified and remedied.¹ However, reporting of error and adverse events by health professionals depends on variables such as their awareness of the events, their judgment about events and their willingness to document the events. Incidence rates for adverse events vary 10-fold when reported by health professionals; differences are attributed to professional attitude and judgment rather than the underlying rate of errors and adverse events.^{2–3} Since patients directly experience harm from adverse events, investigators have proposed patient-report to complement professional reporting.^{4–8} Although the proposition that patient reporting of adverse events has merit, the feasibility of patient-report is not known.

This study investigates how an automated outpatient health assessment system can be used to identify adverse events.

METHOD

The health assessment survey and the study population

More than 20 years ago, the Dartmouth Cooperative Practice-Based Research Network identified important, wide gaps in the communication between patients and doctors.⁹ As an example, this research documented the need for and stimulated the development of patient-report measures of functional status.¹⁰ The Cooperative Practice-Based Research Network now widely disseminates a patient health assessment survey to improve patient–clinician communication.^{11–16}

Using computer-generated branching logic, up to 120 items of the survey inquire about health status, symptoms, concerns, chronic disease management, preventive interventions, specific healthcare experiences and confidence with self-care (<http://www.howsyourhealth.org>). Based on their own initiative or at the request of their doctor, people complete the health survey to obtain health and self-management information customised to their needs.

As no personal identifiable information is stored by any means, patients decide on how to use their information. For

example, they can deliver a clinical summary of the information (electronically or in person) to their doctor so that they and the doctor can review and manage the important problems identified by the survey in a better way.

Item completion rates are >95%. Whenever a patient reports that he/she has completed the survey in the previous 6 months, the recent information is not stored. The data are exempt from institutional review board review as no personal identifiable data is stored.

This report is based on 44 860 cross-sectional responses by residents of the USA, aged 19–69 years during the 2-year time frame from April 2003 to April 2005.

A measure of adverse events

Near the end of the survey one item asks:

Part A: "Describe here any medical errors (mistakes) that you or your family have experienced. Errors include such things as mixed up medications or poor treatment that result in harm or additional problems. If possible, be sure to tell us the cause of the error and how it might have been avoided. Your response will help us to improve future care delivery."

Part B: "If you wrote in an error or harm, please help us by choosing ANY of the following categories for this error. (Please mark all that apply.) It caused harm, hurt or injury. It happened within the last year. It happened to me."

What is an adverse event?

Although an individual clinician receiving a patient report of an adverse event might be able to corroborate it with the information in the medical record, for this large study population of patients only their verbatim comments were available to us. To contend with the possibility that the patients' reports might represent mere complaints about poor service, we used independent legal review to estimate how many, if any, patient-reports were possibly serious—that is, the percentage of reports that might, after additional review, cross the threshold from nuisance to potentially compensable injury.

A lawyer (MH) with extensive experience in medical malpractice litigation reviewed all patient-reported adverse

Table 1 Health-related adverse events and patient characteristics

	Respondents reporting no health-related adverse event in the previous year 98.6% (n = 44 250)	Respondents reporting a health-related adverse event in the previous year 1.4% (n = 610)
Demographics		
Female	65	79*
Age 19–49 years	68	63
Low financial status	15	27*
Bothersome dysfunction		
Pain	13	44*
Emotional	10	24*
Social support	11	26*
Physical activity	7	23*
Daily activity	4	24*
Social activity	4	22*
Chronic disease		
Hypertension	20	24
Respiratory	10	20*
Arthritis	14	26*
Diabetes	6	11*
Cardiovascular	4	8*
Symptoms		
>1 of 14 bothersome symptoms	33	65*
Composite Burden of Illness Score† (dysfunction, disease, symptoms)		
	3.0	4.8*
Unique prescription medications‡ ≥3		
	20	45*
Any hospital use in the previous year		
	8	33*
Lifestyle		
Not exercising regularly	59	61*
BMI ≥30	31	36
Poor injury prevention or poor eating habits	20	23
Smoking	14	22
Told to “cut down” drinking	7	8*
Composite Health Behaviour Score		
	3.0	3.4*

BMI, body mass index.
 *Significance of difference is $p < 0.01$.
 †Burden of Illness ranges from 0 (none) to 11; Health Behaviour from 0 (no risk) to 15.
 All values are in percentages except for Composite Burden of Illness Score and Composite Health Behaviour Score.

events and selected those that seemed serious enough to represent compensable injury: when the costs of the legal remedy could be greater than the costs of litigation.

We derived a conservative estimate of potentially compensable injuries by subjecting the adverse events identified by the first legal review to another review. The cases that were agreed upon by two attorneys constitute our estimate of potentially serious adverse events. A verbatim list of these events is available at <http://www.howsyourhealth.org/adverse.pdf>.

ANALYSIS

We have limited the analysis to descriptive statistics as the data are cross-sectional and are based on a convenience sample of patients out of a much larger potential population of non-respondents in clinical practices or communities.




Figure 1 Annual rate per 100 of health-related adverse events associated with the patients' burden of illness. Burden of illness is a composite score based on the number of chronic diseases, the number of bothersome symptoms and the number of bothersome functional problems. The 95% CI is shown for each quintile of burden from the lowest to the highest.

RESULTS

From the 2-year convenience sample of 44 860 survey respondents, 2979 indicated a harm, hurt or injury to themselves or a family member, and 610 of these adverse events had happened to the respondents during the previous year.

Association of possible adverse events with burden of illness

Table 1 lists the characteristics of patients who reported health-related adverse events and those who did not. All but four (age, hypertension, exercise and excessive alcohol consumption) of the many listed variables were statistically different between the two groups of patients. In the stepwise regression of the demographic and composite variables listed in table 1, burden of illness had by far the greatest association with a report of a possible adverse event.

Figure 1 depicts the strong association between possible adverse events and burden of illness. The annual percentage of health-related adverse events ranged eightfold, from 0.4% to 3.4% across quintiles of illness burden.

Severity of possible adverse events

In all, 574 of the 610 patients provided descriptive information about what went wrong. The majority of these events occurred in an outpatient setting; only 12% (66) of the possible adverse events transpired in a hospital.

Two expert malpractice attorneys agreed that 9% (52/574) of the adverse events seemed serious enough to represent a possible compensable injury. In absolute terms, twice as many serious events were experienced in the ambulatory setting (n = 35) than in a hospital (n = 17). However, in relative terms, 26% (17/66) of the hospital events were possibly compensable compared with 7% of those in an outpatient setting.

COMMENT

This study demonstrates that patients will use internet technology to report their perceptions of health-related adverse events. Patients with a greater burden of illness are more likely to report adverse events than those with little burden. The legal

review adds credence to the notion that some patient reports will be more serious than mere complaints about poor service.

These results suggest an inexpensive mechanism by which a clinician can learn about a patient's perceptions of previous poor care, acknowledge the problem, investigate its cause and initiate a remedy.¹⁷ By being patient-centered and giving the perceptions of possible adverse events a human face, such systems could be useful for influencing the way the outpatient clinicians think about safety.¹⁸ Similar systems are also available for inpatients (<http://www.howsyourcare.org>).

Despite the promise of such automated systems for assessing health and healthcare, the findings have at least three significant limitations.

First, since the patient denominator is a non-random sample from an unknown population, the adverse events reported in this study should not be interpreted as a benchmark rate based on a national probability sample of patients. A patient-reported adverse event will be biased by the background of the person who chose to respond to an online "health check-up" and their opportunities to interact with the healthcare system.

Second, in addition to the limitations intrinsic to our specific approach, both professional and patient-derived reports of "when things go wrong" have biases that reflect the attitudes and experiences of the reporters, their incentives or disincentives to report, and the ease of use and fidelity of the reporting system.^{2-8 19}

Finally, as yet there is no proof that patient-reported perceptions of health-related harms will improve the safety and quality of care.

In summary, we have shown how an internet-based, patient assessment survey can produce information about health-related adverse events. More research is needed to demonstrate the value of such automated systems in improving healthcare.

This research was supported by grants from The Commonwealth Fund and the Robert Wood Johnson Foundation.

ACKNOWLEDGEMENTS

We thank Judy Knapp and L Gordon Moore, MD, for assisting with the typology of adverse events.

Authors' affiliations

John H Wasson, Dartmouth Medical School, Hanover, New Hampshire, USA

Todd A MacKenzie, Norris Cotton Cancer Center, One Medical Center Drive, Hanover, New Hampshire, USA

Michael Hall, Hall Stewart, Manchester, New Hampshire, USA (posthumous)

Competing interests: The results are based on the use of an internet technology www.howsyourhealth.org. Although the website is freely available without advertising or support from any commercial entities, it does cover its costs through a nominal fee for custom use by some subscribers. Under a license agreement with the trustees of Dartmouth College, the corresponding author has developed and distributed the [HowsYourHealth.org](http://www.HowsYourHealth.org) website for the collection of data in this report. In this respect, the corresponding author has a conflict of interest. The authors have no other potential conflicts of interest.

REFERENCES

- 1 Kohn LT, Corrigan JM, Donaldson MS, eds. *To err is human*. Washington DC: National Academy Press, 1999.
- 2 Milch CE, Salem DN, Pauker SG, et al. Voluntary electronic reporting of errors and adverse events. *J Gen Intern Med* 2006;**21**:165-70.
- 3 Elder NC, Vonder Meulen, Cassidy A. The identification of medical errors by family physicians during outpatient visits. *Ann Fam Med* 2005;**2**:125-9.
- 4 Agoritas T, Bovier P, Perneger T. Patient reports of undesirable events during hospitalization. *J Gen Intern Med* 2005;**20**:922-8.
- 5 Michel P, Quenon JL, de Sarasqueta AM, et al. Comparison of three methods for estimating rates of adverse events and rates of preventable events in acute care hospitals. *BMJ* 2004;**328**:199-202.
- 6 Weingart SN, Pagovich O, Sands DZ, et al. What can hospitalized patients tell us about adverse events? *J Gen Intern Med* 2005;**20**:830-6.
- 7 Burroughs TE, Waterman AD, Gallagher TH, et al. Patient concerns about medical errors in emergency departments. *Acad Emerg Med* 2005;**12**:57-64.
- 8 Adams RE, Boscarino JA. A community survey of medical errors in New York. *Int J Qual Health Care* 2004;**16**:353-62.
- 9 Nelson EC, Conger B, Douglass R, et al. Functional health status levels of primary care patients. *J Gen Intern Med* 1983;**249**:3331-8.
- 10 Nelson EC, Landgraf JM, Hays RD, et al. The functional status of patients: how can it be measured in physicians' offices? *Med Care* 1990;**28**:1111-26.
- 11 Magari ES, Hamel MB, Wasson JH. An easy way to measure quality of physician-patient interactions. *J Ambulatory Care Manage* 1998;**21**:27-33.
- 12 Wasson JH, Stukel TA, Weiss JE, et al. A randomized trial of using patient self-assessment data to improve community practices. *Eff Clin Pract* 1999;**2**:1-10.
- 13 Ahles T, Wasson J, Seville, Johnson D, et al. A controlled trial of methods for managing pain in primary care patients with or without co-occurring psychosocial problems. *Ann Fam Med* 2006;**4**:1-10.
- 14 Wasson JH, James C. Implementation of a web-based interaction technology to improve the quality of a city's health care. *J Ambulatory Care Manage* 2001;**24**:1-12.
- 15 Luce P, Phillips J, Benjamin R, et al. Technology to support community health alliances. *J Ambulatory Care Manage* 2004;**27**:399-407.
- 16 Wasson JH, Johnson DJ, Benjamin R, et al. Patients report positive impacts of collaborative care. *J Ambul Care Manag* 2006;**29**:201-8.
- 17 Leape LL, Berwick DM, Bates DW. What practices will most improve safety: evidence based medicine meets patient safety. *JAMA* 2002;**288**:501-7.
- 18 Stryer D, Clancy C. Patients' safety. Progress is elusive because culture in health care has not changed. *BMJ* 2005;**330**:553-4.
- 19 Kuzel AJ, Woolf SH, Gilchrist VJ, et al. Patient reports of preventable problems in primary health care. *Ann Fam Med* 2004;**2**:333-40.