

Clarity and strength of implications for practice in medical journal articles: an exploratory analysis

Joanne Lynn,¹ Alessia P Owens,² Jean M Bartunek³

► An additional appendix is published online only. To view this file please visit the journal online (<http://qualitysafety.bmj.com>).

¹Colorado Foundation for Medical Care, Chevy Chase, Maryland, USA

²Howard University School of Social Work, Washington, DC, USA

³Department of Organization Studies, Boston College, Chestnut Hill, Massachusetts, USA

Correspondence to

Joanne Lynn, Colorado Foundation for Medical Care, 2318 Ashboro Drive, Chevy Chase, MD 20815, USA; drjoannelynn@gmail.com

Accepted 3 November 2010

ABSTRACT

Objective: To examine how leading clinical journals report research findings, aiming to assess how they frame their implications for medical practice and to compare that literature's patterns with those of the management literature.

Data Source: Clinically relevant research articles from three leading clinical journals (*N Engl J Med*, *JAMA*, and *Ann Intern Med*).

Methods: Review of wording of a sequential sample from 2010, with categorisation, comparison among journals, and comparison with management literature.

Results: Clinical journals usually state that one approach did or did not differ from another approach (35 of 51 articles, 68.6%), but they recommended a specific course of action ('therefore, x should be done') in just 25.5%. One article gave instruction on how to implement the changes. Two-thirds of the reports called for further research. Half used tentative language. Management research articles nearly always specified who should use the information and drew from over 60 types of potential users, whereas the clinical literature named the audience in only 23.5% of clinicians.

Conclusions: Authors and editors of the clinical literature could test being more clear and direct in presenting implications of research findings for practice, including stating when the findings do not justify changes in practice.

INTRODUCTION

In March 2010, Bartunek and Rynes¹ reviewed the leading professional literature in organisational management, showing that a minority of papers provide concrete and clear advice arising from the research and that most used tentative language in conveying their advice. We were curious as to the patterns and practices in the medical literature.

Clarke and Chalmers² in 1998 reviewed 26 randomised clinical trials and found only two articles that deliberately integrated their new

findings into existing literature. Lewis³ suggested that passive voice and unclear comments limited the usefulness of articles to healthcare decision makers. Lucas⁴ confirmed that medical journals usually give tentative result statements. Tunis and colleagues⁵ reported that clinical trial research regularly fails to give decision makers enough information to make well-informed decisions, perhaps because investigators have little incentive to reach out to practising clinicians or policy makers.

This paper examines recent research reports in leading medical journals to gain insight into how authors interpret the implications of their findings for medical practice.

METHODS

Journal and article selection

Since we were seeking to characterise broadly influential journals, we selected those with the highest 'Journal Impact Factor (JIF)', which tallies the citations of articles in one journal in all professional journals.⁶ The top three US medical journals by (2004–2009) JIF score are *The New England Journal of Medicine* (NEJM), *The Journal of the American Medical Association* (JAMA) and *Annals of Internal Medicine* (Annals). The subscribers are mostly clinicians; for example, JAMA reports 100% of their circulation goes to physicians, medical students, hospitals and firms associated with the medical profession.⁷

Two authors (APO and JL) reviewed the table of contents for each issue of each journal, starting with the first edition of 2010, until we had identified more than 20 articles with titles that appeared to report primary research relevant to the medical care of human beings, excluding reviews, editorials and other items.



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://qualitysafety.bmj.com/site/about/unlocked.xhtml>

Review of selected articles

Each author independently reviewed each article. We excluded articles that did not report findings relevant to medical treatment (because they reported basic science, epidemiological description or methodological work). Our reviews focused upon identifying statements that articulated implications for practice in the Abstract or Discussion sections. In addition, we identified statements that the findings had implications for future research or for public health. We did not code guidelines or medical advice that did not arise from the current research. Annals publishes a short comment from the Editor, which we discuss separately.

After initial review, we developed categories to code and all three authors independently reviewed each article, identified the appropriate text and coded it. We resolved any coding disagreements by discussion. The Bartunek and Rynes article¹ identified that managers prefer straightforward, clearly implementable prescriptions for practice.⁸ Thus, we examined the text that articulated implications for practice, being careful to identify tentative language (eg, words such as 'may', 'speculate', and 'potentially'). We also tallied prescriptive language, such as 'ought' and 'must'. We tabulated findings that applied to a subset of situations, which we called 'contingencies'. Finally, we coded whether the article identified who (eg, the physician), if anyone, should take account of the findings.

We then compared the three journals, using overall χ^2 tests; and, when appropriate, we compared pairs of journals using Bonferroni's correction for significance level. We also compared the medical literature with the management literature¹ using χ^2 tests.

We tabulated text findings on Excel and managed coding and analyses with SPSS (PASW V.18).

RESULTS

The cohort and coding

Aiming to sample ≥ 20 articles per journal, we reviewed NEJM through 28 February 2010; JAMA through 10 March 2010 and Annals through 6 April 2010. The initial cohort totalled 65 articles, with 21 from NEJM, 21 from JAMA and 23 from Annals. The initial review excluded 14 articles, leaving a cohort of 51 that reported research that could be important to medical practice: 18 from NEJM,^{9–26} 12 from JAMA^{27–38} and 21 from Annals.^{39–59}

Figure 1 illustrates the derivation of this analytic cohort.

Thirty-five articles gave advice that 'X was better than Y with regard to Z' or 'X is no better than Y with regard to Z'. These statements often hedged: 'X seemed to be better than Y' or 'X may prove to be better than Y'. These statements did not intimate what other considerations might affect the practitioner's decision as to




Figure 1 Article cohort derivation.

whether to change. We called these statements 'Minimal Advice', and 28 articles gave only this kind of phrasing. We identified four additional patterns for stating the implications of findings for practice: 'practitioner should consider' (labelled 'Consider'); 'patient should be informed' (labelled 'Inform'); 'practitioner should do' (labelled 'Act'); and 'practitioner should do and here's how' (labelled 'Technical Assistance').

Statements about implications for practice

One article by Ray *et al*⁵⁵ stated factual findings with no evaluative statement at all. This article reported results from a large observational study that assessed two kinds of medications. The outcomes were complex, showing competing gains and risks. The information would be useful for a practitioner to consider, though the article did not explicitly state that conclusion. The information could not have yielded advice to prefer one treatment strategy, except perhaps to inform the patient.

Table 1 shows the number and proportion of the other 50 articles that provided the five kinds of advice, overall and for each journal and figure 2 illustrates the overlap among the kinds of advice given.

Table 1 Proportion of articles providing each kind of implication for practice, by journal

Journals	Categories, N=51				
	Minimal advice, n (%)	Consider, n (%)	Inform, n (%)	Act, n (%)	Technical assistance, n (%)
NEJM (n=18)	16 (88.9)	4 (22.2)	1 (5.6)	1 (5.6)	0
JAMA (n=12)	10 (83.3)	1 (8.3)	1 (1.9)	2 (16.7)	0
Annals (n=21)	9 (42.9)	6 (28.6)	0	10 (47.6)	1 (4.8)
Total* (N=51)	35 (68.6)	11 (21.6)	2 (3.9)	13 (25.5)	1 (2.0)

*Totals sum to >100% because categories are not mutually exclusive.

Eleven articles called on someone to ‘consider’ acting in response to the findings. In no case did the article give additional advice as to how that person might weigh the issues or what other considerations should enter into the deliberation. For instance, ‘Early resumption of low-dose aspirin therapy with proton pump inhibitors in patients with bleeding ulcers and cardiovascular disease should be considered’.³⁹

Two articles called for informing the patient as part of the implementation of findings, one concerning influenza immunisation¹¹ and one concerning feeding tube placement.²⁹ For example, ‘As the current pandemic unfolds, pregnant and postpartum women should be counseled about the importance of vaccination’.¹¹

Thirteen articles called for specific action, generally stating ‘The physician should...’. For example, ‘Our findings also suggest that compression ultrasonography might be considered for patients with symptomatic SVT

at presentation to evaluate the extent of the thrombosis and diagnose potential DVT, that physicians should suspect and test for pulmonary embolism in patients with suggestive symptoms...’.⁵⁰ The first clause counts as a ‘Consider’ statement; the second as an ‘Act’ statement.

One article provided a unique approach. Goldstein *et al*⁵⁴ published a ‘Brief Communication’ in Annals that gave research findings and stated clear directives for action and also provided an on-line appendix with details about implementation.


Often, the authors appear to assume that the reader would know the implications for practice and do not state them. Van der Gaag *et al*,¹⁶ for example, found that ‘Routine perioperative biliary drainage in patients undergoing surgery for cancer of the pancreatic head increases the rate of complications’. However, the article does not state the obvious implication for practice: do not do the drainage procedure in this setting.

Additional attributes of statements about implications of the research

We describe various attributes of how the articles provide implications for practice in table 2.

Two-thirds (66.7%) of the articles called for future research, usually using a broad and brief formulation. For example, Goebel *et al*⁴⁵ claimed: ‘Additional studies are needed to determine which patients are likely to benefit and which IVIG doses and schedules are most effective’. In 9.8% of the cases, the articles stated an implication for public health actions such as developing guidelines or including certain services in funding. Nachega *et al*,⁴¹ for example, called for monitoring adherence to anti-retroviral therapy to be part of ‘the package of care in anti-retroviral therapy programs in all settings’, which would require action by funders and governments.

Forty-seven per cent of the articles used tentative language in stating their implications for practice. For example, ‘Our findings suggest that the use of morphine during trauma care may reduce the risk of subsequent development of PTSD after serious injury’.¹⁴ As in this example, many used two or more tentative terms. Sometimes, the writing strongly avoided being direct, for example, ‘Therefore, we believe that prescribing an



* One article in each group also met criteria for ‘Inform’

^ One article in this group also met criteria for ‘Technical Assistance’

Figure 2 Number of articles by category.

Table 2 Proportion of articles with particular attributes by journal and overall

Journals	Implication for research, n (%)	Implication for public health, n (%)	Tentative language present, n (%)	Subset language present, n (%)	Prescriptive language present, n (%)	Audience specified, n (%)
NEJM (n=18)	10 (55.6)	4 (22.2)	8 (44.4)	3 (16.7)	4 (22.2)	2 (11.1)
JAMA (n=12)	11 (91.7)	0	6 (50.0)	3 (25.0)	4 (33.3)	2 (16.7)
Annals (n=21)	13 (61.9)	1 (4.8)	10 (47.6)	9 (42.9)	12 (57.1)	8 (38.1)
Total (N=51)	34 (66.7)	5 (9.8)	24 (47.1)	15 (29.4)	20 (39.2)	12 (23.5)

NSAID to treat an asymptomatic postoperative pericardial effusion should no longer be advised'.⁴⁷ The reader might reasonably wonder whether this is a matter of faith that requires belief and also who should no longer be advising whom.

About 39% of the articles had some prescriptive language. For example, 'Until then, clinical practice should not be guided by (point-of-care) platelet function testing'.³¹ Contingent (subset) statements were present in about 29%. For example, 'In short, most women contemplating estrogen plus progestin therapy for the relief of menopausal symptoms should not expect protection against CHD'.⁴⁹

Each journal's website makes a general statement about intended readers: NEJM targets 'medical researchers', JAMA claims 'physician readers' and Annals focuses on 'practising physicians'. The stated audience does not include non-medical professionals, laypersons or patients. Perhaps this strong assumption about the readership explains why only 23.5% of the articles specify the people whose actions should reflect the reported findings. Authors did specify physicians in 10% of articles and healthcare providers or clinicians in 8%.

Differences among journals

We analysed the data in tables 1 and 2 for differences among journals and only two comparisons met conventional standards for significance: Annals was significantly less likely to publish 'Minimal Advice' than NEJM ($\chi^2=8.925$, $df=2$, $p=0.006$). Conversely, Annals was significantly more likely than NEJM to recommend action ($\chi^2=8.469$, $df=2$, $p=0.005$).

In Annals, the Editors' note provides a very short summary of findings and cautions. Sometimes, as in Meurin *et al*,⁴⁷ the Editors give a stronger recommendation than the authors did: 'Physicians should stop prescribing NSAIDs for postoperative pericardial effusion because these agents have no or only small beneficial effects' by the Editors and 'In patients with pericardial effusion after cardiac surgery, diclofenac neither reduced the size of the effusions nor prevented late cardiac tamponade' by the authors. On the other hand, sometimes the Editors' comments worded the authors' conclusions more tentatively than the authors,

as in Schaer *et al*,⁴³ where the Editors said 'In hypertensive patients, mechanisms other than lowering blood pressure may be important causes of atrial fibrillation', while the authors claimed 'In hypertensive patients, long-term receipt of ACE inhibitors, ARBs, or beta-blockers reduces the risk for atrial fibrillation compared with receipt of calcium-channel blockers'.

Comparison with management research reporting

Management researchers advocating changes in practice virtually always stated their claims in the active voice. In contrast, in 33% of cases medical journals used passive voice, for example, 'New strategies to reduce the risk of transmission of HIV-1 are needed for HIV-1-serodiscordant couples'.²⁶ Implications for practice sections in the management literature used tentative language significantly more (74%) than the medical literature (47%), ($\chi^2=17.48$, $df=1$, $p<0.001$). The management literature (55%) was also significantly more likely to use prescriptive language than the medical literature (39%), ($\chi^2=4.59$, $df=1$, $p=0.04$). However, contingent advice was similar in frequency in the management (38%) and medical (29%) literature.

DISCUSSION

This project examined how three influential US medical journals present implications of research findings for practice. Most research reports provide a summary of the findings with a flat descriptive statement that a clinical outcome is better or worse with the tested intervention (68.6%). The one article that did not give this level of advice had findings that would not support doing so. Only about one-fourth (25.5%) of the reports gave explicit recommendations for action and only one article provided instruction as to how to implement the recommended actions. The language in about half of the articles was tentative (47.1%) and a general call for further research was common (66.7%). Reporting in the three journals was generally similar, but as a group they differed from the management journals in including fewer calls to undertake action, and in not specifying who should act.

In this project we studied only a small sample of articles in a confined time frame, which limited the study's

power and generalisability. Furthermore, our review did not examine editorials or journal review services, which often serve to put the findings of single studies into a broad context and provide substantial advice to the practitioner as to how to interpret the results.

While the target for implications for practice in medical journals was either not named or was specified only as 'physicians' or 'clinicians', the management journals virtually always named the target audience. Bartunek and Rynes identified more than 60 categories of individuals or groups to whom articles addressed their implications.¹ The clinician audience for these medical journals may be so obvious as to need no specification.

The project raised some intriguing questions as to optimal reporting. In the management literature assessment, the authors advocated more statements of the implications for practice¹; in contrast, many of the articles we sampled from clinical literature had methodological limitations that precluded strong statements guiding practitioners towards changed practices. The findings could often have needed replication or could need to be weighed along with other considerations in making individual treatment decisions. Strong statements about medical practice might more often be justified in systematic reviews, or clinical practice guidelines based on extensive literature reviews. One well-known example is the US Preventive Services Task Force, which provides authoritative reviews and grades both the strength of the evidence and of their recommendations.^{60 61}

However, clarity in assessing and reporting the impact of new findings is desirable, even when the report should have limited impact. Authors can clearly state what their work has added to the body of evidence and what its implications for practice are, even if the clearly stated advice would often be that a reasonable practitioner should not change anything yet. This proposal would operationalise for all research reports the contention of Clarke and Chalmers² regarding randomised clinical trials. They advocated that each report should state, in the 'discussion' section, how the new findings reshape the prior body of evidence concerning the topic.

Editors should also consider undertaking formal tests of whether authors can clearly and succinctly state implications for practice, perhaps by providing explicit guidelines on grades of evidence and levels of recommendation and whether such explicit and consistent reporting helps readers. If past research and the new data support a strong recommendation for implementation, then the article should explicitly state the major counter-considerations or contingencies and often can provide 'technical assistance' for implementation. If the authors hold that the findings are not yet reliable enough for action, they could state that

judgement clearly, along with their opinion regarding who should undertake what research or other actions.

Medical research generally builds in small increments and few articles make a dramatic difference to what practitioners should do. However, our findings suggest that the current system for reporting the implications of single clinical studies deserves serious re-examination by authors, editors and readers. Stating the strength of evidence and recommendations for action in those studies as directly, unambiguously and consistently as possible seems to us likely to prove helpful in translating their results into the most appropriate actions.

Acknowledgements The authors are grateful to the Health Foundation in the UK which brought two of the authors together to examine the epistemology of quality improvement in April 2010, a meeting which sparked our interest in the topic. The authors are grateful to the National Academy of Social Insurance, which supported Alessia P Owens as a Somers Scholar, enabling her to take the lead in this work. The authors also acknowledge the substantial help from anonymous reviewers and the editor, Dr Frank Davidoff.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

1. Bartunek JM, Rynes SL. The construction and contributions of "implications for practice": what's in them and what might they offer? *Academy of Management Learning and Education* 2010;9:100–17.
2. Clarke M, Chalmers I. Discussion sections in reports of controlled trials published in general medical journals: islands in search of continents? *JAMA* 1998;280:280–2.
3. Lewis S. Toward a general theory of indifference to research-based evidence. *J Health Serv Res Policy* 2007;12:166–72.
4. Lucas BP, Evans AT, Reilly BM, *et al*. The impact of evidence on physicians' inpatient treatment decisions. *J Gen Intern Med* 2004;19:402–9.
5. Tunis SR, Stryer DB, Clancy CM. Practical clinical trials: increasing the value of clinical research for decision making in clinical and health policy. *JAMA* 2003;290:1624–32.
6. Journal Citation Reports® for 2009: Thomson Reuters, 2010. http://thomsonreuters.com/content/corporate/PDF/IR/annReports/tr2009_annReport.pdf (accessed 25 Oct 2010).
7. BPA Worldwide. Business publication circulation statement for the 6 month period ended January 2010. *JAMA* 2010;903:1–5. <http://pubs.ama-assn.org/misc/JAMAcirc.pdf> (accessed 29 Jul 2010).
8. Bazerman MH. Conducting influential research: the need for prescriptive implications. *Acad Manage Rev* 2005;30:25–31.
9. Bode LG, Kluytmans JA, Wertheim HF, *et al*. Preventing surgical-site infections in nasal carriers of *Staphylococcus aureus*. *N Engl J Med* 2010;362:9–17.
10. Darouiche RO, Wall MJ Jr, Itani KM, *et al*. Chlorhexidine-Alcohol versus Povidone-Iodine for Surgical-Site Antisepsis. *N Engl J Med* 2010;362:18–26.
11. Louie JK, Acosta M, Jamieson DJ, *et al*. Severe 2009 H1N1 influenza in pregnant and postpartum women in California. *N Engl J Med* 2010;362:27–35.
12. Libster R, Bugna J, Coviello S, *et al*. Pediatric hospitalizations associated with 2009 pandemic influenza A (H1N1) in Argentina. *N Engl J Med* 2010;362:45–55.
13. Mansfield AJ, Kaufman JS, Marshall SW, *et al*. Deployment and the use of mental health services among U.S. Army wives. *N Engl J Med* 2010;362:101–9.
14. Holbrook TL, Galarneau MR, Dye JL, *et al*. Morphine use after combat injury in Iraq and post-traumatic stress disorder. *N Engl J Med* 2010;362:110–17.
15. Griffiths CE, Strober BE, van de Kerkhof P, *et al*. Comparison of ustekinumab and etanercept for moderate-to-severe psoriasis. *N Engl J Med* 2010;362:118–28.
16. van der Gaag NA, Rauws EA, van Eijck CH, *et al*. Preoperative biliary drainage for cancer of the head of the pancreas. *N Engl J Med* 2010;362:129–37.

17. Delaere P, Vranckx J, Verleden G, *et al*. Tracheal allotransplantation after withdrawal of immunosuppressive therapy. *N Engl J Med* 2010;362:138–45.
18. Lowy I, Molrine DC, Leav BA, *et al*. Treatment with monoclonal antibodies against *Clostridium difficile* toxins. *N Engl J Med* 2010;362:197–205.
19. Madhi SA, Cunliffe NA, Steele D, *et al*. Effect of human rotavirus vaccine on severe diarrhea in African infants. *N Engl J Med* 2010;362:289–98.
20. Richardson V, Hernandez-Pichardo J, Quintanar-Solares M, *et al*. Effect of rotavirus vaccination on death from childhood diarrhea in Mexico. *N Engl J Med* 2010;362:299–305.
21. Bakker JJ, Verhoeven CJ, Janssen PF, *et al*. Outcomes after internal versus external tocodynamometry for monitoring labor. *N Engl J Med* 2010;362:306–13.
22. Patel NC, Hertel PM, Estes MK, *et al*. Vaccine-acquired rotavirus in infants with severe combined immunodeficiency. *N Engl J Med* 2010;362:314–19.
23. Kappos L, Radue EW, O'Connor P, *et al*. A placebo-controlled trial of oral fingolimod in relapsing multiple sclerosis. *N Engl J Med* 2010;362:387–401.
24. Cohen JA, Barkhof F, Comi G, *et al*. Oral fingolimod or intramuscular interferon for relapsing multiple sclerosis. *N Engl J Med* 2010;362:402–15.
25. Giovannoni G, Comi G, Cook S, *et al*. A placebo-controlled trial of oral cladribine for relapsing multiple sclerosis. *N Engl J Med* 2010;362:416–26.
26. Celum C, Wald A, Lingappa JR, *et al*. Acyclovir and transmission of HIV-1 from persons infected with HIV-1 and HSV-2. *N Engl J Med* 2010;362:427–39.
27. Hemmelgarn BR, Manns BJ, Lloyd A, *et al*. Relation between kidney function, proteinuria, and adverse outcomes. *JAMA* 2010;303:423–9.
28. O'Brien PE, Sawyer SM, Laurie C, *et al*. Laparoscopic adjustable gastric banding in severely obese adolescents: a randomized trial. *JAMA* 2010;303:519–26.
29. Teno JM, Mitchell SL, Gozalo PL, *et al*. Hospital characteristics associated with feeding tube placement in nursing home residents with advanced cognitive impairment. *JAMA* 2010;303:544–50.
30. Paynter NP, Chasman DI, Pare G, *et al*. Association between a literature-based genetic risk score and cardiovascular events in women. *JAMA* 2010;303:631–7.
31. Breet NJ, van Werkum JW, Bouman HJ, *et al*. Comparison of platelet function tests in predicting clinical outcome in patients undergoing coronary stent implantation. *JAMA* 2010;303:754–62.
32. Jones AE, Shapiro NI, Trzeciak S, *et al*. Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial. *JAMA* 2010;303:739–46.
33. Fowkes FG, Price JF, Stewart MC, *et al*. Aspirin for prevention of cardiovascular events in a general population screened for a low ankle brachial index: a randomized controlled trial. *JAMA* 2010;303:841–8.
34. Wunsch H, Guerra C, Barnato AE, *et al*. Three-year outcomes for Medicare beneficiaries who survive intensive care. *JAMA* 2010;303:849–56.
35. Brookhart MA, Schneeweiss S, Avorn J, *et al*. Comparative mortality risk of anemia management practices in incident hemodialysis patients. *JAMA* 2010;303:857–64.
36. Briel M, Meade M, Mercat A, *et al*. Higher vs lower positive end-expiratory pressure in patients with acute lung injury and acute respiratory distress syndrome: systematic review and meta-analysis. *JAMA* 2010;303:865–73.
37. Loeb M, Russell ML, Moss L, *et al*. Effect of influenza vaccination of children on infection rates in Hutterite communities: a randomized trial. *JAMA* 2010;303:943–50.
38. Segev DL, Muzaale AD, Caffo BS, *et al*. Perioperative mortality and long-term survival following live kidney donation. *JAMA* 2010;303:959–66.
39. Sung JJ, Lau JY, Ching JY, *et al*. Continuation of low-dose aspirin therapy in peptic ulcer bleeding: a randomized trial. *Ann Intern Med* 2010;152:1–9.
40. Yeh HC, Duncan BB, Schmidt MI, *et al*. Smoking, smoking cessation, and risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med* 2010;152:10–17.
41. Nachega JB, Leisegang R, Bishai D, *et al*. Association of antiretroviral therapy adherence and health care costs. *Ann Intern Med* 2010;152:18–25.
42. Hayward RA, Krumholz HM, Zulman DM, *et al*. Optimizing statin treatment for primary prevention of coronary artery disease. *Ann Intern Med* 2010;152:69–77.
43. Schaer BA, Schneider C, Jick SS, *et al*. Risk for incident atrial fibrillation in patients who receive antihypertensive drugs: a nested case-control study. *Ann Intern Med* 2010;152:78–84.
44. Dunn KM, Saunders KW, Rutter CM, *et al*. Opioid prescriptions for chronic pain and overdose: a cohort study. *Ann Intern Med* 2010;152:85–92.
45. Goebel A, Baranowski A, Maurer K, *et al*. Intravenous immunoglobulin treatment of the complex regional pain syndrome: a randomized trial. *Ann Intern Med* 2010;152:152–8.
46. Schnoll RA, Patterson F, Wileyto EP, *et al*. Effectiveness of extended-duration transdermal nicotine therapy: a randomized trial. *Ann Intern Med* 2010;152:144–51.
47. Meurin P, Tabet JY, Thabut G, *et al*. Nonsteroidal anti-inflammatory drug treatment for postoperative pericardial effusion: a multicenter randomized, double-blind trial. *Ann Intern Med* 2010;152:137–43.
48. Rodriguez J, Jiang R, Johnson WC, *et al*. The association of pipe and cigar use with cotinine levels, lung function, and airflow obstruction: a cross-sectional study. *Ann Intern Med* 2010;152:201–10.
49. Toh S, Hernandez-Diaz S, Logan R, *et al*. Coronary heart disease in postmenopausal recipients of estrogen plus progestin therapy: does the increased risk ever disappear? A randomized trial. *Ann Intern Med* 2010;152:211–17.
50. Decousus H, Quere I, Presles E, *et al*. Superficial venous thrombosis and venous thromboembolism: a large, prospective epidemiologic study. *Ann Intern Med* 2010;152:218–24.
51. Harvey WF, Yang M, Cooke TD, *et al*. Association of leg-length inequality with knee osteoarthritis: a cohort study. *Ann Intern Med* 2010;152:287–95.
52. Baggish AL, Hutter AM Jr, Wang F, *et al*. Cardiovascular screening in college athletes with and without electrocardiography: A cross-sectional study. *Ann Intern Med* 2010;152:269–75.
53. Wheeler MT, Heidenreich PA, Froelicher VF, *et al*. Cost-effectiveness of preparticipation screening for prevention of sudden cardiac death in young athletes. *Ann Intern Med* 2010;152:276–86.
54. Goldstein N, Carlson M, Livote E, *et al*. Brief communication: Management of implantable cardioverter-defibrillators in hospice: A nationwide survey. *Ann Intern Med* 2010;152:296–9.
55. Ray WA, Murray KT, Griffin MR, *et al*. Outcomes with concurrent use of clopidogrel and proton-pump inhibitors: a cohort study. *Ann Intern Med* 2010;152:337–45.
56. Yeh JM, Nekhlyudov L, Goldie SJ, *et al*. A model-based estimate of cumulative excess mortality in survivors of childhood cancer. *Ann Intern Med* 2010;152:409–17, W131–8.
57. Trifirò G, Gambassi G, Sen E, *et al*. Association of community-acquired pneumonia with antipsychotic drug use in elderly patients. *Ann Intern Med* 2010;152:418–25, W139–40.
58. Rosenquist JN, Murabito J, Fowler JH, *et al*. The spread of alcohol consumption behavior in a large social network. *Ann Intern Med* 2010;152:426–33, W141.
59. Stein PD, Chenevert TL, Fowler SE, *et al*. Gadolinium-enhanced magnetic resonance angiography for pulmonary embolism: a multicenter prospective study (PIOPED III). *Ann Intern Med* 2010;152:434–43, W142–3.
60. U.S. Preventive Services Task Force. Procedure Manual. AHRQ Publication No. 08-05118-EF, 2008. <http://www.uspreventiveservicestaskforce.org/uspstf08/methods/procmanual.htm> (accessed 28 Oct 2010).
61. Harris RP, Helfand M, Woolf SH, *et al*; Methods Work Group, Third US Preventive Services Task Force. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med* 2001;20:21–35.