



OPEN ACCESS

Fragmented care in the era of limited work hours: a plea for an explicit handover curriculum

Max Wohlaue

Correspondence to

Dr Max Wohlaue,
Department of Surgery,
University of Colorado School
of Medicine, 12631 E. 17th
Ave., C302, Aurora, CO
80045, USA;
max.wohlaue@ucdenver.edu

Accepted 8 October 2012

ABSTRACT

The current challenge in surgical residency education, preserving a sense of accountability in an era of limited work hours and multiple transitions of care, can be accomplished through the promotion of effective handovers. Approaches include increasing the transparency of handovers, enhancing the 'situational awareness' of individuals involved, and incorporating narratives and social interactions, and expanding common ground among professionals who collectively have responsibility for patient care. Done well, handovers can preserve the physician-patient relationship and provide a continuity bridge during a vulnerable time for patients.

As a surgical resident in the era of duty hour limits (in my case the US Accreditation Council for Graduate Medical Education's 2011 duty hour standards), I have seen the changes in the healthcare system and bear witness to the importance of effective handovers. In an era when healthcare depends on the effective exchange of information among individuals and teams who collectively provide care for patients across time and different healthcare settings, the importance of the handover is growing. In the European Community, which has work hour restrictions for all healthcare professionals, and where the healthcare system separates primary care and the hospital setting, handover skills are of paramount importance. Taking pride in continuity of care expressed by an effective handover elevates patients above their disease processes and considers their individuality, despite multiple professionals and settings involved and the increasing reliance on technology.

While standardisation can be an effective way to deal with the increasingly complex hospital environment, adopting checklists and tools without teaching residents the principles of risk management, systems thinking and effective handovers does not solve the

problem.¹ Impressing the importance of handovers by making them part of the explicit educational curriculum is the answer.

PRINCIPLES OF EFFECTIVE HANDOVERS

Handovers are complex and multifaceted events that occur daily in the inpatient setting and between the hospital and the patient's primary care setting.²⁻³ Handovers within the hospital span a variety of settings, including transitions from the emergency department to the operating room, the operating room to the intensive care unit, and the intensive care unit to the inpatient floor. In all of these contexts, the purpose of the handover is to safely effect a transition of care; promoting patient continuity and safety in a specific physical and cultural environment. A handover is more than the passive transfer of information; a *decisive feature* is the transfer of responsibility for the patient's care among the participants.⁴ Data elements important to effective handovers include patient identification, a diagnostic summary, current condition and disease trajectory, plan of care, a list of care task (the to-do list), and a plan for anticipated events and contingencies.⁵ An effective handover is a bidirectional conversation whatever the setting or context, and includes active involvement of the sender and the receiver.⁶⁻⁸

FRAGMENTATION OF CARE: THE UNINTENDED CONSEQUENCE OF SHIFT WORK AND DISTRIBUTION OF MEDICAL SERVICES

Physicians several decades ago often cared for their patients across the different settings, following them through the acute care admission, to home and upon return to the ambulatory clinic for follow-up. The growing

boundaries between specialisation and the separation of primary care and hospital services have resulted in the responsibility for continuity of care being shared by many professionals. Work hour restrictions, designed to improve patient care, have resulted in less time with the patients, shorter shifts and successive transfers of responsibility for the patient among multiple professionals, further contributing to fragmented patient care.⁹ When a patient presents to the hospital with an illness, it takes time for physicians to understand the precipitating events, gather medical records, order tests, etc. This process is called *sensemaking*, the activity of gathering information to accomplish a task.¹⁰ In the hospital setting sensemaking involves bringing the incoming provider up to speed to facilitate a smooth transition of care at admission or during shift change.¹¹

The initial investment to 'process' the patient in the inpatient context can be a substantial one. Understanding a patient's disease trajectory helps his or her physicians anticipate likely and less expected contingent events and deal with the inherent uncertainty of taking care of an ill patient. If the initial time and effort spent can be considered the *investment*, then establishing a plan of care is the *return on investment*. This may be a reason why some physicians who trained in the era before work hour restrictions look fondly upon these experiences. They feel that the time invested getting to know a patient was time well spent. Conversely, a frequent complaint is that junior residents trained in the current work-restricted era do not know their patients. A similar complaint emerged from the healthcare professionals participating in qualitative studies conducted as part of the European HANDOVER Project.¹² Primary care physicians lamented that time constraints and disconnectedness between the hospital and the primary care settings that cared for patients following discharge resulted in haphazard information exchanges that did not allow them to know their patients. Healthcare professionals in the hospital cited time constraints as reasons they were not able to make a full 'investment' in their patient at discharge, such as a consultation that assesses the patients' preparedness for discharge, and the provision of timely and detailed information to the primary care physician. Another study found that transfers between intensive care and a regular nursing floor were haphazard, and critical personnel were routinely absent during this important exchange.¹³ The root causes of these common complaints about discontinuity of care are threefold: the investment of getting to know the patient is perceived to be too great; time constraints prevent a meaningful exchange of information that allows for sensemaking; and/or the relatively short exposure contributes to a lack of emotional connection with the patient, leading to a relative disinterest on the part of the healthcare professional in gaining more detailed knowledge and building trust.

IMPORTANCE OF SURGICAL HANDOVERS IN THE TWENTY-FIRST CENTURY

Traditionally, surgical care continuity has involved minimising handovers, often at the cost of prolonging hours at work in the name of professionalism.¹⁴ Generations of surgical trainees have provided continuity irrespective of hours worked. However, in an era of limits on residents' duty hours, sick patients can be very inconvenient to a resident nearing the end of a shift. The issue is continuity of care, and experts on the topic argue that even a fatigued intern with detailed knowledge about a patient may be able to render more appropriate care than a well rested one who is less familiar with the patient and also busy admitting other patients to the hospital.¹⁵ How can this be possible? Transitions of care create vulnerability, and these moments of vulnerability may contribute to patient adverse events, near misses and duplicative work by healthcare providers.¹⁶ However, the benefits of effective handovers are plentiful, including providing a fresh perspective when insight is greatly needed and more humanistic and supportive learning conditions.¹⁷

In spite of their proven importance and ubiquity, most surgery residency programmes lack an explicit handover curriculum. I learnt how to perform handovers from a chief resident who understood their importance and taught me an implicitly effective method. While this approach worked on an individual basis, the approach to handover education in residency training is hit or miss and unreliable. Had I been on service with a different chief resident, I might have learned a less effective technique or received no guidance at all. Rather than accepting a haphazard approach to educating physicians about handovers, this fundamental skill in the twenty-first century should be formally integrated into the residency programme curriculum and assessed, potentially using entrustable professional activities.¹⁸

Learning about shift changes when patients are at high risk offers insight into improving transitions of care. One approach involves increasing the transparency of handovers, and enhancing the 'situational awareness' of the individuals involved.¹⁹ Recent research has shown that incorporating narratives, social interactions, and cultural norms can be enabling factors in designing an effective handover, and a growing body of literature aims to improve communication.²⁰ As the vast majority of communication breakdowns are verbal rather than written, a computerised sign-out list is a valuable adjunct to the process.^{21–23} Other options to increase the reliability of handovers include overlapping shifts and expanding common ground by formally including other caregivers into the exchange of handover information,¹⁰ or by reducing abbreviated 'coded' communication

terms when third parties may become unintended recipients of handover information shared with data systems or the electronic health record.²⁴ We also understand more and need to be prepared for addressing how various factors such as personality and experience of incoming or outgoing provider could impact the effectiveness of communication during handover.⁷

In conclusion, if a Spartan ethos enabled the twentieth century resident to live at the hospital and provide uninterrupted patient care, then team work and effective handovers are the modern tools empowering the twenty-first century resident to maintain a sense of accountability and patient ownership in an era of increasingly fragmented care.²⁵ Handovers preserve the physician–patient relationship and provide a continuity bridge during a vulnerable time for patients. It is time for physician training programmes and education programmes for other health professions with responsibility for patient care to make training in handovers an explicit part of their curriculum, elevating this essential twenty-first century skill to the heightened status to which it belongs.

Competing interests None.

Provenance and peer review Commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 3.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/3.0/>

REFERENCES

1. Symons NR, Wong HW, Manser T, *et al.* An observational study of teamwork skills in shift handover. *Int J Surg* 2012;10:355–9.
2. Horwitz LI, Krumholz HM, Green ML, *et al.* Transfers of patient care between house staff on internal medicine wards: a national survey. *Arch Intern Med* 2006;166:1173–7.
3. Hesselink G, Schoonhoven L, Barach P, *et al.* Improving patient handovers from hospital to primary care: a systematic review. *Ann Intern Med* 2012;157:417–28.
4. Cohen MD, Hilligoss PB. The published literature on handoffs in hospitals: deficiencies identified in an extensive review. *Qual Saf Health Care* 2010;19:493–7.
5. Borowitz SM, Waggoner-Fountain LA, Bass EJ, *et al.* Adequacy of information transferred at resident sign-out (in-hospital handover of care): a prospective survey. *Qual Saf Health Care* 2008;17:6–10.
6. Philibert I. Use of strategies from high-reliability organisations to the patient hand-off by resident physicians: practical implications. *Qual Saf Health Care* 2009;18:261–6.
7. Gibson SC, Ham JJ, Apker J, *et al.* Communication, communication, communication: the art of the handoff. *Ann Emerg Med* 2010;55:181–3.
8. Sharit J, McCane L, Thevenin DM, *et al.* Examining links between sign-out reporting during shift changeovers and patient management risks. *Risk Anal* 2008;28:983–1001.
9. Laine C, Goldman L, Soukup JR, *et al.* The impact of a regulation restricting medical house staff working hours on the quality of patient care. *JAMA* 1993;269:374–8.
10. Weick KE, Sutcliffe KM, Obstfeld D. Organizing for high reliability: processes of collective mindfulness In Research in Organizational Behavior; R. Sutton and B. Staw (Eds.); 1999; Greenwich, CT: JAI, 81–124.
11. Sharma N. Sensemaking handoff: when and how? *Proc Am Soc Inf Sci Technol* 2008;45:1–12.
12. Hesselink G, Flink M, Olsson M, *et al.* Are patients discharged with care? A qualitative study of perceptions and experiences of patients, family members and care providers. *BMJ Qual Saf*, December 2012.
13. Toccafondi G, Albolino S, Tartaglia R, *et al.* Analysis of the interface between high acuity and low acuity care: the collaborative communication model for patient handover. *BMJ Qual Saf*, December 2012.
14. Philibert I, Leach DC. Re-framing continuity of care for this century. *Qual Saf Health Care* 2005;14:394–6.
15. Petersen LA, Brennan TA, O'Neil AC, *et al.* Does housestaff discontinuity of care increase the risk for preventable adverse events? *Ann Intern Med* 1994;121:866–72.
16. Horwitz L, Moin T, Krumholz H, *et al.* Consequences of inadequate sign-out for patient care. *Arch Intern Med* 2008;168:1755–60.
17. Patterson ES, Roth EM, Woods DD, *et al.* Handoff strategies in settings with high consequences for failure: lessons for health care operations. *Int J Qual Health Care*. 2004;16:125–32.
18. ten Cate Th J O, Young J. The patient handover as an entrustable professional activity: adding meaning in teaching and practice. *BMJ Qual Saf*, December 2012.
19. Patterson ES. Communication strategies from high-reliability organizations: translation is hard work. *Ann Surg* 2007;245:170–2.
20. Patterson ES, Wears RL. Patient handoffs: standardized and reliable measurement tools remain elusive. *Jt Comm J Qual Patient Saf* 2010;36:52–61.
21. Greenberg CC, Regenbogen SE, Studdert DM, *et al.* Patterns of communication breakdowns resulting in injury to surgical patients. *J Am Coll Surg* 2007;204:533–40.
22. Van Eaton EG, Horvath KD, Lober WB, *et al.* Organizing the transfer of patient care information: the development of a computerized resident sign-out system. *Surgery* 2004;136:5–13.
23. Wohlaer MV, Rove KO, Pshak TJ, *et al.* The computerized rounding report: implementation of a model system to support transitions of care. *J Surg Res* 2012;172:11–7.
24. Patterson E. Technology support of the handover: promoting observability, flexibility and efficiency. *BMJ Qual Saf*, December 2012.
25. Van Eaton EG, Horvath KD, Pellegrini CA. Professionalism and the shift mentality: how to reconcile patient ownership with limited work hours. *Arch Surg* 2005;140:230–5.