

decrease in medical costs by 64%, lost days by 69% and treatment delays by 77%. Texas adopted EBTG in 2006; Total costs have declined by 50%, patients are recovering more quickly, more providers are willing to treat these patients, opioid abuses have declined and many states are looking to adopt this model. Other state outcomes and essential elements needed in an EBTG will be discussed.

050 DEVELOPING EVIDENCE-BASED CLINICAL PRACTICE GUIDELINES FOR PRIMARY CARE PHYSIOTHERAPY: A LOCAL INITIATIVE

¹S Bernhardtsson, ^{1,2}M Larsson. ¹Region Västra Götaland, Primary Care, Gothenburg, Sweden; ²Sahlgrenska Academy at Gothenburg University, Institute of Neuroscience and Physiology, Gothenburg, Sweden

10:1136/bmjqs-2013-002293.81

Background In primary care physiotherapy in Sweden, there is a lack of evidence-based clinical practice guidelines to support clinicians and patients in clinical decision making. A local initiative to develop and implement evidence-based clinical practice guidelines for PTs in primary care was initiated by a regional health authority.

Objectives To develop evidence-based clinical practice guidelines for physiotherapy treatment of low back pain, neck pain, and subacromial pain, for use in primary care physiotherapy in Sweden.

Methods The guideline development was performed by a project team of five primary care physiotherapists in a systematic process that was guided by a 7-step guideline development model by Grol et al. We performed systematic database searches, critical appraisal of the evidence base using GRADE, and formulated evidence-based practice recommendations.

Results The guideline format follows recommendations from AGREE II. The guidelines consist of a brief summary on the first page; a brief introduction to the topic with up-to-date information on definition of the condition, prevalence and prognosis; recommendations on patient management according to strength of evidence; a discussion section, and a detailed reference list. Detailed search strategy and search results, summaries of the body of evidence, recommended outcome measures, and patient information were provided in appendices to the guidelines.

Discussion Rigorous guideline development methodology was considered important to get clinicians' attention and approval - a requisite for regular use. Clinicians welcomed the initiative.

Implications for Guideline Developers It is important for clinician acceptance that guidelines are brief, to the point and relevant.

051 CAN WE AUTOMATICALLY PRODUCE GENERIC DECISION AIDS FOR THE CLINICAL ENCOUNTER DIRECTLY FROM GRADE GUIDELINE RECOMMENDATIONS? EXPERIENCE FROM THE SHARE-IT PROJECT

¹T Agoritsas, ²L Brandt, ²A Anja Heen, ²A Kristiansen, ³P Alonso-Coello, ⁴E Akl, ¹I Neumann, ¹K Tikkinen, ⁵V Montor, ¹G Guyatt, ²P Vandvik. ¹Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, Canada; ²Norwegian Knowledge Centre for the Health Services, Oslo, Norway; ³Iberoamerican Cochrane Center (CIBERESP), Hospital de Sant Pau, Barcelona, Spain; ⁴Department of Medicine, State University of New York City, Buffalo, USA; ⁵Knowledge and Evaluation Research Unit, Mayo Clinic, Rochester, MN, USA

10:1136/bmjqs-2013-002293.82

Background Although decision aids (DA) can help to communicate evidence to patients, their production is time consuming, often not based on the best available evidence or rapidly outdated. Linking trustworthy guidelines and DA for shared-decision making could both overcome these limitations and enhance guideline dissemination.

Objectives To test the feasibility of automatically translating any recommendations from GRADE guidelines into generic and interactive DA accessible on tablet computers for clinicians and their patients in the clinical encounter.

Methods As part of the DECIDE project, we developed a framework for translating components of GRADE into DA, following the International Patient Decision Aid Standards. Using a recently published guideline, we implemented that framework in our MAGIC (Making Grade the Irresistible Choice) application - a prototype electronic guideline tool and publication platform that can automatically display recommendations in multilayered presentation formats.

Results Our prototype was able to automatically translate a large number of GRADE recommendations and their supporting evidence into electronic and interactive DA. Preliminary results of user-testing in real patient-clinician interactions suggest that these DA can be used at the point of care to discuss estimates of treatment effects for patient relevant outcomes, confidence in estimates, burden of treatment, and cost issues.

Discussion This study provides a proof-of-concept that components of GRADE recommendations can be interactively displayed in generic tools for interactive shared-decision making in a wide range of treatment alternatives.

Implications for Guideline Developers/Users Our electronic DA offer promising opportunities to disseminate guidelines at the point of care.

052 TOOLS FOR COMMUNICATING MAIN MESSAGES CONVEYED BY CLINICAL PRACTICE GUIDELINES (CPGS) - MAKING PATIENT RESOURCES USER FRIENDLY

¹C Moga, ²D Angus, ¹D Chojecki, ¹C Harstall, ¹J Moffatt, ³P Taenzer. ¹Institute of Health Economics, Edmonton, Canada; ²Alberta Innovates - Health Solutions, Edmonton, Canada; ³University of Calgary, Faculty of Medicine, Calgary, Canada

10:1136/bmjqs-2013-002293.83

Background The Alberta Ambassador Guideline Adaptation Programme developed two CPGs on the management of low back pain and headache in primary care. Various guideline companion documents were developed for patients.

Objectives To review the process and outcomes of involving patients/public in the development of guideline resource documents.

Methods Focus groups of different compositions and using various sampling strategies were called upon: self-selected members from national patient pain association (patient information sheets); members of the public who serve on a standing Lay Committee (patient information sheets, comic book); purposive recruitment by market research company (comic book). Participants answered questions related to the content, format, and presentation of patient resources using different interview methods.

Results Discussion themes included wordiness of documents, language accessibility, use of logos, and trustworthiness. Multiple suggestions for improvements that were sensitive to the needs of participants were considered before publication of final products. The comic book was not published.