

086 **GUIDELINES ADAPTATION IN LOW AND MIDDLE INCOME COUNTRIES (LMIC): RESULTS AND LESSONS LEARNT FROM AN 8-YEAR-CAPACITY BUILDING (CB) EXPERIENCE IN ARGENTINA (2005–2012)**

<sup>1,2</sup>M Esandi, <sup>1</sup>M De Luca, <sup>1</sup>Z Ortiz. <sup>1</sup>National Academy of Medicine, Ciudad Autónoma de Buenos Aires, Buenos Aires, Argentina; <sup>2</sup>National Southern University, Bahía, Blanca, Argentina

10:1136/bmjqs-2013-002293.117

**Background** Capacity building (CB) is an essential element for guideline adaptation in the context of LMIC.

**Objectives** To describe the approach and results from a CB process for guidelines adaptation implemented by the National Academy of Medicine (NAM) in Argentina between 2005 and 2012.

**Methods** The CB process is described on the basis of the matrix of capacity-building strategies. Duration, objectives, entities targeted and results through the different stages of the CB process are provided.

**Results** The CB process has been supported on a “learning by doing” approach, and comprised 2 stages: a local capacity development stage (2005–2008) and a knowledge transfer (KT) stage (2008–2012). As a result of the 1st stage, 120 health professionals were involved during the adaptation process; 3 guidelines were produced and a guide to adapt guidelines was published. KT started in 2008 and was initially performed through e-learning courses targeted to individuals. In 2009, a strategy based on continuous online support through a virtual campus and workshops and targeted to institutions was adopted. Four institutions were involved: 1 did not progress; 1 completed the whole process and published an evidence-based guideline and 2 are ongoing.

**Implications for Guideline Developers/Users** In the context of LMIC, CB processes based on the “learning by doing” approach and focused to institutions seem to be more appropriate although challenging: not only technical capacities have to be built, but also those related to human resources management, group-working and use of Internet resources. Different level of achievement of these capacities could explain the results observed alongside the CB process implemented by NAM.

087 **CAPACITIES, PRACTICES AND PERCEPTIONS OF EVIDENCE-BASED PUBLIC HEALTH IN EUROPE**

<sup>1</sup>A Jansen, <sup>2</sup>F Forland, <sup>3</sup>L Murajda, <sup>1</sup>J Latham. <sup>1</sup>ECDC, Stockholm, Sweden; <sup>2</sup>KIT, Stockholm, Sweden; <sup>3</sup>EPIET, Stuttgart, Germany

10:1136/bmjqs-2013-002293.118

Evidence-based methodologies are used to synthesise systematic high-quality evidence and were first applied in clinical practice. Evidence-based public health, however, is still in its early stages. The European Centre for Disease Prevention and Control sought the insight of European organisations working in the field of public health on current practices, capacities, perceptions and predictions of evidence-based public health. A survey was sent to 76 organisations. A response rate of 36% was achieved, representing 27 organisations from 16 countries. Systematic reviews were the most commonly offered service, followed by health technology assessments and rapid assessments. Fifty-four per cent of respondents believed that evidence-based methodologies were poorly integrated into public health. The main perceived barriers to the further development of evidence-based public

health included ‘lack of formalised structure or system’, ‘resource constraints’ ‘lack of understanding of evidence-based methodologies by policy makers’ and ‘lack of data’. Nevertheless, 81% of respondents believed that evidence-based methodologies will play an increasingly important role in public health in future. However, several barriers need to be overcome. Consistent frameworks and consensus on best practices were identified as the most pressing requirements. Steps should be taken to address these barriers and facilitate integration and ultimately public health policies.

088 **COHORT OF CLINICAL PRACTICE GUIDELINES FROM THE SPANISH NATIONAL GUIDELINE PROGRAMME: A SURVIVAL ANALYSIS**

<sup>1</sup>L Martínez García, <sup>1</sup>A Sanabria, <sup>1</sup>R David, <sup>2</sup>L Barajas, <sup>2,3</sup>P Díaz del Campo Fontecha, <sup>3,4</sup>M Estrada Sabadell, <sup>4,5</sup>I Etxeandia Ikobaltzeta, <sup>5,6</sup>E García Álvarez, A Louro González <sup>6,7</sup>F Salcedo Fernández, <sup>7,8</sup>M Trujill, <sup>1</sup>P Alonso-Coello. <sup>1</sup>Iberoamerican Cochrane Centre - IIB Sant Pau, Barcelona, Spain; <sup>2</sup>Centro Cochrane Iberoamericano, Barcelona, Spain; <sup>3</sup>Subdirección General de Tecnología e Innovación Sanitarias, Madrid, Spain; <sup>4</sup>Catalan Agency for Health Information, Assessment and Quality (CAHIAQ), Barcelona, Spain; <sup>5</sup>OSTEBA-OsasunTecnologienEbaluazioa/Basque OfficeHealth Technology Assessment, Vitoria, Spain, <sup>6</sup>Servicio Gallego de Salud (SERGAS), A Coruña, Spain; <sup>7</sup>GuíaSalud-Aragon Institute of Health Sciences, Zaragoza, Spain; <sup>8</sup>Fundación Canaria de Investigación y Salud (FUNCIS) Las Palmas de Gran Canaria, Spain

10:1136/bmjqs-2013-002293.119

**Background** Clinical Practice Guidelines (CPGs) recommendations need to be updated to maintain their validity.

**Objectives** To provide empirical estimates of the average time after which CPGs recommendations become obsolete.

**Methods** We developed a strategy to assess the validity of CPG recommendations, which included assessing their validity by surveying clinical experts, updating the literature search, screening references by pertinence and matching them with recommendations, and identifying pertinent, relevant and key references, and potential changes in each recommendation. A convenience sample of four CPGs was selected. We piloted our strategy in 20% of the recommendations from these CPGs (feasibility test) and we estimated our sample size. We performed a survival analysis and considered a CPG outdated when more than 20% of recommendations needed to be updated.

**Results** The four CPGs included 250 recommendations. A total of 39.133 (range 3.343–14.784) references were identified in the exhaustive literature search in a time frame of 3–5 years. The feasibility test identified 16 key references updating 8 recommendations. The number of recommendations required for the study was 113. A total of 674 references were marked as pertinent to these recommendations.

**Discussion** We developed a rigorous, replicable evaluation strategy to assess the validity of recommendations and estimate CPG obsolescence. Full final results will be present at the GIN meeting.

**Implications for Guideline Developers/Users** Our work is relevant for guideline developers because it provides information about the expected validity of CPGs recommendations.

089 **AN INTERNATIONAL COMPARISON OF OCCUPATIONAL GUIDELINES FOR THE MANAGEMENT OF MENTAL DISORDERS**

<sup>1</sup>M Joosen, <sup>1</sup>E Brouwers, <sup>1</sup>K van Beurden, <sup>2</sup>B Terluin, <sup>3</sup>J van der Klink, <sup>4</sup>J Verbeek, <sup>5</sup>H Eguchi, <sup>6</sup>J Woo, <sup>1</sup>J van Weeghel. <sup>1</sup>Tranzo, Tilburg University, Tilburg School of Social