



OPEN ACCESS

# How sensitive are avoidable emergency department attendances to primary care quality? Retrospective observational study

Beth Parkinson <sup>1</sup>, Rachel Meacock <sup>1</sup>, Kath Checkland,<sup>1</sup> Matt Sutton <sup>1,2</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2020-011651>).

<sup>1</sup>Health, Organisation, Policy and Economics (HOPE) Group, Centre for Primary Care & Health Services Research, University of Manchester, Manchester, UK

<sup>2</sup>Melbourne Institute of Applied Economic and Social Research, Faculty of Business and Economics, The University of Melbourne, Parkville, Victoria, Australia

## Correspondence to

Mrs Beth Parkinson;  
beth.parkinson@manchester.ac.uk

Received 28 May 2020

Revised 9 October 2020

Accepted 14 October 2020

Published Online First

3 November 2020



Check for updates

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

**To cite:** Parkinson B, Meacock R, Checkland K, et al. *BMJ Qual Saf* 2021;**30**:884–892.

## ABSTRACT

**Background** Improvements in primary care quality are often proposed as a solution to rises in emergency department (ED) attendances. However, there is little agreement on what constitutes an avoidable attendance, and the relationship between primary care quality and ED demand remains poorly understood.

**Objective** To estimate the size of the associations between primary care quality and volumes of ED attendances classified as avoidable.

**Methods** Retrospective observational study of all attendances at EDs in England during 2015/2016, applying three definitions of avoidable attendance. We linked practice-level counts of attendances to seven measures of primary care access, patient experience and clinical quality for 7521 practices. We used count data regressions to associate attendance counts with levels of quality. We then calculated proportions of attendances associated with levels of primary care quality below the national average.

**Results** Attendance volumes were negatively related to three of the seven quality measures. Incidence rate ratios (IRRs) for all attendances associated with 10 percentage-point differences in quality were 0.987 for clinical quality and 0.987 for easy telephone access and 0.978 for ability to get an appointment. These associations were relatively stronger for narrower definitions of avoidable attendances (for the narrowest definition, IRRs=0.966, 0.976 and 0.934, respectively) but represented fewer attendances in absolute terms. 341 000 (2.4%) attendances were associated with levels of primary care quality below the national average in 2015/2016.

**Conclusion** ED attendances are sensitive to primary care quality, but magnitudes of these associations are small. Attendances are much less responsive to differences in primary care quality than indicated by estimates of the prevalence of avoidable attendances. This may explain the failure of initiatives to reduce attendances through primary care improvements.

## INTRODUCTION

Increases in the volume of emergency department (ED) attendances have put pressure on healthcare services worldwide, resulting in adverse impacts on the quality and safety of patient care.<sup>1</sup> It is

often suggested that many attendances are avoidable and that these patients could have been more appropriately treated elsewhere, most often in a primary care setting.<sup>2</sup> Several interventions have been introduced to manage demand for EDs, with improving accessibility of primary care services during and outside of normal working hours a key strategy.<sup>3</sup> Such policies assume that a proportion of current attendances could have been avoided if access to primary care was improved. However, the relationship between the quality of primary care services and demand for ED services is poorly understood.

A recent systematic review of studies assessing the effect of primary care practice-level factors on ED attendances found that continuity of care and access to general practice were generally associated with fewer ED attendances, while evidence relating to quality of care was limited and inconclusive.<sup>4</sup> However, the included studies often focused on specific geographical areas or patient populations. To replicate the findings from their systematic review, Tammes and colleagues conducted both cross-sectional<sup>5</sup> and longitudinal studies<sup>6</sup> using data from across England. Both studies showed some evidence of an association between accessibility of primary care and the volume of attendances that were self-referred and did not result in an admission. However, another recent study in England found no association between patient experience of general practice and the volume of ED attendances.<sup>7</sup>

There have been many attempts to calculate the proportion of ED attendances that are ‘avoidable’, yet there is

currently no agreement regarding what constitutes an avoidable attendance, and no standardised definition exists.<sup>2 8 9</sup> Definitions generally attempt to capture attendances that are non-urgent, are self-presentations, receive care that could be delivered by a general practitioner (GP) and do not require admission.<sup>2</sup> However, different criteria have been used to identify these patients empirically.

The lack of agreement as to what constitutes an avoidable ED attendance likely explains the inconsistency in the attempts to link primary care quality with the volume of avoidable attendances. It also means there is no reliable estimate of the magnitude of the problem, that is, the number of ED attendances that are sensitive to primary care quality.

In this paper, we aimed to estimate the size of the associations between primary care quality and volumes of ED attendances classified as avoidable. Furthermore, we quantify the proportion of ED attendances that are associated with levels of primary care quality below the national average. We extend the existing literature by using three definitions of avoidable attendances and several aspects of primary care quality. Thus, we examine comprehensively the sensitivity of the volume of ED attendances to primary care quality.

## METHODS

### Study setting: emergency care in England

In England, patients register with a single general practice of their choice to access appointments for both routine and urgent primary care. GPs act as gatekeepers to specialist services provided by hospitals. While general practices do provide out-of-hours services, most appointments occur during normal working hours.

If patients perceive they need more urgent or more specialised services than are immediately available from their GP, they can obtain urgent care from an ED or walk-in centre. There are three categories of ED. The majority of emergency attendances are at type 1 departments, which are consultant-led with full resuscitation facilities open 24 hours a day.<sup>10</sup> Patients cannot book an appointment, but there is a maximum waiting time target of 4 hours, which should be met for 95% of patients.<sup>11</sup>

General practices are paid by capitation. EDs are reimbursed per attendance under an activity-based financing scheme.<sup>12</sup> The costs of attendances are borne by local third-party payer organisations, Clinical Commissioning Groups. General practices thus have little direct financial incentive to avoid patients attending EDs, although most Clinical Commissioning Groups will intervene if practice rates fall outside of norms.

Primary and secondary care are provided free at point of use, funded through taxation. This makes England a useful setting for examining the relationship

between primary care quality and ED attendances, as it is not confounded by variations in insurance coverage.

## Data

Outcome variables: 'avoidable' ED attendances

We used patient-level data on attendances at EDs in England for the financial year 1 April 2015–31 March 2016. These records are taken from Hospital Episode Statistics (HES), an administrative data set capturing all hospital activity in England.<sup>13</sup> We focused on attendances at major (type 1) EDs. To control for the availability of other emergency care services in the area, we included the practice-level rate of attendance at type 2 (single specialty centres), type 3 (minor injury units) and type 4 (walk-in centres) EDs.

We identified three definitions which had been used to retrospectively identify potentially avoidable attendances in previous studies in England.<sup>14</sup> They vary in the number and strictness of the criteria applied:

1. 'Self-referred discharges': patients who were 'self-referred and sent home (discharged) (from the ED) either with no follow-up or follow-up treatment to be provided by their GP' as in Tammes *et al.*<sup>5 6</sup>
2. 'GP treatable': 'first attendance with some recorded treatments or investigations, all of which may have been reasonably provided by a GP, followed by being sent home or to GP care'. This is currently used in the online data tool on 'unnecessary Accident and Emergency (A&E) attendances' published by NHS Digital, the national NHS information organisation in England.<sup>15</sup>
3. 'No treatment and discharged': patients who were 'self-referred; was an initial (rather than follow-up) attendance for this condition; received no investigation and either no treatment or 'guidance/advice only'; and were sent home with either no follow-up or follow-up with primary care' as in McHale *et al.*<sup>16</sup>

We generated indicators to identify attendances belonging to these three definitions by mapping the characteristics described in the studies to variables in HES attendance records. Online supplemental appendix 1 provides details of this process.

We then used the general practice identifier to generate practice-level counts of each type of attendance, as well as total attendances (online supplemental appendix 2 provides details of the aggregation process).

### General practice quality variables

We included explanatory variables measuring realised access, continuity of care, clinical quality and patient experience, obtained from the General Practitioner Patient Survey (GPPS) and the Quality and Outcomes Framework (QOF). Online supplemental appendix 3 provides a description of all explanatory variables, including sources and how variables were constructed.

The GPPS is a biannual postal survey conducted on behalf of NHS England. Participants are sampled from practice registration lists to collect patients'

views and experiences of the services provided.<sup>17</sup> We use data from the July 2016 publication that contains data collected from July to September 2015 and from January to March 2016. Across these two waves, 2 148 791 surveys were distributed with a response rate of 38.9% (836 312 returned surveys).<sup>18</sup>

From the GPPS, we obtained proportions of patients who: would recommend the practice to someone who has just moved to their local area; reported good overall experience of their GP surgery; found it easy to get through to their GP surgery on the phone; were able to make an appointment to see or speak to someone when they wanted to; were able to make a same day appointment; and, among patients with a preferred GP, were able to speak to their preferred GP always or a lot of the time.

To ensure representativeness of practices' registered populations, responses were published weighted by characteristics of the practice list.<sup>18</sup> The only measure for which we did not use the weighted version is 'the proportion of patients that were able to speak to their preferred GP always or a lot of the time', as the weighted version of this variable is missing for practices with low response rates to this question (n=397). We examined the sensitivity of the results to using the weighted versus unweighted version of this measure. We also controlled for the GPPS response rate at each practice.

The QOF is a national scheme that provides financial incentives for general practices to improve their quality of care.<sup>19</sup> Practices score points for achieving targets on a range of indicators. The QOF contains three domains: clinical, public health, and public health additional services. We used the proportion of clinical QOF points achieved in 2015/2016 as an indicator of clinical quality.

We tested for multicollinearity among practice quality and accessibility variables using the variance inflation factor. For all variables, these inflation factors were under the commonly used threshold for high multicollinearity of 10.<sup>20</sup>

#### Practice and patient population characteristics

We included several characteristics of practices and their registered populations as potential predictors.

We adjusted the attendance counts for the size of registered patient populations and control for the following characteristics: age distribution (proportion 0–4 years, 5–15 years, 45–54 years, 55–64 years, 65–74 years, 75–84 years and 85+ years),<sup>21</sup> income deprivation,<sup>22</sup> proportion unemployed<sup>23</sup> and proportion whose ethnic group was UK white.<sup>23</sup> We also controlled for differences in the health of the registered populations by including prevalence rates of five health conditions, obtained from the QOF<sup>24</sup>: chronic obstructive pulmonary disease, heart failure, asthma, atrial fibrillation, and coronary heart disease.

We controlled for practice size using the average number of GPs working at the practice over the year.<sup>25</sup> We also included several geographical characteristics: a rural indicator,<sup>26</sup> indicators for 13 NHS regions<sup>27</sup> and distance between each GP practice and the nearest type 1 ED in England as an estimate of patients' travel time to the nearest ED.

A full description of how all explanatory variables were constructed and sourced is provided in online supplemental appendix 3. Data on the outcome measures and practice factors were merged using GP practice identifier codes (see online supplemental appendix 2 for full details of the merging process).

#### Statistical methods

We used negative binomial regressions to model the three different counts of avoidable ED attendances as a function of primary care quality, population and practice characteristics. We also modelled the count of all ED attendances for comparison. The negative binomial model is a generalisation of the Poisson model, including an additional parameter to account for overdispersion, a common feature of healthcare use data.

We included practice list size as the exposure variable; its coefficient is constrained to unity so that the coefficients in the model can be interpreted as the effect on the annual rates of attendance per registered patient. Coefficients are reported as incidence rate ratios (IRRs), meaning that values greater than 1 indicate a positive association with the volume of attendances, while values less than 1 indicate a negative association. For continuous variables, the IRR can be interpreted as a 100(IRR–1)% difference in the rate of attendances per one-unit difference in the explanatory variable, while for categorical variables (eg, deprivation decile), this would be a 100(IRR–1)% difference relative to the reference group. We scaled all the quality variables so that one-unit differences represent differences of 10 percentage points.

We used robust SEs, clustered at the Clinical Commissioning Group level. Since each model considers the annual rate of attendance per registered patient and coefficients are reported as IRRs, we are able to directly compare the magnitudes of the coefficients across different definitions of attendance. The coefficients represent relative measures of the risk of attendance incidents associated with measures of primary care quality. If the three definitions of 'avoidable' attendances accurately capture patients that are either divertible or deferrable to primary care, then these measures should be more sensitive than total attendance volumes to primary care quality. This should be both in terms of the magnitude and statistical significance of the coefficients.

We then explored the sensitivity of ED attendance volumes to primary care quality by estimating the number of attendances that were associated with below-average levels of the seven primary care quality

variables. This is calculated as the difference between the numbers of attendances predicted by the models using (1) the observed levels of quality recorded for each practice and (2) replacing values of the quality variables below the national average with the national average. These estimates therefore represent the annual number of attendances in England associated with below-average primary care quality.

#### Sensitivity analyses

We ran three sensitivity checks. First, we removed practices within 20 km of the borders with Scotland or Wales ( $n=323$ ) as we were unable to measure attendances at EDs outside of England. Second, we included ratings from the Care Quality Commission (CQC), the independent regulator of health and social care services in England, as an additional measure of primary care quality. CQC inspections are carried out at least every 5 years, and so ratings do not necessarily correspond to quality during the year of the analysis. Third, we tested the sensitivity of the results to using weighted rather than unweighted 'proportions of patients able to see their preferred GP'.

## RESULTS

The mean number of ED attendances was 257 per 1000 registered patients (table 1). On average, there were 94 attendances per 1000 registered patients classified as avoidable using definition 1 (self-referred discharges); 34 classified as avoidable using definition 2 (GP treatable), and 16 classified as avoidable using definition 3 (no treatment and no follow-up attendances).

Average practice achievement of clinical QOF points was 96% (table 1). On average, 78% of patients would recommend their practice; 85% reported good overall experience of their GP surgery; 62% were able to see their preferred GP; 73% found it easy to get through to their practice on the phone; 85% were able to make an appointment; and 37% of patients were able to get a same-day appointment.

The associations between attendance volumes and practice and population characteristics are presented in table 2. A higher proportion of registered patients aged 0–4 years, higher levels of deprivation and higher unemployment are all associated with higher rates of attendances across all four measures. Distance to the nearest ED, the response rate to the GPPS and the rate of attendance at other emergency care facilities are negatively associated with ED attendances across all four measures. Higher proportions of registered patients aged 75–84 and 85 and over are associated with higher rates of total attendances but not with any of the three measures of avoidable attendances.

The associations between attendance volumes and the primary care quality measures from the same models are presented in table 3. The proportion of patients reporting easy phone access and the proportion of patients able to make an appointment

both demonstrate a significant negative association with rates of attendances across all four attendance measures. The magnitude of the relationship between attendance rates and these two measures of access increases as the strictness of the definition of avoidable attendances increases, from total attendance volumes down to 'no treatment and no follow-up' avoidable attendances. This indicates that measures of avoidable attendances are more sensitive to primary care access than total attendance volumes. Total attendances and 'self-referred discharged' avoidable attendances were also found to be negatively associated with primary care quality as measured by clinical QOF scores. The magnitude of the association is stronger for 'self-referred discharged' avoidable attendances than for total ED attendances.

We estimate that levels of primary care quality below the national average are associated with 2.48% of total ED attendances, 3.70% of self-referred discharged attendances, 4.10% of GP treatable attendances and 5.26% no treatment and no follow-up attendances. However, when we estimate the absolute number of attendances that could potentially be avoided if levels of quality that were below the average were raised to the average, our models generate values of 346 334 total attendances, 187 815 self-referred discharged attendances, 74 464 GP treatable attendances and 44 002 no treatment and no follow-up attendances. While the relative association between primary care quality and attendances strengthens as stricter definitions of avoidability are applied, our estimates of the absolute number of attendances that could potentially be avoided suggests some attendances outside of these definitions of avoidable are nevertheless sensitive to primary care quality.

The results were robust to three sensitivity analyses (online supplemental appendix 4). Removing practices within 20 km of the borders with Scotland or Wales did not notably change the magnitude or implications of the results. CQC inspection ratings were not found to be statistically significantly associated with the volume of attendances. Using the weighted proportion of patients able to see their GP did not change the results.

## DISCUSSION

### Summary of findings

Improvements in primary care quality are often proposed as a solution to the problem of continually increasing ED attendances. However, the relationship between quality of primary care and the demand for ED services is poorly understood. We find that regardless of whether we examine all attendances or narrow down to definitions of an avoidable attendance, lower attendance volumes are associated with better access to primary care services in the form of ability to get an appointment and ease of telephone contact. This relationship exists even after controlling for a wide range

**Table 1** Descriptive statistics on the 7521 general practices included in the analysis

Rates of ED attendance per 1000 population	Mean	SD	25th percentile	Median	75th percentile
Total ED attendances	256.5	89.8	195.1	247.6	311.4
Definition 1: self-referred discharged (Tammes <i>et al</i> ) <sup>5,6</sup>	93.6	50.4	56.7	86.9	124.0
Definition 2: GP treatable (NHS Digital) <sup>15</sup>	33.7	23.6	17.2	28.2	44.0
Definition 3: no treatment and no-follow up (McHale <i>et al</i> ) <sup>16</sup>	15.5	12.8	6.8	12.5	20.5
Primary care quality measures					
Proportion of clinical Quality and Outcomes Framework points achieved	0.954	0.070	0.943	0.977	0.994
Proportion of patients who would recommend the practice	0.774	0.124	0.701	0.793	0.869
Proportion of patients reporting good overall experience of GP surgery	0.852	0.092	0.803	0.870	0.921
Proportion of patients who could see their preferred GP	0.620	0.173	0.500	0.630	0.750
Proportion of patients reporting easy phone access	0.731	0.168	0.625	0.761	0.865
Proportion of patients able to get an appointment to see or speak to someone	0.847	0.081	0.801	0.860	0.906
Proportion of patients who were able to get same day appointment	0.367	0.145	0.258	0.350	0.458
Control variables					
Size of the registered population	7459	4444	4086	6629	9965
Proportion of registered population aged 0–4 years	0.060	0.016	0.049	0.058	0.069
Proportion of registered population aged 5–15 years	0.126	0.027	0.112	0.124	0.139
Proportion of registered population aged 16–44 years	0.388	0.086	0.333	0.373	0.426
Proportion of registered population aged 45–54 years	0.143	0.021	0.134	0.146	0.156
Proportion of registered population aged 55–64 years	0.113	0.026	0.097	0.116	0.130
Proportion of registered population aged 65–74 years	0.093	0.035	0.068	0.094	0.117
Proportion of registered population aged 75–85 years	0.055	0.022	0.039	0.055	0.069
Proportion of registered population aged 85+ years	0.022	0.012	0.014	0.021	0.029
Income deprivation score	0.176	0.110	0.087	0.150	0.245
Proportion of registered patients of UK white ethnicity	0.764	0.254	0.649	0.879	0.949
Proportion of registered patients unemployed	0.051	0.047	0.017	0.039	0.072
Response rate to General Practitioner Patient Survey	0.412	0.110	0.330	0.419	0.496
Atrial fibrillation prevalence (proportion of registered patients)	0.020	0.254	0.011	0.017	0.021
COPD prevalence (proportion of registered patients)	0.027	0.669	0.013	0.018	0.024
Asthma prevalence (proportion of registered patients)	0.078	1.660	0.051	0.059	0.067
Heart failure prevalence (proportion of registered patients)	0.009	0.138	0.005	0.007	0.009
Coronary heart disease prevalence (proportion of registered patients)	0.042	0.853	0.025	0.032	0.040
Distance from GP practice to nearest type one ED (km)	6.858	6.677	2.503	4.497	9.014
Rate of attendance at type 1–3 EDs (per 10 patients)	0.830	1.012	0.105	0.388	1.253
Number of GPs	5.349	3.388	3.000	4.915	7.099
Variables included in sensitivity analyses					
Rated good or outstanding by CQC on first inspection	0.828	0.378	1.000	1.000	1.000
Weighted proportion of patients who could see their preferred GP	0.590	0.171	0.469	0.595	0.715

COPD, chronic obstructive pulmonary disease; CQC, Care Quality Commission; ED, emergency department; GP, general practitioner.

of practice and population characteristics. However, we do not find attendance volumes to be associated with patient experience or the proportion of patients able to see their preferred GP, which is often interpreted as a measure of care continuity.

There is a lack of agreement in the literature as to what constitutes an avoidable ED attendance, and this has resulted in estimates of the proportion of ED use classified as inappropriate ranging from 10% to 90%.<sup>8</sup>

We compared across three commonly used definitions of an avoidable attendance to examine whether some were more sensitive than others to primary care quality. We found that the magnitudes of the associations with indicators of primary care quality were larger for stricter definitions of avoidable attendances. Stricter measures of avoidable attendances are therefore more sensitive to primary care quality in relative terms.

**Table 2** Associations between ED attendance volumes and practice and population characteristics

	All attendances	Self-referred discharged attendances	GP treatable attendances	No treatment and no follow-up attendances
	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)
Proportion of patients aged 0–4†	1.444*** (1.316 to 1.584)	1.501*** (1.258 to 1.791)	1.390** (1.118 to 1.728)	1.444** (1.096 to 1.904)
Proportion of patients aged 5–15†	0.966 (0.910 to 1.025)	1.086 (0.982 to 1.202)	1.091 (0.956 to 1.247)	1.242* (1.050 to 1.469)
Proportion of patients aged 45–54†	1.159** (1.061 to 1.266)	1.089 (0.923 to 1.285)	1.074 (0.895 to 1.290)	0.973 (0.782 to 1.211)
Proportion of patients aged 55–64†	0.990 (0.903 to 1.085)	1.050 (0.893 to 1.236)	0.966 (0.809 to 1.154)	1.023 (0.816 to 1.283)
Proportion of patients aged 65–74†	0.979 (0.885 to 1.082)	0.990 (0.804 to 1.218)	0.824 (0.627 to 1.082)	0.831 (0.577 to 1.195)
Proportion of patients aged 75–84†	1.194** (1.059 to 1.346)	1.126 (0.889 to 1.426)	1.349 (0.971 to 1.872)	1.260 (0.862 to 1.844)
Proportion of patients aged 85 and over†	1.396*** (1.207 to 1.615)	0.946 (0.768 to 1.165)	0.980 (0.651 to 1.474)	0.984 (0.663 to 1.462)
Income deprivation score†	1.044*** (1.030 to 1.057)	1.029* (1.006 to 1.052)	1.052** (1.021 to 1.084)	1.045** (1.013 to 1.078)
Rural practice	1.055*** (1.023 to 1.089)	1.108*** (1.043 to 1.176)	1.065 (0.996 to 1.139)	1.070 (0.987 to 1.160)
Number of GPs	0.999 (0.997 to 0.002)	0.995 (0.990 to 1.000)	0.998 (0.992 to 1.005)	0.993 (0.986 to 0.000)
Proportion of patients of UK white ethnicity†	1.016* (1.001 to 1.031)	1.020 (0.990 to 1.052)	1.025 (0.990 to 1.062)	1.062* (1.014 to 1.113)
Proportion of patients unemployed†	1.101*** (1.060 to 1.144)	1.072*** (1.031 to 1.115)	1.137*** (1.072 to 1.205)	1.119*** (1.060 to 1.181)
Atrial fibrillation prevalence§	0.957* (0.925 to 0.990)	0.962 (0.898 to .030)	0.939 (0.873 to 1.011)	0.945 (0.864 to 1.034)
COPD prevalence§	1.046*** (1.028 to 1.063)	1.048** (1.013 to 1.084)	1.041 (0.999 to 1.085)	1.047 (0.996 to 1.100)
Asthma prevalence§	0.988** (0.979 to 0.996)	0.985* (0.971 to 0.999)	1.002 (0.982 to 1.023)	0.985 (0.963 to 1.007)
Heart failure prevalence§	1.005 (0.960 to 1.053)	0.971 (0.900 to 1.048)	0.987 (0.904 to 1.078)	0.929 (0.835 to 1.034)
Coronary heart disease prevalence§	1.002 (0.985 to 1.018)	1.009 (0.975 to 1.045)	0.984 (0.940 to 1.031)	1.023 (0.969 to 1.080)
Rate of attendance at type 2–4 EDs	0.928*** (0.906 to 0.950)	0.834*** (0.793 to 0.876)	0.824*** (0.775 to 0.876)	0.775*** (0.720 to 0.834)
Distance to nearest ED	0.983*** (0.979 to 0.986)	0.970*** (0.962 to 0.977)	0.974*** (0.966 to 0.982)	0.975*** (0.965 to 0.985)
Responses to GPPS†	0.931*** (0.911 to 0.952)	0.933** (0.890 to 0.977)	0.921** (0.867 to 0.978)	0.913** (0.853 to 0.977)
N	7521	7521	7521	7521

\*P<0.05, \*\*P<0.01, \*\*\*P<0.001.

†These variables are scaled so that the IRRs reflect the effects of 10 percentage point differences.

§These variables are scaled so that the IRRs reflect the effects of 1 percentage point difference. Models also include measures of primary care quality, coefficients for which are presented in table 3. They also include indicators for the 13 NHS England local offices in which a general practice is located (NHS England region: London (n=1354), Wessex (n=303), Cheshire and Merseyside (n=379), Cumbria and North East (n=450), Lancashire and Greater Manchester (n=704), Yorkshire and Humber (n=743), Central Midlands (n=550), East (n=533), North Midlands (n=489), West Midlands (n=653), South Central (n=414), South East (n=563) and South West (n=386)).

COPD, chronic obstructive pulmonary disease; ED, emergency department; GP, general practitioner; GPPS, General Practitioner Patient Survey; IRR, incidence rate ratio.

However, the magnitudes of these associations were small. We found that less than 3% of total ED attendances were associated with levels of primary care quality that were below the national average. Even when applying the strictest definition of an avoidable attendance (no treatment and no follow-up), we found that only 5.3% of these attendances were associated with levels of primary care quality below the national average. These results suggest that ED attendances may be far less responsive to changes in primary care quality than is implied by previous estimates of the prevalence of avoidable attendances. This may in turn explain the failure of many initiatives to reduce ED attendances through improvements to primary care.

While the proportion of attendances that are sensitive to primary care quality increases with the strictness of the definition, the absolute number of primary care sensitive attendances falls, from 346 000 of all attendances to 188 000 self-referred discharged attendances, 75 000 GP treatable attendances, and 44 000 attendances with no treatment and no follow-up. This suggests that while the attendances characterised by

the stricter definitions are more likely to be avoidable, some attendances not meeting these avoidability criteria are nevertheless sensitive to primary care quality. Focusing only on attendances meeting these avoidability criteria will therefore miss some attendances that are sensitive to primary care quality.

A potential explanation for this finding lies in the definitions of avoidable attendances commonly used in the literature and applied in this paper. A recent classification suggests that there are three distinct categories of avoidable ED attendances, classified based on the care that was required: clinically divertable, clinically preventable and clinically unnecessary attendances.<sup>14</sup> The definitions of an avoidable attendance that we have applied here encompass a combination of clinically divertable and clinically unnecessary attendances. However, none of the definitions capture clinically preventable attendances.

The omission of clinically preventable attendances from the definitions of avoidable attendances may therefore explain why we find that some attendances outside of the applied definitions of avoidable are

**Table 3** Associations between ED attendance volumes and primary care quality

	All ED attendances	Self-referred discharged attendances	GP treatable attendances	No treatment and no follow-up attendances
	IRR (95% CI)	IRR (95% CI)	IRR (95% CI)	IRR 95% CI)
Clinical QOF score achievement†	0.987* (0.975 to 0.998)	0.977* (0.954 to 1.000)	0.982 (0.950 to 1.014)	0.966 (0.924 to 1.010)
Proportion of patients that would recommend the practice†	0.995 (0.983 to 1.007)	0.978 (0.954 to 1.002)	0.978 (0.951 to 1.006)	0.978 (0.943 to 1.015)
Proportion of patients reporting good overall experience†	1.007 (0.989 to 1.026)	1.014 (0.983 to 1.047)	1.020 (0.984 to 1.058)	1.018 (0.973 to 1.065)
Proportion of patients that could see their preferred GP†	0.998 (0.993 to 1.004)	1.006 (0.994 to 1.017)	1.005 (0.993 to 1.018)	1.012 (0.996 to 1.029)
Proportion of patients reporting easy phone access†	0.987*** (0.981 to 0.994)	0.985* (0.970 to 0.999)	0.978** (0.963 to 0.994)	0.976* (0.955 to 0.998)
Proportion of patients that were able to get an appointment to see or speak to someone†	0.978*** (0.967 to 0.990)	0.963** (0.941 to 0.985)	0.963* (0.934 to 0.993)	0.934*** (0.898 to 0.971)
Proportion of patients that were able to get a same day appointment†	0.997 (0.991 to 1.003)	0.992 (0.981 to 1.004)	0.992 (0.977 to 1.006)	0.990 (0.972 to 1.009)
N	7521	7521	7521	7521
Pseudo-R <sup>2</sup>	0.045	0.039	0.047	0.042
Number of primary care quality sensitive attendances	346 334	187 815	74 464	44 002
Proportion of attendances sensitive to primary care quality (%)	2.48	3.70	4.10	5.26

\*P< 0.05, \*\*P< 0.01, \*\*\*P< 0.001. The reported pseudo-R<sup>2</sup> statistic is McFadden's pseudo R<sup>2</sup> calculated as one minus the ratio of the log-likelihood of the estimated model to the log-likelihood of a model containing no covariates.

†These variables are scaled so that the IRRs reflect the effects of 10 percentage point differences. Models also include practice and population characteristics, coefficients for which are presented in table 2 and indicators for the 13 NHS England local offices in which a general practice is located (NHS England region: London (n=1354), Wessex (n=303), Cheshire and Merseyside (n=379), Cumbria and North East (n=450), Lancashire and Greater Manchester (n=704), Yorkshire and Humber (n=743), Central Midlands (n=550), East (n=533), North Midlands (n=489), West Midlands (n=653), South Central (n=414), South East (n=563); South West (n=386)).

ED, emergency department; GP, general practitioner; IRR, incidence rate ratio; QOF, Quality and Outcomes Framework.

nevertheless sensitive to primary care quality. While the phenomenon of preventability through earlier intervention in primary care has received wide attention when patients require admission to the hospital in the form of ambulatory care-sensitive conditions,<sup>28</sup> this notion of preventability is largely absent in definitions of avoidable ED attendances.

In the absence of definitions able to capture all three categories of avoidable ED attendances empirically, it may be preferable to examine total attendance volumes. When evaluating the impact of interventions aimed at improving primary care quality, while a stricter definition of avoidable attendances may provide a stronger signal, examining total ED attendances will capture a greater absolute number of responsive attendances.

**Strengths and limitations**

We used national data covering all ED attendances in England over the period of a year. We applied three different definitions of avoidable attendances used in previous literature, in addition to examining total attendance volumes, and examined the sensitivity of these volumes to seven different measures of primary care quality covering access, patient experience and clinical quality.

While we attempted to include a wide variety of quality measures, information on waiting times and

rates of use were not available. We therefore had to rely on patient-reported measures of access to primary care. Our ability to detect significant associations between attendances and primary care quality was constrained by the range of variations in quality in these variables. It is also important to recognise that the variables we use to measure quality contain measurement error, particularly those from the GPPS. This too will have affected our ability to measure the associations between attendances and primary care quality. It will also mean that some of the below-average quality that we include in our impact measurement will have occurred only by chance.

We controlled for a wide range of practice and population characteristics that may have confounded the relationships between quality and the volume of avoidable attendances, including the prevalence of five chronic conditions, but were unable to control for rates of multimorbidity because these are not available from national data.<sup>29</sup>

Our calculations were made on the basis of cross-sectional associations, which are likely to overestimate the true causal magnitude of the relationships. Furthermore, raising the quality of primary care services to current average levels at a minimum across England would represent a huge achievement, far larger than any previous quality improvement programme.

Although some aspects of primary care quality have been declining over recent years,<sup>30</sup> these declines are substantially smaller than the changes in quality that we consider when raising below-average quality to national average levels. For example, the percentage of GPPS responders who report very or fairly easy telephone access in 2015/2016 was five percentage points lower than what has been previously reported for the period 2011/2012–2013/2014.<sup>31</sup> By way of comparison, the average increase in quality that we considered when raising below-average quality to national average levels on this same access measure was 15.8 percentage points. Furthermore, while the use of the contemporaneous national average is a somewhat arbitrary benchmark, we do not consider a national quality deficit to be a plausible explanation for our results.

### Relation to existing literature

Our finding that practice-level ED attendance volumes are sensitive to access to primary care services is consistent with existing research.<sup>4</sup> However, previous studies have generally only examined the relationship using one measure of ED attendances<sup>5–7 32</sup> and/or a single dimension of primary care quality.<sup>7 32</sup> Our more comprehensive examination confirms previous findings that attendance volumes are not associated with overall patient experience,<sup>7</sup> but reveals that ED attendances are sensitive to measures of primary care access and clinical quality. There remains a discrepancy between the findings on the relationship between access and use at the patient level<sup>31 33</sup> and at the practice level,<sup>5–7 32</sup> which should be a priority for future research.

The concept of avoidable ED attendances has been operationalised through consultation with clinical experts who have identified characteristics of attendances which could indicate that patients did not require ED care.<sup>34 35</sup> Our results suggest that while these criteria identify the types of attendances that are more sensitive to primary care quality, over 90% of attendances identified as potentially avoidable are not associated with below-average primary care quality regardless of which definition of avoidable is used. Some attendances classified as avoidable because the patient did not require any clinical care are in fact driven by complex social problems.<sup>14</sup> Further research is needed to explore the sensitivity of ED attendance volumes to social and welfare services, and to investigate potential solutions outside of the healthcare system.

### Policy implications

Although the volume of attendances typically thought to indicate an avoidable attendance is sensitive to primary care quality, the proportion of attendances associated with poor quality of primary care services is much smaller than the numbers suggested by previous prevalence studies and in policy documents.

The potential reduction in ED attendance volumes that could therefore be achieved through quality improvement in primary care is likely to be small.

**Twitter** Beth Parkinson @bethparky24, Rachel Meacock @RachelMeacock and Matt Sutton @MattXSutton

**Contributors** BP undertook the analysis and wrote the first draft of the manuscript. All authors contributed to the design of the analysis and the writing of the final manuscript.

**Funding** This work was funded by a Wellcome Trust PhD Studentship: 208183/Z/17/Z.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party and are not publicly available. Data are either publicly available online from NHS Digital or are available upon request from NHS Digital.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

### ORCID iDs

Beth Parkinson <http://orcid.org/0000-0003-4120-0933>  
Rachel Meacock <http://orcid.org/0000-0001-8933-5058>  
Matt Sutton <http://orcid.org/0000-0002-6635-2127>

### REFERENCES

- 1 Carter EJ, Pouch SM, Larson EL. The relationship between emergency department crowding and patient outcomes: a systematic review. *J Nurs Scholarsh* 2014;46:106–15.
- 2 Bezzina AJ, Smith PB, Cromwell D, *et al*. Primary care patients in the emergency department: who are they? A review of the definition of the 'primary care patient' in the emergency department. *Emerg Med Australas* 2005;17:472–9.
- 3 Berchet C. *Emergency care services*. Paris: Organisation for Economic Co-operation and Development, 2015. <http://www.oecd-ilibrary.org/content/workingpaper/5jrts344crns-en>
- 4 Huntley A, Lasserson D, Wye L, *et al*. Which features of primary care affect unscheduled secondary care use? A systematic review. *BMJ Open* 2014;4:e004746.
- 5 Tammes P, Morris RW, Brangan E, *et al*. Exploring the relationship between general practice characteristics, and attendance at walk-in centres, minor injuries units and EDS in England 2012/2013: a cross-sectional study. *Emerg Med J* 2016;33:702–8.
- 6 Tammes P, Morris RW, Brangan E, *et al*. Exploring the relationship between general practice characteristics and attendance at walk-in centres, minor injuries units and

- emergency departments in England 2009/10-2012/2013: a longitudinal study. *BMC Health Serv Res* 2017;17:546.
- 7 Cowling TE, Majeed A, Harris MJ. Patient experience of general practice and use of emergency hospital services in England: regression analysis of national cross-sectional time series data. *BMJ Qual Saf* 2018;27:643–54.
  - 8 Carret MLV, Fassa ACG, Domingues MR. Inappropriate use of emergency services: a systematic review of prevalence and associated factors. *Cad Saude Publica* 2009;25:7–28.
  - 9 Durand A-C, Gentile S, Devictor B, *et al.* Ed patients: how nonurgent are they? systematic review of the emergency medicine literature. *Am J Emerg Med* 2011;29:333–45.
  - 10 NHS Digital. *Hes data dictionary accident and emergency*. London: NHS Digital, 2018. [https://digital.nhs.uk/binaries/content/assets/legacy/pdf/3/l/hes\\_data\\_dictionary\\_-\\_accident\\_and\\_emergency.pdf](https://digital.nhs.uk/binaries/content/assets/legacy/pdf/3/l/hes_data_dictionary_-_accident_and_emergency.pdf)
  - 11 The Handbook to the NHS constitution for England, 2019. Available: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/770675/The\\_Handbook\\_to\\_the\\_NHS\\_Constitution\\_-\\_2019.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/770675/The_Handbook_to_the_NHS_Constitution_-_2019.pdf)
  - 12 Parkinson B, Meacock R, Sutton M. How do hospitals respond to price changes in emergency departments? *Health Econ* 2019;28:830–42.
  - 13 NHS Digital. *Hospital episode statistics*. Hospital Episode Statistics, 2018. <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics>
  - 14 Parkinson B, Meacock R, Checkland K, *et al.* Clarifying the concept of avoidable emergency department attendance. *J Health Serv Res Policy* 2020:135581962092189.
  - 15 NHS Digital. Unnecessary A and E attendances. Available: <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/innovative-uses-of-data/demand-on-healthcare/unnecessary-a-and-e-attendances> [Accessed 25 May 2018].
  - 16 McHale P, Wood S, Hughes K, *et al.* Who uses emergency departments inappropriately and when - a national cross-sectional study using a monitoring data system. *BMC Med* 2013;11:258.
  - 17 Ipsos MORI. *GP Patient Survey - National summary report*, 2016.
  - 18 Ipsos MORI. *GP Patient Survey - Technical Annex*. 2015-2016 annual report, 2016. Available: <https://gp-patient.co.uk/downloads/archive/2016/July/July%202016%20National%20Technical%20Report.pdf>
  - 19 NHS Digital. Quality and outcomes framework – prevalence, achievements and exceptions report, 2016. Available: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-rep-v2.pdf>
  - 20 Kutner M, Nachtsheim C, Neter J. *Applied linear regression models*. 4th Edn. Chicago: McGraw-Hill/Irwin, 2004.
  - 21 NHS Digital. Numbers of Patients Registered at a GP Practice - July 2015, 2015. Available: <https://digital.nhs.uk/data-and-information/publications/statistical/patients-registered-at-a-gp-practice/july-2015> [Accessed 17 Apr 2019].
  - 22 Ministry of Housing, Communities & Local Government. English indices of deprivation 2015. Available: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015>
  - 23 Ipsos MORI. Gp patient survey. July 2016. practice data (weighted), 2016. Available: <https://www.gp-patient.co.uk/surveysandreports-10-16>
  - 24 NHS Digital. QOF 2015-16: prevalence, all groups at all levels V2, 2016. Available: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-prev-all-lev-v2.xlsx> [Accessed 20 Jul 2018].
  - 25 NHS Digital. Gp and GP practice related data: GPs by GP practices (epracmem), 2018. Available: <https://digital.nhs.uk/services/organisation-data-service/data-downloads/gp-and-gp-practice-related-data> [Accessed 20 Jul 2018].
  - 26 Office for National Statistics O for N. Rural urban classification (2011) of lower layer super output areas in England and Wales, 2018. Available: <https://data.gov.uk/dataset/b1165cea-2655-4cf7-bf22-dfb3cdeb242/rural-urban-classification-2011-of-lower-layer-super-output-areas-in-england-and-wales> [Accessed 17 Oct 2019].
  - 27 ONS. Health geography - Office for National Statistics. Available: <https://www.ons.gov.uk/methodology/geography/ukgeographies/healthgeography>
  - 28 Purdy S, Griffin T, Salisbury C, *et al.* Ambulatory care sensitive conditions: terminology and disease coding need to be more specific to aid policy makers and clinicians. *Public Health* 2009;123:169–73.
  - 29 Kontopantelis E, Mamas MA, van Marwijk H, *et al.* Chronic morbidity, deprivation and primary medical care spending in England in 2015-16: a cross-sectional spatial analysis. *BMC Med* 2018;16:19.
  - 30 Cowling TE, Harris MJ, Majeed A. Evidence and rhetoric about access to UK primary care. *BMJ* 2015;350:h1513.
  - 31 Cowling TE, Harris MJ, Watt HC, *et al.* Access to general practice and visits to accident and emergency departments in England: cross-sectional analysis of a national patient survey. *Br J Gen Pract* 2014;64:e434–9.
  - 32 Cowling TE, Cecil EV, Soljak MA, *et al.* Access to primary care and visits to emergency departments in England: a cross-sectional, population-based study. *PLoS One* 2013;8:e66699.
  - 33 Zhou Y, Abel G, Warren F, *et al.* Do difficulties in accessing in-hours primary care predict higher use of out-of-hours GP services? Evidence from an English national patient survey. *Emerg Med J* 2015;32:373–8.
  - 34 Lowy A, Kohler B, Nicholl J. Attendance at accident and emergency departments: unnecessary or inappropriate? *J Public Health Med* 1994;16:134–40.
  - 35 O’Keeffe C, Mason S, Jacques R, *et al.* Characterising non-urgent users of the emergency department (ED): a retrospective analysis of routine ED data. *PLoS One* 2018;13:e0192855.

**Appendix 1 Descriptions of published definitions of avoidable attendances and the Hospital Episode Statistics Accident & Emergency data variable codes used to identify them**

Definition	Source	Description	Hospital Episode Statistics Accident & Emergency variable codes
1: Self-referred discharges	Tammes et al (1)	Self-referred discharged Emergency Department attendance.	AERFSOURCE 01 = Self-referral; AEATTENDDISP 03 = Discharged – did not require any follow-up treatment, 12 = Left department before being treated, 13 = Left department having refused treatment, or 02 = Discharged – follow-up treatment to be provided by general practitioner.
2: GP treatable	NHS Digital (2)	First attendance with some recorded treatments or investigations all of which may have been reasonably provided by a GP, followed by discharge home or to GP care.	INVEST_N: 06 = Urinalysis, 21 = Pregnancy test, 22 = Dental investigation, or 24 = None; TREAT_N: 221 = Guidance/advice only - written, 222 = Guidance/advice only - verbal, 30 = Recording vital signs, 56 = Dental treatment, 57 = Prescription/medicines prepared to take away, 99 = None, or 07 = Prescriptions; AEATTENDDISP: 02 = Discharged - follow-up treatment to be provided by general practitioner, 03 = Discharged - did not require any follow-up treatment, or 12 = Left department before being treated; AEARRIVALMODE: all codes except for 1 = non-ambulance arrivals.
3: No treatment and no follow-up	McHale et al (3)	Patients who were self-referred; were not attending a follow-up; received no investigation and either no treatment or 'guidance/advice only'; and were discharged with either no follow-up or follow-up with primary care.	AERFSOURCE: 01 = Self-referral; AEATTENDCAT: 1 = First Accident and Emergency attendance, or 3 = Follow-up Accident and Emergency attendance - unplanned; INVEST_N: 24 = None; TREAT_N: 221 = Guidance/advice only - written, 222 = Guidance/advice only - verbal, or 99 = None, AEATTENDDISP: 02 = Discharged - follow-up treatment to be provided by general practitioner, or 03 = Discharged - did not require any follow-up treatment

## Appendix 2: Data management

We applied the following process to generate annual practice-level counts of each type of avoidable emergency department attendance from the patient-level attendances data set:

Of the 14,864,083 attendances at type 1 emergency departments in England during the financial year 1<sup>st</sup> April 2015 to 31<sup>st</sup> March 2016, 64,706 (0.4%) were recorded as having no registered GP. 480,705 (3.2%) attendances were recorded as having a general practice code that was not applicable, not known, or had no practice code submitted. A further 137,274 (0.9%) of attendances had a practice code that did not belong to the master list of 7,758 practices, obtained from practice registrations data from July 2015 (4). This left n=14,181,398 attendances which we were able to link to the general practice with which the patient is registered. These attendances were then aggregated to practice-level counts of total emergency department attendances and counts of the three definitions of avoidable attendances.

Data on the practice-level counts of emergency department attendances were then merged with practice factors using GP practice identifier codes as follows: The master file of practice codes, the number of registered patients, and age-gender distribution of patients were obtained from NHS Digital (n=7,758 general practices) (4). Practices with a registered population of less than 1,000 patients were dropped (n=49). We also drop practices that were established after 1<sup>st</sup> April 2015 (n=1) and those that closed before 1<sup>st</sup> April 2016 (n=5), using practice open and close dates obtained from NHS Digital (5), since these practices were not active for the full financial year we examine. Furthermore, we drop practices that serve atypical populations (n=16), such as those that solely register students, homeless people and asylum seekers. This was done by searching practice names for the terms 'Student', 'University', 'Homeless', 'Asylum' and 'Vulnerable'. These restrictions leave n=7,687 general practices. Using this master list of general practices we then merge in attendance counts (not merged n=0), GP Patient Survey (GPPS) data (not merged n=2), and Quality and Outcomes Framework (QOF) data (not merged n=127). We exclude practices that are missing any of the covariates in the analysis (n=37), leaving a final sample of 7,521 general practices.

**Appendix 3: Description of the explanatory variables included in our analysis and the data from which these were sourced**

Variable	Description	Source
<b>Primary care quality measures</b>		
Clinical quality	Proportion of total clinical Quality and Outcomes Framework (QOF) score achieved	Quality and Outcomes Framework (6)
Recommend the practice	Proportion of patients answering 'Yes, would definitely recommend' or 'Yes, would probably recommend' to the question 'Would you recommend your GP surgery to someone who has just moved to your local area?' vs 'Not sure', 'No, would probably not recommend', 'No, would definitely not recommend', or 'Don't know'.	GP Patient Survey (7)
Good overall experience	Proportion of patients answering 'Very good or 'Fairly good' to the question 'Overall, how would you describe your experience of your GP surgery?' vs 'neither good nor poor', 'fairly poor', or 'very poor'.	GP Patient Survey (7)
See their preferred GP	Proportion of patients answering 'Always or almost always' or 'A lot of the time' to the question 'How often do you see or speak to the GP you prefer?' vs 'some of the time', 'never or almost never' or 'not tried at this GP surgery' [among patients with preferred GP].	GP Patient Survey (7)
Easy phone access	Proportion of patients answering 'Very easy' or 'Fairly easy' to the question 'Generally, how easy is it to get through to someone at your GP surgery on the phone?' vs 'Not very easy', 'Not at all easy', or 'Haven't tried'	GP Patient Survey (7)
Able to get an appointment to see or speak to someone	Proportion of patients answering 'Yes' or 'Yes, but had to call back closer to the day I wanted the appointment' to the question 'Were you able to get an appointment to see or speak to someone?' vs 'No' and 'Can't remember' [at the last time they wanted to see or speak to a GP or nurse from their GP surgery]	GP Patient Survey (7)
Able to get a same day appointment	Proportion of patients answering 'On the same day' to the question 'How long after initially contacting the surgery did you actually see or speak to them?' vs 'on the next working day', 'a few days later', 'a week or more later', or 'can't remember' [at the last time they wanted to see or speak to a GP or nurse from their GP surgery].	GP Patient Survey (7)
<b>Practice characteristics</b>		
List size	Total number of patients registered at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 0-4 years	Proportion of total list size aged 0-4 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 5-15 years	Proportion of total list size aged 5-15 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 16-44 years	Proportion of total list size aged 16-44 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 45-54 years	Proportion of total list size aged 45-54 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 55-64 years	Proportion of total list size aged 55-64 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 65-74 years	Proportion of total list size aged 65-74 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 75-84 years	Proportion of total list size aged 75-84 years at a practice as at July 2015	NHS Digital, Number of patients

Proportion of list size aged 85+ years	Proportion of total list size aged 85+ years at a practice as at July 2015	registered at a GP Practice (4) NHS Digital, Number of patients registered at a GP Practice (4)
Income deprivation	2015 Index of Multiple Deprivation (IMD) income domain score of the Lower Super Output Area (LSOA) within which the general practice is located	Ministry of Housing, Communities & Local Government. English indices of deprivation 2015 (8)
Rurality	Classifies GP practices as either 'Rural' or 'Urban' based on the LSOA of the practice	Office for National Statistics, Rural Urban Classification (2011) of Lower Layer Super Output Areas in England and Wales (9)
NHS Region	Classifies GP practices into the 13 NHS England Regions based on their Clinical Commissioning Group	ONS Clinical Commissioning Group to NHS Region (Geography) to NHS Commissioning Region (July 2015) Lookup in England (10)
Ethnicity	Proportion of respondents who answered 'UK White' to 'What is your ethnic group?'	GP patient survey (7)
Unemployment	Proportion of respondents who answered 'Unemployed' to 'Which of these best describes what you are doing at present?'	GP patient survey (7)
GP Patient Survey (GPPS) response rate	Percentage of returned surveys as a proportion of total surveys sent	GP patient survey (7)
Atrial Fibrillation prevalence	Number of patients with an 'initial event'; paroxysmal (intermittent); persistent and permanent atrial fibrillation, as a proportion of total patients registered at the practice	Quality and Outcomes Framework (11)
Chronic Obstructive Pulmonary Disease prevalence	Number of patients with a diagnosis of chronic obstructive pulmonary disease, as a proportion of total patients registered at the practice	Quality and Outcomes Framework (11)
Asthma prevalence	Number of patients with a diagnosis of asthma, as a proportion of total registered patients	Quality and Outcomes Framework (11)
Heart Failure prevalence	Number of patients with diagnosis of heart failure, as proportion of total registered patients	Quality and Outcomes Framework (11)
Coronary Heart Disease prevalence	Number of patients with a diagnosis of coronary heart disease, as a proportion of total registered patients	Quality and Outcomes Framework (11)
Distance from practice to closest emergency department	Geodetic distance in kilometres between practice postcode and closest type 1 emergency department. Postcodes and corresponding coordinates of GP practices and emergency departments were obtained and Stata's 'geonear' command was used to calculate distance to the nearest emergency department from each GP practice and the corresponding geodetic distance. When an emergency department opened or closed in 2015/16, we used the distance between each GP practice and emergency department at the start and end of the year, weighted by the proportion of the year each emergency department was open.	Practice postcodes: NHS digital, GP Practices (epracur) (5) Emergency department postcodes: department of health (12) and NHS digital (13). ONS Postcode directory (14).
Use/availability of alternative urgent care services	Rate of attendance at type 1-3 emergency departments per 1,000 registered patients	Hospital episode statistics
Total number of GPs	The average number of GPs working at the practice within 2015/16. Data contains a list of all GPs that have worked in general practice, and the periods that they have worked at each practice. We calculated the number of days that each GP worked at each practice in each year. We then summed these periods for each practice-	NHS Digital, GPs by GP practice (epracmem) (15)

year to calculate the average number of GPs working at the practice across the days of that year. We were unable to use NHS Digital's full-time equivalent GP data for the 2015/16 financial year due to the amount of missing data and changes in the methodology for calculating FTEs in 2015/16.

---

**Appendix 4: Sensitivity analysis: associations between total emergency department attendances and primary care quality**

	I Removing practices within 20km Scottish & Welsh boarder	II Including CQC inspection rating as additional practice variable	III Including unweighted 'see preferred GP' variable for practices with complete data for weighted variable	III Including weighted 'see preferred GP' variable for practices with complete data
Proportion of clinical QOF score achievement	0.988 <sup>*</sup> [0.00597]	0.982 <sup>**</sup> [0.00635]	0.988 <sup>*</sup> [0.00583]	0.988 <sup>*</sup> [0.00582]
Proportion that would recommend the practice	0.995 [0.00638]	0.992 [0.00637]	0.988 [0.00626]	0.989 [0.00625]
Proportion reporting good overall experience	1.008 [0.00966]	1.008 [0.00907]	1.013 [0.00925]	1.013 [0.00928]
Proportion that could see their preferred GP	0.998 [0.00296]	0.999 [0.00288]	0.998 [0.00285]	0.997 [0.00271]
Proportion reporting easy phone access	0.987 <sup>***</sup> [0.00338]	0.988 <sup>***</sup> [0.00311]	0.988 <sup>***</sup> [0.00331]	0.988 <sup>***</sup> [0.00330]
Proportion able to get an appointment to see or speak to someone	0.977 <sup>***</sup> [0.00604]	0.982 <sup>**</sup> [0.00560]	0.980 <sup>**</sup> [0.00604]	0.981 <sup>**</sup> [0.00607]
Proportion able to get a same day appointment	0.997 [0.00293]	0.996 [0.00289]	0.995 [0.00287]	0.995 [0.00288]
Practice received CQC rating good or outstanding	- -	0.991 [0.00922]	- -	- -
<i>N</i>	7189	7209	7124	7124
pseudo <i>R</i> <sup>2</sup>	0.045	0.047	0.046	0.046
Primary care quality sensitive attendances	338,798	335,058	341,947	350,183
Proportion of attendances which are sensitive to PC	2.52%	2.46%	2.47%	2.53%

Notes: Standard errors clustered at Clinical Commissioning Group level in brackets. Models also include practice and population characteristics and indicators for the 13 NHS England local offices in which a general practice is located (NHS England Region: London (N=1,354); Wessex (N=303); Cheshire and Merseyside (N=379); Cumbria and North East (N=450); Lancashire and Greater Manchester (N=704); Yorkshire and Humber (N=743); Central Midlands (N=550); East (n=533); North Midlands (N=489); West Midlands (N=653); South Central (N= 414); South East (N=563); South West (N=386)).  $p < 0.05$ ,  $** p < 0.01$ ,  $*** p < 0.001$ .

## Sources

1. Tammes P, Morris RW, Brangan E, Checkland K, England H, Huntley A, et al. Exploring the relationship between general practice characteristics, and attendance at walk-in centres, minor injuries units and EDs in England 2012/2013: a cross-sectional study. *Emerg Med J EMJ*. 2016 Oct;33(10):702–8.
2. NHS Digital. Unnecessary A and E attendances [Internet]. NHS Digital. [cited 2018 May 25]. Available from: <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/innovative-uses-of-data/demand-on-healthcare/unnecessary-a-and-e-attendances>
3. McHale P, Wood S, Hughes K, Bellis MA, Demnitz U, Wyke S. Who uses emergency departments inappropriately and when - a national cross-sectional study using a monitoring data system. *BMC Med*. 2013 Dec 13;11:258.
4. NHS Digital. Numbers of Patients Registered at a GP Practice - July 2015 [Internet]. 2015 [cited 2019 Apr 17]. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/patients-registered-at-a-gp-practice/july-2015>
5. NHS Digital. GP and GP practice related data: GP Practices (epraccur) [Internet]. NHS Digital; 2018 May [cited 2018 Jul 20]. Available from: <https://digital.nhs.uk/services/organisation-data-service/data-downloads/gp-and-gp-practice-related-data/www.england.nhs.uk/statistics/statistical-work-areas/gp-patient-survey/>
6. NHS Digital. QOF 2015-16: Achievements at practice level, all domains [Internet]. 2016. Available from: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-prac-dom-ach.xlsx>
7. Ipsos MORI. GP Patient Survey. July 2016. Practice data (weighted). [Internet]. 2016. Available from: <https://www.gp-patient.co.uk/surveysandreports-10-16>
8. Ministry of Housing, Communities & Local Government. English indices of deprivation 2015 [Internet]. GOV.UK. 2015 [cited 2018 Jul 30]. Available from: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015>
9. Office for National Statistics O for N. Rural Urban Classification (2011) of Lower Layer Super Output Areas in England and Wales [Internet]. 2018 [cited 2019 Oct 17]. Available from: <https://data.gov.uk/dataset/b1165cea-2655-4cf7-bf22-dfbd3cdeb242/rural-urban-classification-2011-of-lower-layer-super-output-areas-in-england-and-wales>
10. Clinical Commissioning Group to NHS Region (Geography) to NHS Commissioning Region (July 2015) Lookup in England [Internet]. 2015 [cited 2019 Oct 17]. Available from: <https://geoportal.statistics.gov.uk/items?tags=Local%20Commissioning%20Group>
11. NHS Digital. QOF 2015-16: Prevalence, all groups at all levels v2 [Internet]. NHS Digital; 2016 Oct [cited 2018 Jul 20]. Available from: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-prev-all-lev-v2.xlsx>
12. Department of Health. Address of Major A&E departments in England and Wales. (Obtained under the Freedom of Information Act (2000) from The Department of Health; 26th October 2012; reference DE00000729700) [Internet]. 2012 [cited 2019 Nov 3]. Available from: [https://www.whatdotheyknow.com/request/address\\_of\\_major\\_ae\\_departments](https://www.whatdotheyknow.com/request/address_of_major_ae_departments)
13. NHS Digital. Addresses of major (type 1) A&E departments in England. (Obtained under the Freedom of Information Act (2000) from NHS Digital; 12 April 2017; Reference NIC-98262-D1L6S) [Internet]. 2017 [cited 2019 Nov 3]. Available from: [https://www.whatdotheyknow.com/request/address\\_of\\_major\\_ae\\_departments](https://www.whatdotheyknow.com/request/address_of_major_ae_departments)

14. Office for National Statistics. ONS Postcode Directory (May 2016) [Internet]. 2017 [cited 2019 Feb 1]. Available from: <https://data.gov.uk/dataset/bf86a637-7e4e-43df-ac96-2cb4670f799b/ons-postcode-directory-may-2016>
15. NHS Digital. GP and GP practice related data: GPs by GP Practices (epracmem) [Internet]. 2018 May [cited 2018 Jul 20]. Available from: <https://digital.nhs.uk/services/organisation-data-service/data-downloads/gp-and-gp-practice-related-data>

**Appendix 1 Descriptions of published definitions of avoidable attendances and the Hospital Episode Statistics Accident & Emergency data variable codes used to identify them**

Definition	Source	Description	Hospital Episode Statistics Accident & Emergency variable codes
1: Self-referred discharges	Tammes et al (1)	Self-referred discharged Emergency Department attendance.	AERFSOURCE 01 = Self-referral; AEATTENDDISP 03 = Discharged – did not require any follow-up treatment, 12 = Left department before being treated, 13 = Left department having refused treatment, or 02 = Discharged – follow-up treatment to be provided by general practitioner.
2: GP treatable	NHS Digital (2)	First attendance with some recorded treatments or investigations all of which may have been reasonably provided by a GP, followed by discharge home or to GP care.	INVEST_N: 06 = Urinalysis=, 21 = Pregnancy test, 22 = Dental investigation, or 24 = None; TREAT_N: 221 = Guidance/advice only - written, 222 = Guidance/advice only - verbal, 30 = Recording vital signs, 56 = Dental treatment, 57 = Prescription/medicines prepared to take away, 99 = None, or 07 = Prescriptions; AEATTENDDISP: 02 = Discharged - follow-up treatment to be provided by general practitioner, 03 = Discharged - did not require any follow-up treatment, or 12 = Left department before being treated; AEARRIVALMODE: all codes except for 1 = non-ambulance arrivals.
3: No treatment and no follow-up	McHale et al (3)	Patients who were self-referred; were not attending a follow-up; received no investigation and either no treatment or 'guidance/advice only'; and were discharged with either no follow-up or follow-up with primary care.	AERFSOURCE: 01 = Self-referral; AEATTENDCAT: 1 = First Accident and Emergency attendance, or 3 = Follow-up Accident and Emergency attendance - unplanned; INVEST_N: 24 = None; TREAT_N: 221 = Guidance/advice only - written, 222 = Guidance/advice only - verbal, or 99 = None, AEATTENDDISP: 02 = Discharged - follow-up treatment to be provided by general practitioner, or 03 = Discharged - did not require any follow-up treatment

## Appendix 2: Data management

We applied the following process to generate annual practice-level counts of each type of avoidable emergency department attendance from the patient-level attendances data set:

Of the 14,864,083 attendances at type 1 emergency departments in England during the financial year 1<sup>st</sup> April 2015 to 31<sup>st</sup> March 2016, 64,706 (0.4%) were recorded as having no registered GP. 480,705 (3.2%) attendances were recorded as having a general practice code that was not applicable, not known, or had no practice code submitted. A further 137,274 (0.9%) of attendances had a practice code that did not belong to the master list of 7,758 practices, obtained from practice registrations data from July 2015 (4). This left  $n=14,181,398$  attendances which we were able to link to the general practice with which the patient is registered. These attendances were then aggregated to practice-level counts of total emergency department attendances and counts of the three definitions of avoidable attendances.

Data on the practice-level counts of emergency department attendances were then merged with practice factors using GP practice identifier codes as follows: The master file of practice codes, the number of registered patients, and age-gender distribution of patients were obtained from NHS Digital ( $n=7,758$  general practices) (4). Practices with a registered population of less than 1,000 patients were dropped ( $n=49$ ). We also drop practices that were established after 1<sup>st</sup> April 2015 ( $n=1$ ) and those that closed before 1<sup>st</sup> April 2016 ( $n=5$ ), using practice open and close dates obtained from NHS Digital (5), since these practices were not active for the full financial year we examine. Furthermore, we drop practices that serve atypical populations ( $n=16$ ), such as those that solely register students, homeless people and asylum seekers. This was done by searching practice names for the terms 'Student', 'University', 'Homeless', 'Asylum' and 'Vulnerable'. These restrictions leave  $n=7,687$  general practices. Using this master list of general practices we then merge in attendance counts (not merged  $n=0$ ), GP Patient Survey (GPPS) data (not merged  $n=2$ ), and Quality and Outcomes Framework (QOF) data (not merged  $n=127$ ). We exclude practices that are missing any of the covariates in the analysis ( $n=37$ ), leaving a final sample of 7,521 general practices.

**Appendix 3: Description of the explanatory variables included in our analysis and the data from which these were sourced**

Variable	Description	Source
<b>Primary care quality measures</b>		
Clinical quality	Proportion of total clinical Quality and Outcomes Framework (QOF) score achieved	Quality and Outcomes Framework (6)
Recommend the practice	Proportion of patients answering 'Yes, would definitely recommend' or 'Yes, would probably recommend' to the question 'Would you recommend your GP surgery to someone who has just moved to your local area?' vs 'Not sure', 'No, would probably not recommend', 'No, would definitely not recommend', or 'Don't know'.	GP Patient Survey (7)
Good overall experience	Proportion of patients answering 'Very good or 'Fairly good' to the question 'Overall, how would you describe your experience of your GP surgery?' vs 'neither good nor poor', 'fairly poor', or 'very poor'.	GP Patient Survey (7)
See their preferred GP	Proportion of patients answering 'Always or almost always' or 'A lot of the time' to the question 'How often do you see or speak to the GP you prefer?' vs 'some of the time', 'never or almost never' or 'not tried at this GP surgery' [among patients with preferred GP].	GP Patient Survey (7)
Easy phone access	Proportion of patients answering 'Very easy' or 'Fairly easy' to the question 'Generally, how easy is it to get through to someone at your GP surgery on the phone?' vs 'Not very easy', 'Not at all easy', or 'Haven't tried'	GP Patient Survey (7)
Able to get an appointment to see or speak to someone	Proportion of patients answering 'Yes' or 'Yes, but had to call back closer to the day I wanted the appointment' to the question 'Were you able to get an appointment to see or speak to someone?' vs 'No' and 'Can't remember' [at the last time they wanted to see or speak to a GP or nurse from their GP surgery]	GP Patient Survey (7)
Able to get a same day appointment	Proportion of patients answering 'On the same day' to the question 'How long after initially contacting the surgery did you actually see or speak to them?' vs 'on the next working day', 'a few days later', 'a week or more later', or 'can't remember' [at the last time they wanted to see or speak to a GP or nurse from their GP surgery].	GP Patient Survey (7)
<b>Practice characteristics</b>		
List size	Total number of patients registered at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 0-4 years	Proportion of total list size aged 0-4 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 5-15 years	Proportion of total list size aged 5-15 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 16-44 years	Proportion of total list size aged 16-44 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 45-54 years	Proportion of total list size aged 45-54 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 55-64 years	Proportion of total list size aged 55-64 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 65-74 years	Proportion of total list size aged 65-74 years at a practice as at July 2015	NHS Digital, Number of patients registered at a GP Practice (4)
Proportion of list size aged 75-84 years	Proportion of total list size aged 75-84 years at a practice as at July 2015	NHS Digital, Number of patients

Proportion of list size aged 85+ years	Proportion of total list size aged 85+ years at a practice as at July 2015	registered at a GP Practice (4) NHS Digital, Number of patients registered at a GP Practice (4)
Income deprivation	2015 Index of Multiple Deprivation (IMD) income domain score of the Lower Super Output Area (LSOA) within which the general practice is located	Ministry of Housing, Communities & Local Government. English indices of deprivation 2015 (8)
Rurality	Classifies GP practices as either 'Rural' or 'Urban' based on the LSOA of the practice	Office for National Statistics, Rural Urban Classification (2011) of Lower Layer Super Output Areas in England and Wales (9)
NHS Region	Classifies GP practices into the 13 NHS England Regions based on their Clinical Commissioning Group	ONS Clinical Commissioning Group to NHS Region (Geography) to NHS Commissioning Region (July 2015) Lookup in England (10)
Ethnicity	Proportion of respondents who answered 'UK White' to 'What is your ethnic group?'	GP patient survey (7)
Unemployment	Proportion of respondents who answered 'Unemployed' to 'Which of these best describes what you are doing at present?'	GP patient survey (7)
GP Patient Survey (GPPS) response rate	Percentage of returned surveys as a proportion of total surveys sent	GP patient survey (7)
Atrial Fibrillation prevalence	Number of patients with an 'initial event'; paroxysmal (intermittent); persistent and permanent atrial fibrillation, as a proportion of total patients registered at the practice	Quality and Outcomes Framework (11)
Chronic Obstructive Pulmonary Disease prevalence	Number of patients with a diagnosis of chronic obstructive pulmonary disease, as a proportion of total patients registered at the practice	Quality and Outcomes Framework (11)
Asthma prevalence	Number of patients with a diagnosis of asthma, as a proportion of total registered patients	Quality and Outcomes Framework (11)
Heart Failure prevalence	Number of patients with diagnosis of heart failure, as proportion of total registered patients	Quality and Outcomes Framework (11)
Coronary Heart Disease prevalence	Number of patients with a diagnosis of coronary heart disease, as a proportion of total registered patients	Quality and Outcomes Framework (11)
Distance from practice to closest emergency department	Geodetic distance in kilometres between practice postcode and closest type 1 emergency department. Postcodes and corresponding coordinates of GP practices and emergency departments were obtained and Stata's 'geonear' command was used to calculate distance to the nearest emergency department from each GP practice and the corresponding geodetic distance. When an emergency department opened or closed in 2015/16, we used the distance between each GP practice and emergency department at the start and end of the year, weighted by the proportion of the year each emergency department was open.	Practice postcodes: NHS digital, GP Practices (epracur) (5) Emergency department postcodes: department of health (12) and NHS digital (13). ONS Postcode directory (14).
Use/availability of alternative urgent care services	Rate of attendance at type 1-3 emergency departments per 1,000 registered patients	Hospital episode statistics
Total number of GPs	The average number of GPs working at the practice within 2015/16. Data contains a list of all GPs that have worked in general practice, and the periods that they have worked at each practice. We calculated the number of days that each GP worked at each practice in each year. We then summed these periods for each practice-	NHS Digital, GPs by GP practice (epracmem) (15)

year to calculate the average number of GPs working at the practice across the days of that year. We were unable to use NHS Digital's full-time equivalent GP data for the 2015/16 financial year due to the amount of missing data and changes in the methodology for calculating FTEs in 2015/16.

---

**Appendix 4: Sensitivity analysis: associations between total emergency department attendances and primary care quality**

	I Removing practices within 20km Scottish & Welsh boarder	II Including CQC inspection rating as additional practice variable	III Including unweighted 'see preferred GP' variable for practices with complete data for weighted variable	III Including weighted 'see preferred GP' variable for practices with complete data
Proportion of clinical QOF score achievement	0.988 <sup>*</sup> [0.00597]	0.982 <sup>**</sup> [0.00635]	0.988 <sup>*</sup> [0.00583]	0.988 <sup>*</sup> [0.00582]
Proportion that would recommend the practice	0.995 [0.00638]	0.992 [0.00637]	0.988 [0.00626]	0.989 [0.00625]
Proportion reporting good overall experience	1.008 [0.00966]	1.008 [0.00907]	1.013 [0.00925]	1.013 [0.00928]
Proportion that could see their preferred GP	0.998 [0.00296]	0.999 [0.00288]	0.998 [0.00285]	0.997 [0.00271]
Proportion reporting easy phone access	0.987 <sup>***</sup> [0.00338]	0.988 <sup>***</sup> [0.00311]	0.988 <sup>***</sup> [0.00331]	0.988 <sup>***</sup> [0.00330]
Proportion able to get an appointment to see or speak to someone	0.977 <sup>***</sup> [0.00604]	0.982 <sup>**</sup> [0.00560]	0.980 <sup>**</sup> [0.00604]	0.981 <sup>**</sup> [0.00607]
Proportion able to get a same day appointment	0.997 [0.00293]	0.996 [0.00289]	0.995 [0.00287]	0.995 [0.00288]
Practice received CQC rating good or outstanding	- -	0.991 [0.00922]	- -	- -
<i>N</i>	7189	7209	7124	7124
pseudo <i>R</i> <sup>2</sup>	0.045	0.047	0.046	0.046
Primary care quality sensitive attendances	338,798	335,058	341,947	350,183
Proportion of attendances which are sensitive to PC	2.52%	2.46%	2.47%	2.53%

Notes: Standard errors clustered at Clinical Commissioning Group level in brackets. Models also include practice and population characteristics and indicators for the 13 NHS England local offices in which a general practice is located (NHS England Region: London (N=1,354); Wessex (N=303); Cheshire and Merseyside (N=379); Cumbria and North East (N=450); Lancashire and Greater Manchester (N=704); Yorkshire and Humber (N=743); Central Midlands (N=550); East (n=533); North Midlands (N=489); West Midlands (N=653); South Central (N= 414); South East (N=563); South West (N=386)).  $p < 0.05$ ,  $** p < 0.01$ ,  $*** p < 0.001$ .

## Sources

1. Tammes P, Morris RW, Brangan E, Checkland K, England H, Huntley A, et al. Exploring the relationship between general practice characteristics, and attendance at walk-in centres, minor injuries units and EDs in England 2012/2013: a cross-sectional study. *Emerg Med J EMJ*. 2016 Oct;33(10):702–8.
2. NHS Digital. Unnecessary A and E attendances [Internet]. NHS Digital. [cited 2018 May 25]. Available from: <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/innovative-uses-of-data/demand-on-healthcare/unnecessary-a-and-e-attendances>
3. McHale P, Wood S, Hughes K, Bellis MA, Demnitz U, Wyke S. Who uses emergency departments inappropriately and when - a national cross-sectional study using a monitoring data system. *BMC Med*. 2013 Dec 13;11:258.
4. NHS Digital. Numbers of Patients Registered at a GP Practice - July 2015 [Internet]. 2015 [cited 2019 Apr 17]. Available from: <https://digital.nhs.uk/data-and-information/publications/statistical/patients-registered-at-a-gp-practice/july-2015>
5. NHS Digital. GP and GP practice related data: GP Practices (epraccur) [Internet]. NHS Digital; 2018 May [cited 2018 Jul 20]. Available from: <https://digital.nhs.uk/services/organisation-data-service/data-downloads/gp-and-gp-practice-related-data/www.england.nhs.uk/statistics/statistical-work-areas/gp-patient-survey/>
6. NHS Digital. QOF 2015-16: Achievements at practice level, all domains [Internet]. 2016. Available from: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-prac-dom-ach.xlsx>
7. Ipsos MORI. GP Patient Survey. July 2016. Practice data (weighted). [Internet]. 2016. Available from: <https://www.gp-patient.co.uk/surveysandreports-10-16>
8. Ministry of Housing, Communities & Local Government. English indices of deprivation 2015 [Internet]. GOV.UK. 2015 [cited 2018 Jul 30]. Available from: <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2015>
9. Office for National Statistics O for N. Rural Urban Classification (2011) of Lower Layer Super Output Areas in England and Wales [Internet]. 2018 [cited 2019 Oct 17]. Available from: <https://data.gov.uk/dataset/b1165cea-2655-4cf7-bf22-dfbd3cdeb242/rural-urban-classification-2011-of-lower-layer-super-output-areas-in-england-and-wales>
10. Clinical Commissioning Group to NHS Region (Geography) to NHS Commissioning Region (July 2015) Lookup in England [Internet]. 2015 [cited 2019 Oct 17]. Available from: <https://geoportal.statistics.gov.uk/items?tags=Local%20Commissioning%20Group>
11. NHS Digital. QOF 2015-16: Prevalence, all groups at all levels v2 [Internet]. NHS Digital; 2016 Oct [cited 2018 Jul 20]. Available from: <https://files.digital.nhs.uk/publicationimport/pub22xxx/pub22266/qof-1516-prev-all-lev-v2.xlsx>
12. Department of Health. Address of Major A&E departments in England and Wales. (Obtained under the Freedom of Information Act (2000) from The Department of Health; 26th October 2012; reference DE00000729700) [Internet]. 2012 [cited 2019 Nov 3]. Available from: [https://www.whatdotheyknow.com/request/address\\_of\\_major\\_ae\\_departments](https://www.whatdotheyknow.com/request/address_of_major_ae_departments)
13. NHS Digital. Addresses of major (type 1) A&E departments in England. (Obtained under the Freedom of Information Act (2000) from NHS Digital; 12 April 2017; Reference NIC-98262-D1L6S) [Internet]. 2017 [cited 2019 Nov 3]. Available from: [https://www.whatdotheyknow.com/request/address\\_of\\_major\\_ae\\_departments](https://www.whatdotheyknow.com/request/address_of_major_ae_departments)

14. Office for National Statistics. ONS Postcode Directory (May 2016) [Internet]. 2017 [cited 2019 Feb 1]. Available from: <https://data.gov.uk/dataset/bf86a637-7e4e-43df-ac96-2cb4670f799b/ons-postcode-directory-may-2016>
15. NHS Digital. GP and GP practice related data: GPs by GP Practices (epracmem) [Internet]. 2018 May [cited 2018 Jul 20]. Available from: <https://digital.nhs.uk/services/organisation-data-service/data-downloads/gp-and-gp-practice-related-data>