



OPEN ACCESS

Exploring the actionability of healthcare performance indicators for quality of care: a qualitative analysis of the literature, expert opinion and user experience

Erica Barbazza , Niek S Klazinga , Dionne S Kringos

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2020-011247>).

Department of Public and Occupational Health, Academic Medical Centre, Amsterdam University Medical Centres, University of Amsterdam, Amsterdam, The Netherlands

Correspondence to

Erica Barbazza, Department of Public and Occupational Health, Academic Medical Centre, Amsterdam University Medical Centres, University of Amsterdam, 1105 AZ Amsterdam, The Netherlands; e.s.barbazza@amsterdamumc.nl

Received 29 March 2020
Revised 19 April 2021
Accepted 23 April 2021
Published Online First
7 May 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

To cite: Barbazza E, Klazinga NS, Kringos DS. *BMJ Qual Saf* 2021;**30**:1010–1020.

ABSTRACT

Background This study explores the meaning of actionable healthcare performance indicators for quality of care-related decisions. To do so, we analyse the constructs of *fitness for purpose* and *fitness for use* across healthcare systems and in practice based on the literature, expert opinion and user experience.

Methods A multiphase qualitative study was undertaken. Phases included a literature review, a first round of one-on-one interviews with a panel of academics and thought leaders in the field (n=16), and a second round of interviews with real-world users of performance indicators (n=16). Thematic analysis was conducted between phases in order to triangulate findings in a stepwise process.

Results Common uses of healthcare performance indicators were differentiated within micro-meso-macro contexts of healthcare systems. Each purpose of use signals different decision-making tasks, and in effect information needs. An indicator's fitness for use can be appraised by three clusters of considerations: methodological, contextual and managerial. Methodological considerations gauge an indicator's perceived importance, engagement potential, interpretability, standardisation, feasibility of remedial actions, alignment to care models and sensitivity to change. Information infrastructure, system governance, workforce capacity and learning culture were found as enabling contextual considerations. Managerial considerations influencing an indicator's use in practice were found to span the selection of indicators, data collection, analysis, display of results and delivery of information to decision-makers.

Conclusion The actionability of a healthcare performance indicator should be appraised by its alignment with the intended purpose of use beyond aggregate healthcare system levels, in combination with the extent to which methodological, contextual and managerial fitness for use considerations are met. Striking a better balance between the importance weighted to an indicator's statistical merits and emphasis put to its fitness for purpose and use is needed for indicators that are ultimately actionable for quality of care-related decision-making.

INTRODUCTION

Healthcare performance measurement, and its use as performance intelligence,

plays an important role in guiding the decisions of healthcare system actors with respect to quality of care.¹ Since the early 2000s, the importance of performance measurement in healthcare,² its institutionalisation as standard practice within³ and across healthcare systems,^{4–6} and more recently its professionalisation⁷ has received widespread prioritisation. This attention has increased scientific rigour around criteria for selecting indicators (eg, reliability, validity),^{8–9} development of indicator sets (eg, parsimony, epidemiological relevance),¹⁰ and methods, tools and approaches to guide these processes.^{11–13}

Importantly, adherence to agreed-upon criteria for a statistically sound indicator does not guarantee that it is useful for decision-making. The information needs of decision-makers across healthcare systems, including policy-makers, managers, clinicians and patients, are varied. The type of indicator, data sources, level of precision, timeliness and relevant comparisons are among the key differences.^{14–15} For example, working to improve antibiotic prescribing, a primary care clinician may assess new and re-prescribing of antibiotics in their practice quarterly; an insurer, the adherence of practices to prescribing guidelines for issuing payment incentives annually; and a policy-maker, the total volume of antibiotics prescribed per 100 000 population by region, nationally and in comparison with other countries by policy cycle.

In effect, the ability for an indicator to meet the information needs of decision-makers goes beyond their statistical quality and is rather a measure of their

actionability. To be actionable, it is generally agreed an indicator should be both *fit for purpose*—serving an intended decision-making function—and *fit for use*—getting the right information into the right hands at the right time.^{16–18} While there is agreement on the importance of actionability,^{18–20} and increasing attention put to its two main constructs of fitness for purpose and use, it still remains an elusive concept to define, assess and operationalise. In the absence of a common understanding of the meaning of actionability, the tendency to select indicators on the merit of their *potential* to be actionable perpetuates.^{18 21–23} And while there are implicit criteria that appear to influence the actual use of indicators, such as data availability and ease of interpretation,^{1 15 24–26} how these relate across different healthcare systems remains underexplored.^{1 14 15}

With the advancement of information systems and data analytics, there has been impressive growth in the speed, volume and range of data available for performance measurement.^{27 28} COVID-19 and the ensuing surge in performance data reported is evidence of this.^{29 30} It also serves to illustrate that an abundance of information does not translate to informed decisions. Our attention is increasingly called to this fact and the work still needed to advance methods for measuring quality of care^{31–33} and patient safety³⁴ in order to obtain additional value from our data-rich systems.^{35–38}

In this study, we set out with the aim to gain further insights into the meaning of actionable healthcare performance indicators for quality of care-related decision-making across healthcare systems. To do so, we explore the notions of fitness for purpose and fitness for use derived through the existing literature, expert opinion and experiences of data users in varied developed country contexts. We pose two questions. The first aims to differentiate an indicator's purpose of use by micro-meso-macro decision-making levels, investigating what are the uses of healthcare performance indicators across healthcare systems. The second aims to consolidate the determinants of an indicator's fitness for use, exploring what are the key considerations influencing an indicator's use.

METHODS

Design

We applied qualitative methods³⁹ in a multiphase approach, comprising a review to examine actionability according to the published literature⁴⁰ and multiple perspective semistructured interviews^{39 41 42} to gain insights from two groups (panels) representing the scientific community and data users. We employed one-on-one interviews following our literature review rather than a questionnaire or focus groups for richer exchanges and the possibility to elicit the individual opinions of each participant.⁴³ Our stepwise approach to analysis allowed for the triangulation of findings across phases and to aggregate individual-level



Figure 1 Decision-making contexts across healthcare systems.

results for panel-wide themes.⁴² The study adheres to the Consolidated Criteria for Reporting Qualitative Research.³⁹

Indicators refer to a quantifiable variable measured to provide simplified information about a larger area of interest,⁴⁴ typically measured over time.^{9 45} In the scope of this study, we focus on *healthcare performance indicators*: indicators for quality of care-driven decision-making to improve performance on one or more of the six dimensions of quality: safe, effective, patient-centred, timely, efficient and equitable care.^{8 46} As an exploratory study, we prioritised the generalisability of findings and were inclusive of varied types of healthcare (eg, primary, acute, specialist, long-term care), settings (eg, primary care, hospitals), health system types and countries, although limited to developed country contexts.

To explore our first research question, we took as a basis the characterisation of decision-making in healthcare systems by three contexts: patient care (micro-level), organisational (meso-level) and policy (macro-level), as illustrated in figure 1.^{47 48} Indicators are used to inform decisions in each context, be it quality improvement, services management, population health planning or other strategic and tactical choices.

Data collection and analysis

Phase 1: literature review and content analysis

We reviewed the existing literature with the following aims: to examine the current scientific understanding of actionable healthcare performance indicators; to generate an initial core list of indicator purposes of use and fitness for use considerations; and to identify leading experts in the field. Our search was conducted using PubMed at the outset of the study in early 2019.

The search was limited to the past 10 years and articles published in English using the following key terms in varied combinations: health care performance indicator, actionability, quality of care, measurement and use. We also reviewed reporting of relevant international organisations and networks, namely the WHO and its regional offices, the Organisation for Economic Co-operation and Development (OECD), and the European Commission Expert Group on Health Systems Performance Assessment. Reference lists of articles and reports identified were reviewed in a snowballing approach.

The results of the initial literature search were synthesised and used to inform a provisional approach and visualisation of the uses of healthcare performance indicators by micro-meso-macro context. Recurrent fitness for use considerations were also distilled and clustered. These findings were prepared as an expert panel brief for use as a background document in the second phase (online supplemental appendix 1).

Phase 2: interviews with expert panel and thematic analysis

The first panel aimed to engage prominent academics and thought leaders in the field of healthcare performance measurement and quality of care (hereafter, expert panel). Experts were identified based on the authorship of literature reviewed and with consideration to the following criteria for the panel's composition: a balance of expertise in areas related to quality of care, performance measurement, governance, data and information systems or management; senior academic or technical roles related to their area of expertise; and affiliation to varied healthcare systems and geographical contexts. A target of 15 experts were pursued for manageability and presumed saturation.⁴⁹

One pilot interview was conducted to ensure relevance and clarity. Piloting resulted in the addition of illustrative examples of data users and fitness for use considerations. Panellists were invited to participate via email and received a panel brief in advance. The brief provided relevant study details together with the findings of phase 1. All interviews were conducted by the primary researcher (EB, female) with experience in semistructured interviews and subject matter expertise. Interviews took place between August and September 2019 both in person and at distance based on the proximity and preference of panellists. Interviews lasted between 45 and 60 min. Records of the interviews were prepared as detailed summaries rather than verbatim transcripts in the approach described by Halcomb and Davidson.⁵⁰ The research adheres to the Dutch ethics guidelines stated in the 'Medical Research Act with People (Wet medisch-wetenschappelijk onderzoek met mensen (WMO)) (Dutch), in BWBR0009408, W.a.S. Ministry of Health, Editor. 1998: Hague, Netherlands',⁵¹ for which verbal consent was deemed adequate by the authors as no human data were retained. To ensure informed voluntary participation,

participants provided written agreement to participate in the study during the recruitment stage and restated verbally their consent at the start of all interviews.

The interview records of this first panel were stored in an Excel-based tool for thematically analysing themes (EB). The analysis incorporated a deductive and inductive approach: topics explored in the interviews (online supplemental appendix 1) were used to guide the deductive thematic analysis⁵² and new themes that emerged were identified using an inductive approach.⁵³ The data were also interpreted by redrawing conceptual diagrams. Two others (DSK, NSK) with complementary expertise in quality of care, performance measurement, health governance and management reviewed the findings to ensure consistency and reach agreement on the theme extraction.

Phase 3: interviews with user panel and thematic analysis

The findings from the expert panel were used to refine the mapping of uses of healthcare performance by micro-meso-macro level and fitness for use themes. The revisions were summarised in a new brief prepared for a second panel of one-on-one interviews (online supplemental appendix 2). This panel aimed to engage real-world data users for their first-hand experiences using healthcare performance indicators for quality of care-related decision-making (hereafter, user panel).

A target of 15 data users actively contributing to the further development of this field were pursued as panellists. The selection drew on existing membership lists of international networks, working groups and projects related to healthcare performance indicators, measurement and quality of care, such as the OECD Health Care Quality Indicators Project⁵⁴ and initiatives of the European Commission (eg, HealthPros⁵⁵). The panel composition aimed to capture a range of perspectives, with representation of differing health system types, country affiliations and uses of healthcare performance indicators. Interviews were conducted in the same manner as the first panel and were completed between November 2019 and January 2020.

Interview records were consolidated in the existing Excel-based tool for further thematic analysis. The topics and themes explored were used to refine and/or confirm the classification resulting from the expert panel on uses of healthcare performance indicators and fitness for use considerations. Observing the convergence of themes, with this phase data collection and analysis were considered complete.

RESULTS

Literature review and panel results

Based on the literature synthesis, 19 experts were identified and invited to participate in the first panel. Of these, 16 experts agreed to participate. Non-participants were either unreachable (n=1), unavailable (n=1) or referred to an alternative contact (n=1). Together, expert panellists had published more than

Table 1 Characteristics of panellists

Expert panel	n (%)	User panel	n (%)
Total	16 (–)		16 (–)
Affiliation*		Uses	
Academia	10 (63)	Macro	7 (44)
International organisation	3 (19)	Meso	4 (25)
Think tank	3 (19)	Micro	3 (19)
Expertise		Organisation type	
Measurement	5 (31)	Government	5 (31)
Quality of care	3 (19)	Health services	4 (25)
Governance	3 (19)	Standards	3 (19)
Information systems	3 (19)	Research	2 (13)
Management	2 (13)	Improvement	2 (13)
Region			
Europe	9 (56)		9 (56)
North America	5 (31)		7 (44)
Oceania	2 (13)		–
Sex			
Male	11 (69)		9 (56)
Female	5 (31)		7 (44)

*Primary affiliations.

50 articles or reports on the use, selection or improvement of healthcare performance indicators at the time of study. This literature (online supplemental appendix 3) was reviewed in phase 1 together with other relevant

works.^{22 34 44 48 56–64} Expert panellists were predominantly affiliated to academia and in senior or executive roles spanning eight countries (Australia, Canada, Denmark, Germany, Italy, the Netherlands, UK and USA). A range and balance of areas of expertise that included performance measurement, quality of care, governance, information systems and management were achieved.

The user panel comprised participants spanning the micro-level, meso-level and macro-level of healthcare systems. Participants included representatives of national health authorities, health standards and accreditation agencies, insurers, professional associations, as well as clinicians and patient advocates. In total, 31 participants were contacted, of which 16 agreed to participate (online supplemental appendix 3). Non-participants reported the same reasons as the first panel, with the majority (n=6) referring to an alternative contact and the remainder being either unreachable (n=5) or unavailable (n=4). User panellists spanned seven countries (Belgium, Canada, Germany, Ireland, the Netherlands, UK and USA). **Table 1** summarises the key characteristics across panellists.

From the literature reviewed, 11 clusters of uses of healthcare performance indicators and fitness for use considerations related to the methodological quality of an indicator were identified (**figure 2**). In the second




Figure 2 Summary of key findings across study phases. Note: boxes denote key themes emerging by study phase. Broken lines denote a change in level. Solid lines denote agreement between phases with possible adjustments to phrasing. Darker grey shading denotes the introduction of new elements. Ordering within cells is not indicative of importance.

phase, there was agreement across experts on the relevance and importance to distinguish purposes of use of healthcare performance indicators beyond aggregate micro-level, meso-level and macro-level. The panel shared strong views to avoid a hierarchy within levels, finding this introduced a rigidity that may not translate across contexts. Rather, the framing of uses identified as common or frequent was found more transferable.

The experts introduced further consistency, refinements and additional purposes of use and fitness for use considerations. Specifically, the uses of indicators for functions such as regulation or strategy development were differentiated from mechanisms to achieve these functions, such as international comparisons or public reporting. Refinements to the distribution of uses across levels were introduced for consistency, for example, recategorising *improvements of organisations and networks* to the meso-level. Additions included emphasis on the use of indicators by patients as a decision-maker for informed choice and the cross-cutting function of research. The clustering of fitness for use considerations was disaggregated, with emphasis on the importance of considering an indicator's use in a specific setting (where it is used) and as a process (how it is used).

In the third phase, user panellists agreed with the categorisation of uses by micro-level, meso-level and macro-level. *Accountability* was viewed as an aim rather than specific use and *external assessments* were viewed rather as a mechanism. There were detailed discussions on fitness for use considerations, with agreement to classify considerations that underscored the importance of the setting in which an indicator is used for its contextualisation. The case was made to view *practical* considerations as *managerial* aspects related to the process of using indicators.

Purposes of use of healthcare performance indicators

Through our stepwise approach to data collection and analysis, common uses of healthcare performance indicators were differentiated beyond the aggregate decision-making contexts of patient care (micro-level), organisations (meso-level) and policy (macro-level). In [table 2](#), we list the uses for healthcare performance indicators identified, each serving different managerial decision-making functions, users and information needs. The purposes of use are not exhaustive and may take varied forms by healthcare system. Specifically, expert and user panellists noted variation in the degree of patient choice, role of insurers or mandate of professional bodies.

The detailed differentiation of uses of healthcare performance indicators signals important, yet often overlooked, distinctions in information needs *within* system levels. To illustrate these differences, we take the macro-level as an example. While uses of healthcare performance indicators in this context share an overall aim of informing policy decisions,

distinctions between uses include system performance monitoring—signalling to system stakeholders, often including the public, the performance of the system as a whole, answering ‘How is my health care system doing?’; or strategy development—signalling to ministries, departments of health or similar with the aim of identifying priority areas, monitor trends and ultimately answering ‘Have I chosen the right areas to prioritize?’; or system quality assurance—informing decisions of health service executives, quality inspectors or quality observatories for an overview of care processes and signalling of incidents, answering ‘Is care being delivered as intended?’

Fitness for use of healthcare performance indicators

Three main clusters of considerations influencing the second construct of actionability—fitness for use—were found. These include methodological, contextual and managerial considerations ([table 3](#)).

Methodological considerations

Methodological considerations pertain to the indicator itself, although beyond its statistical quality. Seven recurrent considerations were identified. First, an indicator should measure what matters. User panellists emphasised the importance that the target audience *cares* about the results, explaining an indicator that ‘moves’ people makes everyone uncomfortable that the right thing is not already being done. Second, the extent to which an indicator resonates with a range of stakeholders was emphasised as a key gauge of its ability to facilitate a ‘what can *we* do’ approach, rather than limiting action to an individual user.⁶⁵ Third, an indicator's inherent ease of interpretation was described by panellists and in the literature^{18 66 67} to strongly influence an end user's confidence in their interpretation of its meaning. Fourth, the extent to which an indicator is clearly defined was described as a key contributor to trust in what it signals, as well as the likelihood of wide uptake. Fifth, an indicator should be able to be broken down into its constituent parts to make change points clear,⁸ with panellists finding a remote or disconnected indicator from a user's performance difficult to act on.^{59 63} Sixth, an indicator should measure a phenomenon as true to lived experience as possible.^{27 68} The tendency to focus on specific (siloe) areas of care was described to reduce performance to overly narrow aspects of care and, as one user panellist described, misses the ‘system-ness’ of quality. Lastly, the ability of an indicator to be sufficiently sensitive to change based on its intended use was described by both panels as intuitive, yet often a challenge for an indicator to meet.

Contextual considerations

Contextual considerations refer to critical factors pertaining to the setting in which an indicator is used. Four main clusters emerged. One, the information

Table 2 Differentiating uses of healthcare performance indicators across healthcare systems

Context	Purpose of use	Illustrative uses	Illustrative users	Illustrative information need
Macro	System performance monitoring.	Signalling the performance of the system as a whole; comparing performance internationally; publicly reporting system performance.	Public; ministry of health; regional (provincial, state) authorities; health service executive (authority).	How is my healthcare system doing? How does it compare with others?
	Strategy development.	Setting health policy priorities; identifying emerging health priority areas; and monitoring trends in current priority areas.	Government and ministries; regional (provincial, state) authorities; accountable care organisations; health maintenance organisations.	Have I chosen the right areas to prioritise? What is the impact of strategies that are in place?
	System quality assurance.	Measuring care processes; reporting of incidents and never events.	Quality inspectorate; national quality observatory; health and safety executive.	Is care being delivered as intended? Where do problems in the delivery of care lie?
Meso	Regulation (professional, facility, pharmaceuticals).	Informing accreditation, certification and/or licensing processes.	Medical councils, chambers, college of physicians; medicines and healthcare products regulatory agency.	Does the performance of organisations, facilities, medicines, etc, meet established standards?
	Professional development.	Reporting internally and benchmarking within profession or specialty.	Societies of medical professionals; professional associations; training institutions.	How do healthcare professionals of a specific specialty perform?
	Quality-based financing.	Issuing performance-based payment (pay-for-performance); value-based contracting.	Healthcare insurers; healthcare providers.	Are existing guidelines or standards being adhered to? Does this merit the issuing of incentives?
	Organisation/network performance improvement.	Improving performance of hospitals, networks and care groups; assessing local needs and geographical differences.	Hospital management; integrated care; networks/groups; local collaboratives of care.	Are affiliated practices/facilities performing optimally?
Micro	Practice or team performance improvement.	Convening audit and feedback, plan-do-study-act, and/or collaborative, team-based improvement cycles; comparing across practices.	Primary care practices; specialist departments or units; pathways of care.	How is my team performing? How can we improve our performance? How do I perform relative to my team members?
	Individual performance improvement.	Identifying trends in the management of patients; tailoring services to target groups.	Individual physicians; nurse/practitioners; other healthcare professionals.	How am I managing my practice panel? How can I improve my performance?
	Informed choice.	Selecting a healthcare provider; participating in care decision-making; self-managing care needs.	Patients; family members and carers; public.	What treatment options or providers are best for me?
Cross-cutting	Research.	Exploring the use of indicators across contexts.	Academia and academic networks; think tanks, research groups; topic-specific associations.	Secondary user-directed.

infrastructure was met with consensus across panellists as a key predictor of use, determining the ability to collect, store and extract information. Relevant considerations repeatedly raised included the interoperability of information systems (ie, linkages, output format) and overall data quality (ie, consistency in field, codes, maintenance). Second, characteristics of governance were emphasised, with panellists citing the importance of political will and vision, regulatory arrangements for data exchanges, as well as cross-sector partnerships and aligned financing structures. Third, workforce capacity considerations were underscored, specifically the data literacy skills of actors across the healthcare system and the availability of protected time for the healthcare workforce to use data. Lastly, pertaining to culture and professional norms, be it in clinical practice, healthcare organisations, professional networks or government agencies, the importance of a learning orientation and shared sense of responsibility was

emphasised as a predictor of the importance placed to measurement and ultimately the use of an indicator.

Managerial considerations

The importance of embedding indicators into performance management systems is well established.^{60 63 69–72} Based on the literature and insights from the panels, we conceptualised an indicator's use cycle (figure 3). This cycle was used to consolidate considerations brought forward around embedding indicators into management systems to safeguard an indicator's use in practice. The considerations reflect key decisions to be managed across the cycle and include selecting an indicator with consideration to define clear parameters of its intended use,^{18 38 73} gain clarity around its construction,⁶⁰ assess data needs and define measurement considerations; accessing data to ensure data are available, of quality or can feasibly be collected⁴⁸; applying methods of analysis for the relevant calculation of

Table 3 Overview of methodological, contextual and managerial fitness for use considerations

Clusters	Considerations	Guiding questions for considering an indicator's use
<i>Methodological</i>		
	Measures what matters. Wide engagement. Easily interpreted. Clear standardisation. Alignment of accountability. Measurement matches delivery. Sensitive to meaningful change.	Does anybody care? What can we do? Does the indicator signal a clear direction? Is the indicator clearly defined and replicable? Are entry points for taking action feasible? Is the indicator a reflection of the system? Is the indicator sufficiently sensitive to change?
<i>Contextual</i>		
Information infrastructure	Interoperability. Data quality.	Can needed data be accessed? Is the data of quality?
Governance	Political will and vision. Regulation for data protection. Cross-sector partnerships. Aligned financing structures.	Is there high-level commitment and direction for use? Does existing legislation facilitate use? Are cross-sector partnerships in place? Do financing structures encourage the intended use?
Workforce capacity	Data and quality expertise. Time dedicated to improvement.	Are the competencies to interpret and use data in place? Is time allocated to encourage use?
Culture	Learning orientation. Shared responsibility for health.	Is an environment for learning cultivated? Do users feel accountable for improvement?
<i>Managerial</i>		
Selecting healthcare performance indicators	Clear purpose of use. Target end user is known. Conceptual framework. Indicator quality. Source, type and availability of data. Standards for appraisal. Degree of public disclosure. Accompanying indicators. Previous use.	What is the purpose of use? (eg, strategy development) Is the target audience known? (eg, clinicians, public) Is the dimension of quality pursued clear? Is the indicator scientifically sound? What data are needed and are they available? (eg, administrative, clinical, survey data, wearables) How will improvements in performance be assessed? Is the indicator for internal or external (public) use? Are there relevant accompanied indicators? Has the indicator been used previously?
Accessing data	Representativeness of data. Data linkages. Data collection tools. Unity of language/coding.	Are the data complete? Can relevant data sources be linked? How will data be collected? (eg, paper-based, automated electronically, manual electronic entry) Is there consistency in coding across data to be used?
Applying methods of analysis	Type of analysis. Aggregation of indicators. Reference group. Breakdowns/cohorts. Calculation of values. Time interval. Application of risk adjustments. Managing missing data. Contextualising data.	How will the data be analysed? (eg, benchmarking, time trend, case mix correction) How can composites/indices be used to simplify data? Who is the reference group? How will the data be disaggregated? (eg, age, sex, ethnicity, geographically) How will values be calculated? (eg, mean, median, SD, top 10% mean) Should a time trend be reported and at what interval? How will risk adjustments be applied? (eg, variable specification, source, weighting scheme) How will missed data points be handled? What other data are needed to give the indicator meaning?
Displaying findings	Chart options. Simplification techniques. Customisation of display. Narrated interpretation. Format of reporting.	How will the data be visualised? (eg, chart, map, table) What techniques to simplify the meaning can be applied? (eg, colour, size variation, icons) How can users customise the data? (eg, change of display, change of information) How can the quality and the meaning of data be narrated? How will it be reported? (eg, print, mobile, web-based)
Reaching decision-makers	Frequency of reporting. Dissemination channels. Guidance on use.	What is the relevant reporting cycle (eg, real time, quarterly, annually, biennially) How will users be reached? (eg, mail, email, champions) How can users be supported to make use of findings?




Figure 3 Use cycle for managing healthcare performance indicators.

values that correspond to the intended purpose⁶³; displaying findings⁷⁴, including decisions around how data are visualised⁷⁴ and the degree of *story-telling* to describe and interpret results to support understanding of what is meant and any caveats^{48 75 76}; and actually reaching decision-makers, with decisions needed as to the frequency of dissemination, channel used for delivering information and guidance (if any) to facilitate the use of information provided.⁶³

DISCUSSION

Principal findings

Healthcare performance indicators share a common aim to provide simplified, readily understood information to facilitate decision-making.^{9 44 45} An indicator's ability to do so in practice extends beyond its statistical quality and rather is characterised by its actionability.^{16–18 67} In this study, we explored actionability through the two constructs of fitness for purpose and fitness for use and observe the following main findings into their further operationalisation.

First, the different uses of an indicator within micro-meso-macro and research contexts stress the importance of clarity and precision on the intended use of an indicator. The relevance of precision regarding an indicator's use has been stressed in the literature^{15 18–20 23} and previously explored from the perspective of different end users.¹ Our findings further differentiate uses of indicators across healthcare systems. While not pursuing a universal, exhaustive listing of purposes of use—recognising varied healthcare system types and contextual considerations that deem this irrelevant—our findings signal the imperative of clarity regarding an indicator's intended use and user to gauge its potential usefulness. The taxonomy of uses of healthcare performance indicators can be an input to further operationalise the construct of fitness for purpose.

Second, we find an indicator's fitness for use is captured by three types of considerations. These relate to an indicator's technical qualities, its intended context of use and its handling across what can be characterised as a use cycle. It means, to gauge an

indicator's fitness for use, a range of considerations should be assessed that span, for example, 'Does the indicator signal a clear direction?' to 'Can needed data be accessed?' and 'What is the relevant reporting cycle?' The listed considerations (table 3) based on the literature and views of panellists are a testament to the wide range of variables weighing on an indicator's use that require thoughtful handling.

Third, an indicator's fitness for purpose and fitness for use should be taken together to appraise actionability. For example, a policy-maker may identify a target to be measured in the scope of a strategy, yet for this specific purpose fitness for use considerations may not be met due to information system constraints or other contextual limitations. In another instance, an indicator may meet fitness for use considerations yet lack a clear and specific purpose and, in effect, misses a target audience. In both cases, the actionability of the indicator is compromised.

Lastly, as the expertise and lived experience of panellists served to highlight, the actionability of an indicator is not a guarantee of impact. Literature on the misuse, manipulation of data and unintended consequences of performance measurement depicts this.^{45 73} This distinction between action and impact underscores that while actionable healthcare performance indicators may be a precursor to better decision-making, the impact of an indicator weighs on considerations of its own.

Applications and further research

This study has sought to consolidate the relevant literature and engage informants from differing contexts, areas of expertise and first-hand experiences for diverse insights. Future research should test the findings empirically, investigating purposes of use and fitness for use considerations by specific country contexts, governance structures, services delivery systems or areas of specialisation.

The findings of this study have a range of potential applications. In the context of the COVID-19 pandemic, actionable healthcare performance indicators have proven of paramount importance,^{29 77} and surges in publicly reported data illustrate the increased demand for information.^{78 79} The extent to which this information informs decision-making is a reflection of the alignment between an indicator's intended purpose of use and related fitness for use considerations. The findings could also inform the selection of indicators for measurement frameworks and indicator sets that cascade healthcare system levels by priority areas (eg, tackling the misuse of antibiotic prescribing, strengthening integrated care), where different decision-making functions need to work in combination.

Limitations

These findings may not be generalisable beyond the context of developed countries. The effect of system

conditions, such as level of decentralisation, public-private mix and development status, has not been captured nor investigated given the targeted sample of informants, and as suggested should be explored empirically. The initial literature review was limited to English-language materials, which may also impact the generalisability of findings. Engaging expert panellists beyond English-speaking countries sought to minimise this. Some nuances may have become lost in choosing to summarise rather than transcribe interviews, although the advantages of our approach were found better suited for the study aims and design. In exploring performance indicators in the scope of healthcare, the study has not captured the broader use of indicators for public health despite its importance. Distributing panellists between panels was to the discretion of the study team for the purposes of the two-panel design, although many participants held positions or memberships suitable to both. The value of engaging panellists from different perspectives and stages took precedent. The prominence of panellists meant some were known to the authors. In order to avoid bias, a consistent interviewer was selected with the least previous engagement with panellists.

CONCLUSION

Clarifying the meaning of actionable healthcare performance indicators is a prerequisite to its further operationalisation. This study has explored the body of literature on the actionability of healthcare performance indicators for quality of care-related decision-making together with expert opinion and data user experiences in an effort to unpack the constructs of fitness for purpose and fitness for use. The study aimed to capture these constructs from a system perspective. The findings signal the importance of clarity and precision on an indicator's purpose of use and context for the handling of methodological, contextual and managerial considerations weighing on its use in practice. Striking a better balance between the importance weighted to an indicator's statistical merits and emphasis put to an indicator's fitness for purpose and use is needed for indicators that are actionable for quality of care-related decision-making.

Acknowledgements We thank the following individuals for their contribution at varied stages of this work: JohnMarc Alban; Thomas Boeckz; Jeffrey Braithwaite; Emma Cartwright; Louise Clement; Cheryl Damberg; Greg Dempsey; Gail Dobel; Tejal Gandhi; Oliver Groene; Torsten Hecke; Peter Hibbert; John Lavis; Doreen MacNeil; Jan Mainz; Stephanie Medlock; Rob Nelissen; Sabina Nuti; Jillian Oderkirk; Irene Papanicolas; Lotte Ramerman; Ari Robiseck; Alexandru Rotar; Eric Schneider; Peter Smith; Jorien Soethout; Juan Tello; Rob Tollenaar; Michael van den Berg; Jeremy Veillard; Robert Verheij; and Naira Yeitsvan.

Contributors EB, NSK and DSK conceived and designed the study. EB collected the data. EB, NSK and DSK analysed the data and drafted the manuscript. All authors revised the manuscript.

Funding This study was funded by H2020 Marie Skłodowska-Curie Actions (765141).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request to the corresponding author.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Erica Barbazza <http://orcid.org/0000-0001-7621-1638>

Niek S Klazinga <http://orcid.org/0000-0002-3937-8014>

Dionne S Kringos <http://orcid.org/0000-0003-2711-4713>

REFERENCES

- 1 Smith P. Part 1: Principles of performance measurement. In: *Performance measurement for health system improvement: experiences, challenges and prospects*. Copenhagen: WHO Regional Office for Europe, 2008.
- 2 WHO Regional Office for Europe. *The Tallinn charter: health systems for health and wealth*. Copenhagen: WHO Regional Office for Europe, 2008.
- 3 Fekri O, Macarayan ER, Klazinga N. Health system performance assessment in the WHO European Region: which domains and indicators have been used by Member States for its measurement? In: *Health evidence network synthesis report 55*. Copenhagen: WHO Regional Office for Europe, 2018.
- 4 World Health Organization. *The world health report 2000: health systems: improving performance*. Geneva: World Health Organization, 2000.
- 5 Kelley J, Hurst J. *Health care quality indicators project conceptual framework paper, in OECD health working papers No. 23*. Paris: OECD, 2006.
- 6 Perić N, Hofmarcher-Holzhaecker MM, Simon J. Health system performance assessment landscape at the EU level: a structured synthesis of actors and actions. *Arch Public Health* 2017;75:5.
- 7 Kringos DS, Groene O, Johnsen SP. Training the first generation of health care performance intelligence professionals in Europe and Canada. *Acad Med* 2019;94:747–8.
- 8 Mainz J. Developing evidence-based clinical indicators: a state of the art methods primer. *Int J Qual Health Care* 2003;15:5i–11.
- 9 Mainz J. Defining and classifying clinical indicators for quality improvement. *Int J Qual Health Care* 2003;15:523–30.

- 10 Boerma T, AbouZahr C, Evans D, *et al.* Monitoring intervention coverage in the context of universal health coverage. *PLoS Med* 2014;11:e1001728.
- 11 McGlynn EA. The outcomes utility index: will outcomes data tell us what we want to know? *Int J Qual Health Care* 1998;10:485–90.
- 12 de Koning J, Burgers J, Klazinga N. *Appraisal of indicators through research and evaluation (AIRE)*. Amsterdam: University of Amsterdam, 2008.
- 13 Fitch K. *The RAND/UCLA appropriateness method user's manual*. Santa Monica, CA: RAND, 2001.
- 14 Quentin W. Measuring healthcare quality. In: Busse R, ed. *Improving healthcare quality in Europe: characteristics, effectiveness and implementation of different strategies*. WHO and OECD: Copenhagen, 2019.
- 15 Damberg CL, Sorbero ME, Lovejoy SL, *et al.* An evaluation of the use of performance measures in health care. *Rand Health Q* 2012;1:3.
- 16 OECD. *Health in the 21st century: putting data to work for stronger health systems*. Paris: OECD, 2019.
- 17 McDowell R. *Signs to look for: criteria for developing and selecting fit for purpose indicators*. Wellington, New Zealand: PricewaterhouseCoopers, 2017.
- 18 Smith P, Mossialos E, Papanicolas I. *Performance measurement for health system improvement: experiences, challenges and prospects*. Copenhagen: WHO Regional Office for Europe, 2008.
- 19 Carinci F, Van Gool K, Mainz J, *et al.* Towards actionable international comparisons of health system performance: expert revision of the OECD framework and quality indicators. *Int J Qual Health Care* 2015;27:137–46.
- 20 European Commission Brussels. *Expert group on health system performance assessment, so what? strategies across Europe to assess quality of care*, 2016.
- 21 Smith P. Principles of performance measurement. In: *Performance measurement for health system improvement*. Cambridge: Cambridge University Press, 2009.
- 22 Klazinga N, Stronks K, Delnoij D, *et al.* Indicators without a cause. reflections on the development and use of indicators in health care from a public health perspective. *Int J Qual Health Care* 2001;13:433–8.
- 23 Hilarion P, Suñol R, Groene O, *et al.* Making performance indicators work: the experience of using consensus indicators for external assessment of health and social services at regional level in Spain. *Health Policy* 2009;90:94–103.
- 24 Smith P. *Health system performance assessment*. Brussels: European Commission, 2014.
- 25 Secanell M, Groene O, Arah OA, *et al.* Deepening our understanding of quality improvement in Europe (DUQuE): overview of a study of hospital quality management in seven countries. *Int J Qual Health Care* 2014;26:5–15.
- 26 iHD. Transform introduction enriching knowledge and enhancing care through data, 2016. Available: <https://www.i-hd.eu/index.cfm/resources/ec-projects-results/transform/>
- 27 Nuti S. Let's play the patients music: A new generation of performance measurement systems in healthcare. *Manag Decis* 2018;56:2252–72.
- 28 Verheij R. *Reuse of routine care data for policy and science: how things can be improved [Dutch]*. Utrecht: Nivel, 2019.
- 29 Kringos D, Carinci F, Barbazza E, *et al.* Managing COVID-19 within and across health systems: why we need performance intelligence to coordinate a global response. *Health Res Policy Syst* 2020;18:80.
- 30 The Lancet Infectious Diseases. The COVID-19 infodemic. *Lancet Infect Dis* 2020;20:875.
- 31 National Academies of Sciences, E.a.M. *Crossing the global quality chasm: improving health care worldwide*. Washington, DC: National Academies of Sciences, Engineering and Medicine, 2018.
- 32 World Health Organization, World Bank Group, and Organisation for Economic Co-operation and Development. *Delivering quality health services: a global imperative for universal health coverage*. Geneva: World Health Organization, 2018.
- 33 Kruk ME, Gage AD, Arsenault C, *et al.* High-quality health systems in the sustainable development goals era: time for a revolution. *Lancet Glob Health* 2018;6:e1196–252.
- 34 Salzburg Global Seminar. *The Salzburg statement on moving measurement into action: global principles for measuring patient safety*. Institute for Healthcare Improvement and Salzburg Global Seminar, 2019.
- 35 Panch T, Szolovits P, Atun R. Artificial intelligence, machine learning and health systems. *J Glob Health* 2018;8:020303.
- 36 Panch T, Pearson-Stuttard J, Greaves F, *et al.* Artificial intelligence: opportunities and risks for public health. *Lancet Digit Health* 2019;1:e13–14.
- 37 Verheij RA, Curcin V, Delaney BC, *et al.* Possible sources of bias in primary care electronic health record data use and reuse. *J Med Internet Res* 2018;20:e185.
- 38 Braithwaite J, Hibbert P, Blakely B, *et al.* Health system frameworks and performance indicators in eight countries: a comparative international analysis. *SAGE Open Med* 2017;5:2050312116686516:205031211668651.
- 39 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 40 Baumeister RF, Leary MR. Writing narrative literature reviews. *Rev Gen Psychol* 1997;1:311–20.
- 41 Kvale S, interviews D. *The SAGE qualitative research kit*. Los Angeles, Calif: Sage, 2009.
- 42 Vogl S, Schmidt E-M, Zartler U. Triangulating perspectives: ontology and epistemology in the analysis of qualitative multiple perspective interviews. *Int J Soc Res Methodol* 2019;22:611–24.
- 43 Keeney S, Hasson F, McKeena H. *The Delphi technique in nursing and health research*. nited Kingdom Sinead Keeney, Felicity Hasson and Hugh McKenna, 2011.
- 44 Hammond AL. *Environmental indicators : a systematic approach to measuring and reporting on environmental policy performance in the context of sustainable development*, 1995.
- 45 Astleithner F, Hamedinger A, Holman N, *et al.* Institutions and indicators – the discourse about indicators in the context of sustainability. *J Hous Built Environ* 2004;19:7–24.
- 46 Institute of Medicine. *Crossing the quality chasm: a new health system for the 21st century*. Washington, DC: National Academy Press, 2001.
- 47 Plochg T, Klazinga NS. Community-based integrated care: myth or must? *Int J Qual Health Care* 2002;14:91–101.
- 48 Raleigh VS, Foot C. *Getting the measure of quality: opportunities and challenges*. London: The King's Fund, 2010.
- 49 Fink A, Kosecoff J, Chassin M, *et al.* Consensus methods: characteristics and guidelines for use. *Am J Public Health* 1984;74:979–83.

- 50 Halcomb EJ, Davidson PM. Is verbatim transcription of interview data always necessary? *Appl Nurs Res* 2006;19:38–42.
- 51 Ministry of Health. *Medical research act with people (Wet medisch-wetenschappelijk onderzoek met mensen (WMO)) [Dutch]*, in BWBR0009408. Hague, Netherlands: Ministry of Health, 1998.
- 52 King N. Using templates in the thematic analysis of text. In: *Essential guide to qualitative methods in organizational research*. Sage, 2004.
- 53 Virginia B, Victoria C. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3:77–101.
- 54 OECD. OECD health care quality indicators project - background. Available: <http://www.oecd.org/els/health-systems/oecdhealthcarequalityindicatorsproject-background.htm>
- 55 HealthPros. International training network for healthcare performance intelligence professionals, 2020. Available: <https://www.healthpros-h2020.eu/>
- 56 Freeman T. Using performance indicators to improve health care quality in the public sector: a review of the literature. *Health Serv Manage Res* 2002;15:126–37.
- 57 Grover V, Chiang RHL, Liang T-P, *et al.* Creating strategic business value from big data analytics: a research framework. *J Manag Inf Syst* 2018;35:388–423.
- 58 Esquivel A, Meric-Bernstam F, Bernstam EV. Accuracy and self correction of information received from an Internet breast cancer list: content analysis. *BMJ* 2006;332:939–42.
- 59 Compton-Phillips A, Lee TH. *The “give a darn” method for outcomes measurement*, in *NEJM catalyst*. NEJM Catalyst, 2018.
- 60 van den Berg MJ, Kringos DS, Marks LK, *et al.* The Dutch health care performance report: seven years of health care performance assessment in the Netherlands. *Health Res Policy Syst* 2014;12:1.
- 61 Berwick DM, James B, Coye MJ. Connections between quality measurement and improvement. *Med Care* 2003;41:1–30.
- 62 Arnaboldi M, Lapsley I, Steccolini I. Performance management in the public sector: the ultimate challenge. *Financial Account Manag* 2015;31:1–22.
- 63 Brehaut JC, Colquhoun HL, Eva KW, *et al.* Practice feedback interventions: 15 suggestions for optimizing effectiveness. *Ann Intern Med* 2016;164:435–41.
- 64 Saver BG, Martin SA, Adler RN, *et al.* Care that matters: quality measurement and health care. *PLoS Med* 2015;12:e1001902.
- 65 Nuti S, Vola F, Bonini A, *et al.* Making governance work in the health care sector: evidence from a 'natural experiment' in Italy. *Health Econ Policy Law* 2016;11:17–38.
- 66 Hibbard JH, Peters E. Supporting informed consumer health care decisions: data presentation approaches that facilitate the use of information in choice. *Annu Rev Public Health* 2003;24:413–33.
- 67 Marshall MN, Romano PS, Davies HTO. How do we maximize the impact of the public reporting of quality of care? *Int J Qual Health Care* 2004;16:i57–63.
- 68 Nuti S, De Rosis S, Bonciani M, *et al.* Rethinking healthcare performance evaluation systems towards the People-Centredness approach: their pathways, their experience, their evaluation. *Healthc Pap* 2017;17:56–64.
- 69 CIHI. How is an indicator developed at CIHI? 2019. Available: <https://www.cihi.ca/en/how-is-an-indicator-developed-at-cihi> [Accessed 3 Feb 2020].
- 70 Michel P, Fraticelli L, Parneix P, *et al.* Assessing the performance of indicators during their life cycle: the mixed quid method. *Int J Qual Health Care* 2020;32:12–19.
- 71 Hibbert PD, Wiles LK, Cameron ID, *et al.* CareTrack Aged: the appropriateness of care delivered to Australians living in residential aged care facilities: a study protocol. *BMJ Open* 2019;9:e030988.
- 72 Pronovost PJ, Miller MR, Dorman T, *et al.* Developing and implementing measures of quality of care in the intensive care unit. *Curr Opin Crit Care* 2001;7:297–303.
- 73 Mannion R, Braithwaite J. Unintended consequences of performance measurement in healthcare: 20 salutary lessons from the English National health service. *Intern Med J* 2012;42:569–74.
- 74 Ballard A. Promoting performance information use through data visualization: evidence from an experiment. *Public Perform Manag Rev* 2020;43:109–28.
- 75 Robicsek A. *Six modest proposals for health care measurement in NEJM catalyst*. Waltham, MA: NEJM Catalyst, 2019.
- 76 Canadian Institute for Health Information, Better Information for Improved Health. *A vision for health system use of data in Canada*. Ottawa: CIHI, 2013.
- 77 WHO Regional Office for Europe. *Strengthening and adjusting public health measures throughout the COVID-19 transition phases*. Copenhagen: WHO Regional Office for Europe, 2020.
- 78 Kennedy H. *Simply data visualizations have become key to communicating about the COVID-19 pandemic, but we know little about their impact in LSE impact Blog*. UK: London School of Economics and Political Sciences London, 2020.
- 79 Fisher D, Teo YY, Nabarro D. Assessing national performance in response to COVID-19. *Lancet* 2020;396:653–5.

Online supplementary files

Appendix 1: Questions to expert panel

Summary of users and users of health care performance data

Through our exploratory literature review we have identified a range of data uses that transcend the levels of a health care system. This list is not exhaustive, nor does it intend to be. Importantly, it is also limited to the use of data for the purposes of clinical care rather than a broader population health perspective. Nonetheless, it does ambition to capture a minimum or core set of purposes for use and actors that resonate across different health care systems. Similarly, the listing of actors is not exhaustive and is to the exclusion of other actors that may carry out multiple functions. The classification has attempted to illustrate a unique user by each managerial function and actors that are explicitly assigned a mandate to improve quality of care and made directly accountable to a health care system. As such, actors such as professional and patient associations, academia and other research groups are excluded.

This listing of distinct uses has been distinguished at the micro- (clinical practice), meso- (institutions/organizations) and macro- (policy) level as shown by the nested sub-levels of the figure below and listed again in the table that follows for an editable version.


At each sub-level, a unique user can be identified – characterized as the primary actor with the responsible authority for a specific function (purpose of use). These purposes of use form different layers to decision-making; each embedded within one another. The interactions between layers signal the ways in which the users are related and co-dependent on the use and transfer of information, or as other authors have referred to as a ‘data food chain’, as some may reuse data collected and used by others.

- Is the list of purposes for use complete and accurate? If not, what is missing? Are there purposes for use that appear overlapping?
- Is the list of users complete and accurate? If not, what is missing?
- Is the cascading of users and uses accurate? If not, how can it be improved?

Table A1. Uses and users of health care performance indicators: listed

Level	Managerial function (uses)	User
Macro <i>Policy</i>	International comparisons	International organizations
	Governmental strategy	Government
	Governmental monitoring	Ministry of health
Meso <i>Institutional/ organizations</i>	Supervision/safety	Health care inspectorate
	Public reporting/accountability	Health care institute
	Purchasing/payment	Health care insurer
	Accreditation	Accreditation agency
Micro <i>Clinical practice</i>	Networks of professional performance	Management of institute/service provider network
	Operational quality management	Unit (department, ward) of service provider
	Quality improvement	Professional and professional team/pathways
	Care decision-making	Patient-professional

<i>Other</i>
--------------	-----	-----

Figure A1. Uses and users of health care performance indicators across the health system: visualized

Methodological considerations

For each purpose for use, a range of methodological consequences (data requirements) can be anticipated. For example, the type of measure (e.g. more process measures at the micro-level vs more outcome measures at the macro-level) or confidence intervals (e.g. narrow for precision decisions of health practitioners vs wide for policy-makers). We have consolidated a range of methodologically relevant considerations (features of data) that has been highlighted in the literature as potentially relevant depending on the intended purpose of use. Examples to illustrate the variation for each of these considerations are listed in the table.

To further illustrate these considerations in practice, we have mapped existing indicators for measuring the performance of breast cancer services to different users and uses. The mapping serves to demonstrate the variability in information needs by user/use.

- Are the listed considerations meaningful? What is missed? What would you phrase differently? How can this list be expanded upon to serve its intended purpose of supporting users based on their intended purpose of use?
- How do you expect these considerations to vary based on specific uses and users from the table above?

Table A2. Fitness for use considerations and examples

Relevant considerations	Examples of variations
Type of measure	process, outcome, patient-reported
Sources	administrative, medical records or clinical, registries, patient
Reference group	unit or department, types of units, region, province, country etc.
Reference period	previous 1–6 quarters; 1–12 months; 1–6 years; previous year, etc.
Comparators	past performance, targets, reference group
Compounds (composites)	selection of indicators, weights
Calculation of values	standard deviation, means, median, other percentiles (75th), rank, top 10% mean (ABC method), individual peer scores, ranges
Statistical chart options	funnel plot, Shewhart chart, scan statistics, moving average, sets method
Types of analysis	benchmarking, time trend, international comparison
Confidence intervals	narrow for precision decisions (practitioners) vs wide for policy-makers
Risk adjustments	variable specification (demographic, clinical factors, socio-economic, health related, patient preferences), source, weighting scheme
Others...	...

General reflections


- Do you agree with the construction and approach to exploring fitness for use of health care performance data as pursued in this study?
- Are you aware of any existing studies that should be consulted in the scope of this work?

Appendix 2: Questions to user panel

Purposes of use of health care performance indicators

Uses of health care performance indicators can be differentiated at the micro- (clinical practice), meso- (institutions/organizations) and macro- (policy) level of health systems. Methodologically relevant distinctions by the purposes for using health care performance indicators are shown in the nested sub-levels of the figure below. For example, at the micro-level, uses of health care performance indicators include for informed choice (e.g. by patients), individual professional performance improvement and practice or team performance improvement. Cross-cutting these uses are factors that ultimately weigh on the extent to which a specific purpose of use can be met. These feed-in factors can be clustered around the specific country context, governance and inputs (e.g. research, data infrastructure).

1. Based on the figure (A2), how would you describe your most predominate purpose for using health care performance indicators?
2. For this purpose, what in your opinion constitutes a good indicator? That is, what information do you need? Can you give examples?
3. Can you describe how you make use of health care performance indicators? Who is your target end-user? What specific strategies or mechanisms are relied on (e.g. public reporting, licensing or accreditation, financial incentives, etc.)?


Figure A2. Uses and users of health care performance indicators across the health system: visualize

Considerations for the use of health care performance indicators

We recognize that depending on the purpose of use, each user has different information needs. The type of indicator (e.g. process, structure outcome), sources of data (e.g. administrative, clinical, survey data), timeliness (e.g. weekly, quarterly, annually) or type of analysis (e.g. benchmarking, time trend, international comparison), are among some of these differences.

Given this variability, there are a range of both practical and methodological considerations that may potentially affect the use of health care performance indicators. These considerations cannot be reduced solely to the analysis of indicators. For example, if an indicator is well-defined, collected and analyzed but is not presented optimally, what it signals may not be clear to the intended user for decision-making purposes. In other instances, it may be the data itself that is the main issue, because it cannot be linked, is of poor quality or incomplete. We have clustered five key factors and potentially relevant considerations depending on the intended purpose of use of an indicator that have been highlighted in the literature and by the first round of panelists.

1. Based on your use of health care performance indicators, what are your critical needs across the indicator cycle shown in Fig.A3? That is, for each stage in the cycle as shown, what would make the ideal conditions for your purposes?
2. Which stage in your opinion is most important for the actionability of an indicator?
3. Is the information currently used by your organization or practice useful for your purposes? That is, are you able to make decisions and learn from the information? How could it be better?
4. In your opinion, what are the main obstacles users face to make health care performance data actionable?

Figure A3. Use cycle of health care performance indicators

Appendix 3: Panelists

Expert panelists				User panelists			
#	Expertise (key words)	Literature consulted	Country	#	Organization type	Level	Country
1	Quality of care	[1-6]	Australia	1	Standards	Macro	United States
2	Performance measurement	[7-10]	United States	2	Governmental	Macro	Belgium
3	Performance measurement	[11-16]	Germany	3	Research/academia	Micro	United Kingdom
4	Quality of care	[1,2,5,6,17]	Australia	4	Standards	Meso	Canada
5	Governance	[18-20]	Canada	5	Governmental	Macro	Ireland
6	Performance measurement	[21-23]	Denmark	6	Governmental	Macro	Canada
7	Data/information	[24-26]	Netherlands	7	Health services	Meso	Germany
8	Management	[27-29]	Italy	8	Improvement	Macro	United States
9	Data/information systems	[30-32]	Canada	9	Governmental	Macro	Canada
10	Performance measurement	[33-36]	United Kingdom	10	Health services	Micro	Netherlands
11	Management	[37-39]	Netherlands	11	Research/academia	Meso	Netherlands
12	Quality of care	[40-42]	United States	12	Health services	Meso	United States
13	Performance measurement	[33,35,36,43]	United Kingdom	13	Standards	Meso	Netherlands
14	Governance	[44-46]	Denmark	14	Health services	Micro	Netherlands
15	Governance	[23,47,48]	Canada	15	Improvement	Macro	Netherlands
16	Data/information systems	[49-51]	Netherlands	16	Governmental	Macro	Canada

Appendix references

1. Braithwaite J, Hibbert P, Blakely B, Plumb J, Hannaford N, Long JC, et al. Health system frameworks and performance indicators in eight countries: A comparative international analysis. *SAGE Open Medicine*. 2017;5:2050312116686516.
2. Braithwaite J, Hibbert PD, Jaffe A, White L, Cowell CT, Harris MF, et al. Quality of Health Care for Children in Australia, 2012-2013. *Jama*. 2018;319(11):1113-24.
3. Mannion R, Braithwaite J. Unintended consequences of performance measurement in healthcare: 20 salutary lessons from the English National Health Service. *Intern Med J*. 2012;42(5):569-74.
4. Mumford V, Greenfield D, Hogden A, Debono D, Forde K, Westbrook J, et al. Development and application of an indicator assessment tool for measuring health services accreditation programs. *BMC Res Notes*. 2015;8:363.
5. Hibbert PD, Wiles LK, Cameron ID, Kitson A, Reed RL, Georgiou A, et al. CareTrack Aged: the appropriateness of care delivered to Australians living in residential aged care facilities: a study protocol. *BMJ Open*. 2019;9(6):e030988.
6. Runciman WB, Hunt TD, Hannaford NA, Hibbert PD, Westbrook JI, Coiera EW, et al. CareTrack: assessing the appropriateness of health care delivery in Australia. *Med J Aust*. 2012;197(2):100-5.
7. Damberg CL, Sorbero ME, Lovejoy SL, Lauderdale K, Wertheimer S, Smith A, et al. An Evaluation of the Use of Performance Measures in Health Care. *Rand health quarterly*. 2012;1(4):3-.
8. Damberg CL, Baker DW. Improving the Quality of Quality Measurement. *Journal of general internal medicine*. 2016;31 Suppl 1(Suppl 1):8-9.
9. Friedberg MW, Damberg CL. A Five-Point Checklist To Help Performance Reports Incentivize Improvement And Effectively Guide Patients. *Health Affairs*. 2012;31(3):612-8.
10. Parast L, Doyle B, Damberg CL, Shetty K, Ganz DA, Wenger NS, et al. Challenges in Assessing the Process–Outcome Link in Practice. *Journal of general internal medicine*. 2015;30(3):359-64.
11. Shaw C, Groene O, Berger E. Chapter 8: External institutional strategies: accreditation, certification, supervision. In: Busse R, Klazinga N, Panteli D, Quentin W, editors. *Improving healthcare quality in Europe: characteristics, effectiveness and implementation of different strategies* Copenhagen WHO Regional Office for Europe 2019.
12. Groene O. Improving the capacity for learning and improvement in health care. *International Journal for Quality in Health Care*. 2018;30(3):159-60.
13. Groene O, Sunol R. Patient involvement in quality management: rationale and current status. *J Health Organ Manag*. 2015;29(5):556-69.
14. Taylor A, Neuburger J, Walker K, Cromwell D, Groene O. How is feedback from national clinical audits used? Views from English National Health Service trust audit leads. *Journal of health services research & policy*. 2016;21(2):91-100.
15. Wagner C, Groene O, Thompson CA, Dersarkissian M, Klazinga NS, Arah OA, et al. DUQuE quality management measures: associations between quality management at hospital and pathway levels. *International journal for quality in health care : journal of the International Society for Quality in Health Care*. 2014;26 Suppl 1(Suppl 1):66-73.
16. Hilarion P, Suñol R, Groene O, Vallejo P, Herrera E, Saura RM. Making performance indicators work: the experience of using consensus indicators for external assessment of health and social services at regional level in Spain. *Health Policy*. 2009;90(1):94-103.
17. Hibbert Pea. Performance indicators used internationally to report publicly on healthcare organisations and local health systems. South Wales: Australian Institute of Health Innovation: University of New South Wales; 2013.
18. Lavis JN. Finding and using research evidence: Summary sheet. Hamilton, Canada: McMaster Health Forum 2019.
19. Fretheim A, Oxman AD, Lavis JN, Lewin S. SUPPORT Tools for Evidence-informed Policymaking in health 18: Planning monitoring and evaluation of policies. *Health research policy and systems*. 2009;7(1):S18.

20. Lavis JN, Hammill AC. Governance arrangements. In: JN L, editor. Ontario's health system: Key insights for engaged citizens, professionals and policymakers. Hamilton: McMaster Health Forum 2016.
21. Mainz J. Defining and classifying clinical indicators for quality improvement. *International Journal for Quality in Health Care*. 2003;15(6):523-30.
22. Mainz J. Developing evidence-based clinical indicators: a state of the art methods primer. *International Journal for Quality in Health Care*. 2003;15(suppl_1):i5-i11.
23. Carinci F, Van Gool K, Mainz J, Veillard J, Pichora EC, Januel JM, et al. Towards actionable international comparisons of health system performance: expert revision of the OECD framework and quality indicators. *International Journal for Quality in Health Care*. 2015;27(2):137-46.
24. Medlock S, Wyatt JC. Health Behaviour Theory in Health Informatics: Support for Positive Change. *Studies in health technology and informatics*. 2019;263:146-58.
25. Medlock S, Eslami S, Askari M, Arts DL, van de Glind EM, Brouwer HJ, et al. For which clinical rules do doctors want decision support, and why? A survey of Dutch general practitioners. *Health Informatics J*. 2019;25(3):1076-90.
26. Askari M, Wierenga PC, Eslami S, Medlock S, De Rooij SE, Abu-Hanna A. Studies pertaining to the ACOVE quality criteria: a systematic review. *International journal for quality in health care : journal of the International Society for Quality in Health Care*. 2012;24(1):80-7.
27. Nuti S. Let's play the patients music: A new generation of performance measurement systems in healthcare. *Management Decision*. 2018;56(10):2252-72.
28. Nuti S, De Rosis S, Bonciani M, Murante AM. Rethinking Healthcare Performance Evaluation Systems towards the People-Centredness Approach: Their Pathways, their Experience, their Evaluation. *Healthc Pap*. 2017;17(2):56-64.
29. Nuti S, Bini B, Ruggieri TG, Piaggese A, Ricci L. Bridging the Gap between Theory and Practice in Integrated Care: The Case of the Diabetic Foot Pathway in Tuscany. *International journal of integrated care*. 2016;16(2):9-.
30. OECD. Health in the 21st Century: putting data to work for stronger health systems. Paris: OECD; 2019.
31. OECD. Health data governance: privacy, monitoring and research - policy brief Paris: OECD; 2015.
32. OECD. Strengthening health information infrastructure for health care quality governance: good practices, new opportunities and data privacy challenges Paris: OECD 2013.
33. Smith P, Mossialos E, Papanicolas I. Performance measurement for health system improvement: experiences, challenges and prospects. Copenhagen: WHO Regional Office for Europe; 2008.
34. Papanicolas I, Jha AK. Challenges in International Comparison of Health Care Systems. *JAMA*. 2017;318(6):515-6.
35. Smith P, Mossialos E, Papanicolas I, Leatherman S. Principles of performance measurement In: Smith P, Mossialos E, Papanicolas I, Leatherman S, editors. Performance measurement for health system improvement Cambridge: Cambridge University Press; 2009.
36. Papanicolas I, Smith PC. Health system performance comparison: An agenda for policy, information and research Maidenhead: McGraw-Hill 2013.
37. Rotar AM, van den Berg MJ, Kringos DS, Klazinga NS. Reporting and use of the OECD Health Care Quality Indicators at national and regional level in 15 countries. *International journal for quality in health care : journal of the International Society for Quality in Health Care*. 2016;28(3):398-404.
38. Rotar AM, van den Berg MJ, Klazinga NS. An expert-based mapping of healthcare system strategies to support rational drug prescribing in primary care across 13 European countries. *Health research policy and systems*. 2020;18(1):102.
39. Rotar AM, Botje D, Klazinga NS, Lombarts KM, Groene O, Sunol R, et al. The involvement of medical doctors in hospital governance and implications for quality management: a quick scan in 19 and an in depth study in 7 OECD countries. *BMC Health Serv Res*. 2016;16 Suppl 2(Suppl 2):160.
40. Van der Wees PJ, Nijhuis-van der Sanden MWG, van Ginneken E, Ayanian JZ, Schneider EC, Westert GP. Governing healthcare through performance measurement in Massachusetts and the Netherlands. *Health Policy*. 2014;116(1):18-26.

41. Van Der Wees PJ, Nijhuis-Van Der Sanden MWG, Ayanian JZ, Black N, Westert GP, Schneider EC. Integrating the Use of Patient-Reported Outcomes for Both Clinical Practice and Performance Measurement: Views of Experts from 3 Countries. *The Milbank quarterly*. 2014;92(4):754-75.
42. Geissler KH, Friedberg MW, SteelFisher GK, Schneider EC. Motivators and barriers to using patient experience reports for performance improvement. *Med Care Res Rev*. 2013;70(6):621-35.
43. Smith PC, Anell A, Busse R, Crivelli L, Healy J, Lindahl AK, et al. Leadership and governance in seven developed health systems. *Health Policy*. 2012;106(1):37-49.
44. Barbazza E, Tello J. A review of health governance: Definitions, dimensions and tools to govern *Health Policy* 2014;116:1-11.
45. Tello JE, Barbazza E, Waddell K. Review of 128 quality of care mechanisms: A framework and mapping for health system stewards. *Health Policy*. 2020;124(1):12-24.
46. Tello J, Baez-Camargo C. Strengthening health system accountability: a WHO European Region multi-country study. Copenhagen WHO Regional Office for Europe 2015.
47. Veillard JH, Brown AD, Baris E, Permanand G, Klazinga NS. Health system stewardship of National Health Ministries in the WHO European region: concepts, functions and assessment framework. *Health Policy*. 2011;103(2-3):191-9.
48. Veillard J, Champagne F, Klazinga N, Kazandjian V, Arah OA, Guisset AL. A performance assessment framework for hospitals: the WHO regional office for Europe PATH project. *International journal for quality in health care : journal of the International Society for Quality in Health Care*. 2005;17(6):487-96.
49. Verheij RA, Curcin V, Delaney B, McGilchrist MM. Possible sources of bias in primary care electronic health record data use and reuse *Journal of medical internet research*. 2018;20(5).
50. Verheij R, Witvliet C, Jansen T, Hooiveld M, Hilten O van. Reuse of routine care data for policy and science: how things can be improved [Dutch]. Utrecht: Nivel; 2019.
51. van der Bij S, de Hoon S, Nielen M, de Jong A, de Boer D, Verheij R. Routine recorded care data as a source of quality information [Dutch]. Netherlands Nivel; 2016.