

# Outcome differences between surgeons performing first and subsequent coronary artery bypass grafting procedures in a day: a retrospective comparative cohort study

Danwei Zhang,<sup>1</sup> Dachuan Gu,<sup>1,2</sup> Chenfei Rao,<sup>1,2</sup> Heng Zhang,<sup>1,2</sup> Xiaoting Su,<sup>1</sup> Sipeng Chen,<sup>1</sup> Hanping Ma,<sup>1</sup> Yan Zhao,<sup>1</sup> Wei Feng,<sup>1,2</sup> Hansong Sun,<sup>1,2</sup> Zhe Zheng <sup>1,2,3</sup>

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2021-014244>).

For numbered affiliations see end of article.

## Correspondence to

Dr Zhe Zheng, National Clinical Research Center for Cardiovascular Diseases, State Key Laboratory of Cardiovascular Disease, and Department of Cardiovascular Surgery, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, No.167 North Lishi Road, Xicheng District, Beijing, 100037, People's Republic of China, Beijing, People's Republic of China; [zhengzhe@fuwai.com](mailto:zhengzhe@fuwai.com)

DZ and DG are joint first authors.

Received 14 September 2021  
Accepted 13 May 2022  
Published Online First  
1 June 2022



► <http://dx.doi.org/10.1136/bmjqs-2022-015045>



© Author(s) (or their employer(s)) 2023. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Zhang D, Gu D, Rao C, et al. *BMJ Qual Saf* 2023;**32**:192–201.

## ABSTRACT

**Background** With increasing surgical workload, it is common for cardiac surgeons to perform coronary artery bypass grafting (CABG) after other procedures in a workday. To investigate whether prior procedures performed by the surgeon impact the outcomes, we compared the outcomes between CABGs performed first versus those performed after prior procedures, separately for on-pump and off-pump CABGs as they differed in technical complexity.

**Methods** We conducted a retrospective cohort study of patients undergoing isolated CABG in China from January 2013 to December 2018. Patients were categorised as undergoing on-pump and off-pump CABGs. Outcomes of the procedures performed first in primary surgeons' daily schedule (first procedure) were compared with subsequent ones (non-first procedure). The primary outcome was an adverse events composite (AEC) defined as the number of adverse events, including in-hospital mortality, myocardial infarction, stroke, acute kidney injury and reoperation. Secondary outcomes were the individual components of the primary outcome, presented as binary variables. Mixed-effects models were used, adjusting for patient and surgeon-level characteristics and year of surgery.

**Results** Among 21 866 patients, 10 109 (16.1% as non-first) underwent on-pump and 11 757 (29.6% as non-first) off-pump CABG. In the on-pump cohort, there was no significant association between procedure order and the outcomes (all  $p>0.05$ ). In the off-pump cohort, non-first procedures were associated with an increased number of AEC (adjusted rate ratio 1.29, 95% CI 1.13 to 1.47,  $p<0.001$ ), myocardial infarction (adjusted OR (OR<sub>adj</sub>) 1.43, 95% CI 1.13 to 1.81,  $p=0.003$ ) and stroke (OR<sub>adj</sub> 1.73, 95% CI 1.18 to 2.53,  $p=0.005$ ) compared with first procedures. These increases were only found to be statistically significant when the procedure was performed by surgeons with <20 years' practice or surgeons with a preindex volume <700 cases.

**Conclusions** For a technically challenging surgical procedure like off-pump CABG, prior workload adversely affected patient outcomes.

## WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Increased workload adversely affects surgical outcomes. However, whether prior procedures performed by the surgeon in a workday impact the outcomes of on-pump and the technically more complex off-pump coronary artery bypass grafting (CABG) remains unclear.

## WHAT THIS STUDY ADDS

⇒ For off-pump CABG, prior surgical workload has negative impacts on patient outcomes, particularly if procedures were performed by less experienced or lower volume surgeons. These findings were consistent across several patient subgroups and multiple sensitivity analyses.

## HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE AND/OR POLICY

⇒ Daily schedules should ensure that complex procedures are assigned first in a surgeon's shift, while taking patient and surgeon characteristics into account.

## INTRODUCTION

With increasing surgical workload, surgeons often have multiple operations in a day, especially in countries where surgical resources are relatively scarce.<sup>1–3</sup> Extended work hours and excessive

workload have been proved to bring negative influences on patient outcomes.<sup>4-7</sup> Therefore, various measures to evaluate workload and strategies to reduce workload impacts have been suggested.<sup>8-9</sup> However, whether prior procedures performed by the surgeon in a workday will affect surgical outcomes remains unclear, which could be instructive in arranging surgeons' daily workflow patterns.

Coronary artery bypass grafting (CABG), the standard of care for patients with extensive coronary artery disease, has been among the most common and major surgical procedures.<sup>10</sup> In a CABG procedure, the most critical and challenging step is performing a surgical anastomosis between autologous vessel grafts and coronary arteries, which requires extremely high mental concentration and great physical strength of the surgeon. To achieve the precision requirement for a successful CABG, the heart is typically arrested with a cardiopulmonary bypass (CPB) machine to support the circulation. The off-pump procedure, however, has also evolved as another option that the anastomosis is performed on a beating heart without CPB,<sup>11</sup> which avoids the adverse effect from cardiac arrest and CPB, but adds to the complexity and difficulty of the procedure. Thus, the off-pump CABG is considered to be more technically challenging than the on-pump CABG (with CPB).<sup>12-14</sup> Considering on-pump and off-pump CABGs are two procedures with similar surgical process but different complexity, prior workload may have varying impacts on them. It is highly necessary to examine the association between the procedure order and outcomes, with on-pump and off-pump CABGs regarded as two typical clinical scenarios.

The purpose of this study was to examine whether intraday workload (ie, prior procedures conducted on the same day) would affect the outcomes after CABG, accounting for surgical complexity. We compared the outcome differences between the first CABG and subsequent ones performed in a day, for on-pump CABG and off-pump CABG separately. We also investigated the effect across different patient health conditions and surgeon experience.

## METHODS

### Study design

Data of patients undergoing CABG at a cardiovascular specialty hospital with the annual CABG volume exceeding 4500 cases in China from 1 January 2013 through 31 December 2018 were collected through the hospital electronic health record (EHR) system. Data audit and quality measures have been reported previously.<sup>15-16</sup> Briefly, we established an EHR-derived database by linking the structured EHR to a registry database to enable real-time data collection, including baseline patient characteristics, surgical records and in-hospital outcomes. To assure the accuracy of the linkage, 5% of patients were randomly selected and all variables of these patients were compared between

the original EHR and EHR-derived database, and the EHR-derived data were 100% consistent with the original records.

### Study population

This study included all consecutive patients aged 18 years or older who underwent isolated, elective CABG, which meant that patients undergoing emergent CABG, CABG combined with valvular surgeries or other cardiovascular surgeries were not included. All the CABG surgeries were performed according to the operating standards (online supplemental method 1). Patients who were converted from off-pump to on-pump procedure during the surgery (n=736) were excluded as their outcomes were influenced by both procedures, and it was inappropriate to include them into either cohort. Other exclusion criteria included: (1) patients treated by surgeons after night shift in the preceding overnight hours (00:00 to 07:00, n=175) to avoid bias from extra preceding workload; and (2) patients without surgeon information (n=4). Patients were categorised as undergoing on-pump and off-pump CABGs by the use of CPB, and the association between intraday procedure order and patient outcomes was assessed in both CABG cohorts separately.


### Procedure order and study group

Procedure order was determined within work hours (from 08:00 to 19:00) in a single day, when all the elective CABG procedures were performed. All surgical procedures (included but not limited to CABG) performed by the surgeon were taken into account to determine the order, regardless of whether the surgeon served as a primary surgeon or an assistant. More details on the duties of the primary surgeon and surgical team during the surgery are described in online supplemental method 1.

'First procedure' was defined as the CABG surgery being the first procedure of the primary surgeon in a day, whereas other CABG surgeries were also assigned their number in the ordering but included as the 'non-first procedure' considering the lower volumes observed for the third or later procedures (figure 1). Thus, in both on-pump and off-pump CABG cohorts, the patients were further divided into 'first procedure' and 'non-first procedure' groups separately.

### Outcome measures

The primary outcome was an adverse events composite (AEC), including in-hospital all-cause mortality, non-fatal myocardial infarction (MI), non-fatal stroke, acute kidney injury (AKI) and reoperation, which are widely used for evaluating the quality of CABG procedures.<sup>17-19</sup> The AEC was expressed as the number of adverse events that occurred during the index hospitalisation, and the analysis was at the individual patient level with each type of the adverse event being counted only once.<sup>20</sup> Secondary outcomes



**Figure 1** Study flow chart. The flow chart shows detailed inclusion and exclusion criteria for this study as well as corresponding patient numbers.

\*Multiple exclusions may apply to one patient so that the number in each criterion adds up greater than the total ineligible or excluded patients in the grey boxes. CABG, coronary artery bypass grafting.

included the individual components of the primary outcome mentioned above, expressed as binary variables (yes/no). MI was defined as an absolute rise in cardiac troponin to over 70 times the 99th percentile upper reference limit (URL), or an increase in creatine kinase - muscle/brain (CK-MB) to over 10 times the 99th percentile URL. Stroke was defined as brain, spinal cord or retinal cell death attributable to focal arterial ischaemia. AKI was defined as any one of the following: (1) an increase in the serum creatinine level over three times the latest preoperative value; (2) a serum creatinine level reaching 4.0 mg/dL with a minimum increase of 0.5 mg/dL relative to the last preoperative value; and (3) a new postoperative dialysis. Reoperation was defined as the re-exploration for mediastinal bleeding either in the intensive care unit (ICU) or operating room. Detailed definitions of the outcomes are further described in online supplemental method 2.

#### Patient and surgeon characteristics

Patient characteristics were collected and some were selected as covariates for risk adjustment because they were previously shown to be risk factors among patients undergoing CABG in China,<sup>21</sup> including (1) demographic characteristics: age, sex; (2) comorbidities: body mass index, angina, estimated glomerular filtration rate (eGFR); (3) medical history: history of chronic obstructive pulmonary disease, MI, stroke,

cardiac surgery, percutaneous coronary intervention; and (4) cardiovascular status: New York Heart Association (NYHA) class, preoperative left ventricular ejection fraction (LVEF), left main disease, number of lesions and aortic calcification. Covariates for each model are listed in online supplemental table 2. Other characteristics that were not included in the multivariable regression models (following the general principle of variable selection, like sample size, collinearity or clinical importance) but were often reported in other cardiovascular studies were also listed in table 1, and they were also used to predict the propensity score in the sensitivity analysis (see below).

All surgeons performed first and non-first CABG procedures. Surgeon characteristics including years in practice and CABG volume before the index procedure were calculated dynamically for each patient as surgeon-level covariates. Years in practice was defined as the number of years a surgeon served as a cardiac surgeon until the date of the index CABG procedure. The CABG volume was defined as the cumulative number of CABGs a surgeon had performed from the year 2012 to the date of the index CABG procedure. The annual CABG volume in 2012 was included to indicate a surgeon's surgical experience at baseline.

#### Statistical analyses

We performed the analyses in the on-pump and off-pump CABG cohorts separately. The patient and

**Table 1** Patient demographic and clinical characteristics

	On-pump CABG cohort (n=10 109)				Off-pump CABG cohort (n=11 757)		
	Overall	First procedure (n=8480)	Non-first procedure (n=1629)	P value*	First procedure (n=8279)	Non-first procedure (n=3478)	P value*
Age, mean (SD), years	61.2 (8.6)	60.8 (8.5)	60.5 (8.5)	0.33	61.7 (8.8)	61.2 (8.6)	<b>0.005</b>
Female, n (%)	4975 (22.8)	1893 (22.3)	388 (23.8)	0.19	1887 (22.8)	807 (23.2)	0.63
BMI, mean (SD), kg/m <sup>2</sup> †	25.7 (3.0)	25.8 (3.0)	25.7 (3.1)	0.94	25.6 (3.0)	25.6 (3.1)	0.65
Smoking, n (%)	12 117 (55.4)	4578 (54.0)	866 (53.2)	0.54	4683 (56.6)	1990 (57.2)	0.51
Diabetes, n (%)	4952 (22.6)	2017 (23.8)	353 (21.7)	0.06	1820 (22.0)	762 (21.9)	0.93
Hypertension, n (%)	13 957 (63.8)	5418 (63.9)	1069 (65.6)	0.18	5291 (63.9)	2179 (62.7)	0.20
Hyperlipidaemia, n (%)	16 128 (73.8)	6178 (72.9)	1176 (72.2)	0.58	6172 (74.6)	2602 (74.8)	0.76
Prior COPD, n (%)	139 (0.6)	44 (0.5)	10 (0.6)	0.63	57 (0.7)	28 (0.8)	0.50
Prior MI, n (%)	8189 (37.5)	3209 (37.8)	633 (38.9)	0.44	3085 (37.3)	1262 (36.3)	0.32
Prior stroke, n (%)	1684 (7.7)	632 (7.5)	122 (7.5)	0.96	648 (7.8)	282 (8.1)	0.61
Prior PCI, n (%)	1233 (5.6)	433 (5.1)	89 (5.5)	0.55	523 (6.3)	188 (5.4)	0.06
Prior cardiac surgery, n (%)	828 (3.8)	322 (3.8)	59 (3.6)	0.73	309 (3.7)	138 (4.0)	0.54
Angina, n (%)	18 491 (84.6)	7032 (82.9)	1352 (83.0)	0.94	7061 (85.3)	3046 (87.6)	<b>0.001</b>
NYHA III/IV, n (%)	6923 (31.7)	2367 (27.9)	502 (30.8)	<b>0.017</b>	2809 (33.9)	1245 (35.8)	0.05
eGFR, median (IQR), mL/min/1.73 m <sup>2</sup> ‡	91.6 (78.1–107.0)	92.0 (78.4–107.5)	91.2 (78.4–106.6)	0.41	91.5 (77.8–106.5)	91.2 (78.4–107.2)	0.40
PAD, n (%)	2790 (12.8)	881 (10.4)	146 (9.0)	0.08	1194 (14.4)	569 (16.4)	<b>0.007</b>
LM disease, n (%)	4145 (19.0)	1580 (18.6)	284 (17.4)	0.25	1633 (19.7)	648 (18.6)	0.17
Three-vessel disease, n (%)	13 296 (60.8)	5692 (67.1)	1116 (68.5)	0.27	4507 (54.4)	1981 (57.0)	<b>0.012</b>
Preoperative LVEF, mean (SD)	61.1 (7.4)	60.9 (7.6)	60.6 (7.8)	0.16	61.3 (7.2)	61.5 (6.9)	0.17
Aortic calcification, n (%)	1774 (8.1)	437 (5.2)	110 (6.8)	<b>0.009</b>	863 (10.4)	364 (10.5)	0.95
EuroSCORE >3, n (%)	4990 (22.8)	1727 (20.4)	304 (18.7)	0.33	2075 (25.1)	884 (25.4)	0.69

\*The  $\chi^2$  test is used for categorical variables and the Kruskal-Wallis test is used for continuous variables between first and non-first procedure groups in specific CABG cohorts. Those achieving a statistically significant level of 0.05 are in bold.

†BMI=weight (kg)/(height [m])<sup>2</sup>.

‡eGFR (mL/min/1.73 m<sup>2</sup>)=186×(serum creatinine level [mg/dL])<sup>-1.154</sup>×(age [years])<sup>-0.203</sup>.

BMI, body mass index; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; EuroSCORE, European System for Cardiac Operative Risk Evaluation; LM, left main; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NYHA, New York Heart Association; PAD, peripheral artery disease; PCI, percutaneous coronary intervention.

surgeon characteristics between first and non-first procedure groups were compared, with the  $\chi^2$  test used for categorical variables and the Kruskal-Wallis test for continuous variables. The completeness of patient-level variables ranged from 96% to 100% (online supplemental table 1). For those key variables with missing data, we implemented multiple imputation with the Markov Chain Monte Carlo method, and the predictors included other variables of patient demographic characteristics, comorbidities and medical history with 100% completeness listed in online supplemental table 1. The final imputed value was an average of 20 imputations for all analyses.

To investigate the AEC difference between first and non-first procedure groups, assuming a Poisson distribution for the number of adverse events, we fitted a mixed-effects model with a Poisson link function and surgeon-specific random intercepts to model the AEC as a function of procedure order (non-first procedure vs first procedure; when examining the potential 'dose-effect' relationship between the number of prior procedures performed by the surgeon and patient outcomes, we compared the third or later procedure, the second procedure vs the first procedure). Rate ratios (RR) were calculated to present the outcome differences. For each individual adverse event (binary outcome),

we fitted separate mixed-effects models with a logit link function and surgeon-specific random intercepts, with ORs estimated. All models were adjusted for patient and surgeon-level characteristics listed in the online supplemental table 2, also including an interval time variable corresponding to the date of surgery (2013 (time=0) through 2018 (time=6)) to represent annual changes in outcomes.

To test whether prior workload had greater impact in clinically high-risk subsets,<sup>10 21</sup> we repeated the analysis for AEC in eight prespecified patient subgroups defined by age (<65 or ≥65 years), sex, diabetes, eGFR (<60 or ≥60 mL/min/1.73 m<sup>2</sup>), LVEF (<50% or ≥50%), prior MI, three-vessel disease and European System for Cardiac Operative Risk Evaluation (≤3 or >3). Since surgeons' skill and experience may overwhelm the effect of workload, we further repeated the analyses in two surgeon subgroups defined by years in practice (<20 or ≥20 years), and the CABG volume (<700 or ≥700 cases) before the index procedure.

### Sensitivity analyses

We performed several sensitivity analyses to test the robustness of our findings. First, we aimed to take into account possible confounding by indication as some high-risk patients may be more likely performed as a non-first procedure. We therefore included all patient characteristics (online supplemental table 1) in a multivariable logistic regression model to estimate the propensity for patients to undergo a non-first procedure. The inverse probability of treatment weighting (IPTW) was used based on the inverse of the propensity score and the mixed-effects model analyses repeated. We also performed a 1:1 propensity score matching to form matched on-pump and off-pump CABG cohorts by matching first and non-first patients treated by the same surgeon with similar propensity score using the greedy nearest neighbour matching algorithm, with a calliper width at 0.01 according to the absolute value of the difference between propensity scores. Second, we took the surgeon and assistant as pairs and restricted the analysis to the pairs who had ever performed both first and non-first procedures, with at least five CABG cases in total. The pair identifiers were used as the random intercepts in mixed-effects models rather than the surgeon, thus reducing bias from assistants. Finally, we restricted the analysis to the surgeries that started after 11:00 in case that circadian variation of patients or surgical teams could influence the outcomes.

All statistical tests were two sided, with a significance level of 0.05. We conducted all the analyses using SAS software V.9.4 (SAS Institute).

## RESULTS

### Study cohorts

Among 27 266 consecutive patients undergoing CABG during the study period, 21 866 were included in the final analysis (table 1). The mean age was 61.2 (SD

8.6) years and 22.8% were female. There were 10 109 patients in the on-pump cohort and 16.1% underwent non-first procedures; 11 757 patients in the off-pump cohort and 29.6% underwent non-first procedures.

In the on-pump CABG cohort, patients in the non-first procedure group had higher prevalence of NYHA III/IV (non-first vs first: 30.8% vs 27.9%) and aortic calcification (6.8% vs 5.2%) compared with the first procedure group, indicating higher risk of worse patient outcomes. In the off-pump CABG cohort, patients who underwent non-first procedures were younger (61.2 vs 61.7 years), and had higher prevalence of angina (87.6% vs 85.3%), peripheral artery disease (16.4% vs 14.4%) and three-vessel disease (57.0% vs 54.4%) than those who underwent a first procedure. Apart from the above, patient characteristics were similar between the first and non-first procedures in the on-pump and off-pump CABG cohorts (table 1).

Overall, 47 primary surgeons were included. Surgeons in the non-first procedure group had higher preindex CABG volume in both on-pump and off-pump CABG cohorts, and longer years in practice in the off-pump cohort ( $p<0.001$ , online supplemental table 3).

### Primary outcome

In the on-pump CABG cohort, 8480 patients undergoing first procedures had 1173 AEC in total, and 1629 patients undergoing non-first procedures had 208 AEC (table 2 and online supplemental figure 1). The number of adverse events per 1000 hospitalisations was 127.7 (non-first) vs 138.3 (first). There was no significant difference in AEC between the two groups after risk adjustment (adjusted rate ratio (RR<sub>adj</sub>) 0.90, 95% CI 0.77 to 1.06,  $p=0.19$ ; table 2).

In the off-pump CABG cohort, 8279 patients undergoing first procedures had 759 adverse events in total (patients with AEC: one event, 658; two events, 44; ≥three events, 4), and 3478 patients undergoing non-first procedures had 390 events (one event, 319; two events, 28; three events, 5). The number of adverse events per 1000 hospitalisations was 112.1 (non-first) vs 91.7 (first). After risk adjustment, undergoing a non-first procedure was associated with an increased number of AEC (RR<sub>adj</sub> 1.29, 95% CI 1.13 to 1.47,  $p<0.001$ ; table 2). Further stratifying non-first procedures, it was shown that the third or later procedure (accounting for 6.9% of all non-first procedures) was associated with a higher rate (RR<sub>adj</sub> 1.35, 95% CI 0.89 to 2.04,  $p=0.16$ ) relative to first procedures than second procedures (RR<sub>adj</sub> 1.28, 95% CI 1.12 to 1.46,  $p<0.001$ ), but did not reach a statistical significance (online supplemental figure 2).

### Secondary outcomes

In the on-pump CABG cohort, there was no statistically significant difference between the first and

**Table 2** Outcome differences associated with non-first versus first procedure groups in the on-pump and off-pump CABG cohorts

Outcome	On-pump CABG cohort (n=10 109)			P value	Off-pump CABG cohort (n=11 757)			P value
	First procedure (n=8480)	Non-first procedure (n=1629)	RR <sub>adj</sub> <sup>*</sup> /OR <sub>adj</sub> <sup>†</sup> (95% CI)		First procedure (n=8279)	Non-first procedure (n=3478)	RR <sub>adj</sub> <sup>*</sup> /OR <sub>adj</sub> <sup>†</sup> (95% CI)	
AEC, n (%) <sup>*</sup>			0.90 (0.77 to 1.06)	0.19			1.29 (1.13 to 1.47)	<b>&lt;0.001</b>
One event	984 (11.6)	165 (10.1)			658 (7.9)	319 (9.2)		
Two events	71 (0.8)	17 (1.0)			44 (0.5)	28 (0.8)		
Three events	13 (0.2)	3 (0.2)			3 (0.0)	5 (0.1)		
Four events	2 (0.0)	0 (0.0)			1 (0.0)	0 (0.0)		
Death, n (%)	32 (0.4)	10 (0.6)	1.44 (0.79 to 2.62)	0.23	16 (0.2)	12 (0.3)	2.01 (0.94 to 4.27)	0.07
MI, n (%)	379 (4.5)	71 (4.4)	0.84 (0.64 to 1.09)	0.19	204 (2.5)	114 (3.3)	1.43 (1.13 to 1.81)	<b>0.003</b>
Stroke, n (%)	98 (1.2)	11 (0.7)	0.57 (0.31 to 1.06)	0.08	57 (0.7)	40 (1.2)	1.73 (1.18 to 2.53)	<b>0.005</b>
AKI, n (%)	559 (6.6)	98 (6.0)	0.93 (0.72 to 1.21)	0.60	362 (4.4)	178 (5.1)	1.23 (1.00 to 1.51)	0.054
Reoperation, n (%)	105 (1.2)	18 (1.1)	0.83 (0.50 to 1.38)	0.48	120 (1.4)	46 (1.3)	0.92 (0.64 to 1.32)	0.65

AEC indicates adverse events composite, which is defined as the number of any adverse events that occurred, including death, myocardial infarction, stroke, acute kidney injury or reoperation.

<sup>\*</sup>For the adverse events composite (AEC, count variable), the 'n (%)' presents the number of patients with given number of events and the percentage of those patients among specific procedure group. A mixed-effects model with a Poisson link function and surgeon-specific random intercepts is fitted, adjusting for the patient risk factors and surgeon characteristics mentioned in the Methods section and listed in online supplemental table 2. The adjusted rate ratio (non-first vs first) is estimated, and when it achieves a statistically significant level of 0.05, the p value is in bold.

<sup>†</sup>For every individual event (binary variable; ie, death, MI, stroke, AKI and reoperation), a mixed-effects model with a logit link function and surgeon-specific random intercepts is fitted, also adjusting for the patient and surgeon characteristics listed in online supplemental table 2. The adjusted OR (non-first vs first) for each outcome is estimated, and when it achieves a statistically significant level of 0.05, the p value is in bold.

AKI, acute kidney injury; CABG, coronary artery bypass grafting; MI, myocardial infarction; OR<sub>adj</sub>, adjusted OR; RR<sub>adj</sub>, adjusted rate ratio.


non-first procedure groups for any individual outcomes (table 2).

In the off-pump CABG cohort, non-first procedure group was associated with higher odds of MI (3.3% vs 2.5%, adjusted OR (OR<sub>adj</sub>) 1.43, 95% CI 1.13 to 1.81, p=0.003) and stroke (1.2% vs 0.7%, OR<sub>adj</sub> 1.73, 95% CI 1.18 to 2.53, p=0.005) than the first procedure

group, but were similar in both groups for all other outcomes (death, AKI and reoperation) (table 2).

### Subgroup analyses

The AEC did not differ between first and non-first procedures in the on-pump CABG cohort among all the patient or surgeon subgroups (figure 2). However,



**Figure 2** Association between procedure order and adverse events composite (AEC) in the patient and surgeon subgroups of the on-pump and off-pump CABG cohorts. The forest plots show the subgroup-specific RRs for the AEC in patients of non-first procedure group compared with patients of first procedure group, adjusting for the patient and surgeon characteristics. Panels (A) and (B) show the subgroup-specific RRs in the on-pump and off-pump CABG cohorts, respectively. A RR of greater than 1 indicates an increased number of adverse events for non-first procedure group in specific subgroups. The horizontal lines indicate 95% CIs. CABG, coronary artery bypass grafting; EuroSCORE, European System for Cardiac Operative Risk Evaluation; eGFR, estimated glomerular filtration rate; LVEF, left ventricular ejection fraction; MI, myocardial infarction; RR, rate ratio.

in the off-pump CABG cohort, the number of AEC increased significantly among most patient subgroups, except for patients with diabetes, or with eGFR <60 mL/min/1.73 m<sup>2</sup>. In surgeon subgroups, a non-first procedure was associated with an increased AEC only when the procedure was performed by surgeons with <20 years in practice (RR<sub>adj</sub> 1.87, 95% CI 1.24 to 2.82, p=0.003) or a preindex CABG volume <700 cases (RR<sub>adj</sub> 1.35, 95% CI 1.04 to 1.76, p=0.02).

### Sensitivity analyses

The results were robust and confirmed the primary findings across all sensitivity analyses. In the IPTW-weighted cohorts, similar results were obtained on an increased AEC (RR<sub>adj</sub> 1.28, 95% CI 1.14 to 1.45, p<0.001) associated with undergoing a non-first procedure in the off-pump CABG. Moreover, the odds of death and AKI were now significantly increased in addition to increased odds on MI and stroke (online supplemental table 5). In the 1:1 propensity score-matched cohorts (3084 patients in the on-pump CABG cohort and 6882 in the off-pump), patient characteristics were well balanced between matched first and non-first procedure groups (online supplemental table 4). Non-first procedures were also associated with increased AEC (RR<sub>adj</sub> 1.32, 95% CI 1.13 to 1.54, p<0.001), MI and stroke in the off-pump cohort (online supplemental table 5). For the analysis based on 316 surgeon–assistant pairs, including 8600 on-pump and 11 185 off-pump CABGs, the non-first procedure was associated with increased AEC (RR<sub>adj</sub> 1.29, 95% CI 1.14 to 1.47, p<0.001; online supplemental table 6), death, MI and stroke in the off-pump cohort. Among 4992 patients undergoing off-pump CABG after 11:00, the non-first group was also associated with increased AEC (RR<sub>adj</sub> 1.30, 95% CI 1.04 to 1.64, p=0.02; online supplemental table 7) and MI. As for reoperation, no significant association was observed in any of the sensitivity analyses. Besides, in the on-pump cohort, undergoing a non-first procedure was associated with a significantly lower odds on stroke in several analyses (online supplemental tables 5 and 6).

### DISCUSSION

In this retrospective comparative study, procedure order of the surgeon in a day did not impact the outcomes after on-pump CABG, but patients undergoing a non-first off-pump CABG procedure had increased adverse events compared with those undergoing a first procedure. The results for off-pump CABG were found across most patient subgroups but only for less experienced and lower volume surgeons, as well as being robust across multiple sensitivity analyses.

Outcome differences among scheduled surgeries during routine work hours should not be neglected, especially when increased workload or inappropriate

arrangement could be possible reasons. Nevertheless, evidence is still scarce on surgeons' prior workload impact on subsequent procedures, with inconclusive results of existing studies.<sup>7 22–24</sup> Lee *et al*<sup>22</sup> found that preceding endoscopic procedures impacted polyp detection, independent of surgical time, and detection rates declined as time passed during an endoscopist's schedule. However, Bagrodia *et al*<sup>7</sup> demonstrated that performing several urological procedures consecutively was irrelevant to worse outcomes. Gruskay *et al*<sup>23</sup> observed that postoperative infection was associated with the case order of decompression procedures but not for fusion procedures in spine surgery. The controversy may partly be attributed to the procedure type and selected outcome indicators. Nevertheless, all these studies implied the importance of daily schedule drafting and rational work arrangement.

Our study took CABG as a specific clinical scenario, one of the complicated but protocolised cardiac surgeries, to study whether prior procedures could impact patient outcomes. The procedure order in a day for a surgeon was regarded as a metric to study surgeon workload and workflow pattern. Additionally, we tested our hypothesis in the on-pump and off-pump CABGs separately, giving full consideration to the technical differences. Meaningful results were found that for off-pump CABG, surgical performance was influenced by preceding procedures performed by the surgeon, as it was associated with an increased number of adverse events. Moreover, the outcome of the third or later procedure may deteriorate further, as suggested by the higher point estimate although it did not reach statistical significance in the current sample. But for on-pump CABG, there was no significant association between the procedure order and outcomes. We considered the explanation for the difference between procedures may be the high technical demand and concentration required for off-pump CABG made it more susceptible to be affected by the fatigue status of primary surgeons. It is reasonable to assume that physical and cognitive fatigue have greater detrimental impact on the performance of surgeons who have performed more procedures before the index surgery. Apart from the primary composite endpoint, non-first procedures were also associated with higher odds of MI and stroke after off-pump CABG, the most devastating complications and common causes of early and late mortality after CABG.<sup>25–27</sup> As there is growing research investigating the surgery time or operative hours on surgical outcomes,<sup>28–30</sup> our study adds new evidence on evaluating the workload impact during regular work hours.

The findings were robust across multiple sensitivity analyses. Both the IPTW and the propensity score matching method to reduce confounding by indication resulted in estimates very close to the primary analysis. For concerns on the possible bias from the assistant, surgeon–assistant pairs were used as random

effects instead of primary surgeons alone, and results were consistent. As for other surgical team members like anaesthesiologists, nurses or perfusionists, they were scheduled for the entire day and hardly changed during the daytime surgeries. Considering that inconclusive discussions remain on the natural circadian variability in surgical performance,<sup>31–34</sup> the analysis on surgeries performed after 11:00 demonstrated the procedure order still had a consistent negative impact on outcomes after off-pump CABG. Additionally, higher odds of death and AKI also reached statistical significance in some sensitivity analyses. Larger sample size is needed to confirm the findings particularly for events that occur infrequently such as death. As for reoperation, we hypothesise that the lack of a significant association among primary and sensitivity analyses was because reoperation-related processes, such as sternal closure and haemostasis, were performed by assistant rather than primary surgeons. We also found that in the on-pump cohort, non-first procedure was associated with lower odds of stroke in some sensitivity analyses. This should be interpreted with caution since some risk factors for stroke after on-pump CABG such as aorta atheroma or carotid artery disease were not recorded in our study, and may bias the result.

The subgroup analyses further enriched our findings. In most high-risk subsets, the primary findings were consistent for both on-pump and off-pump CABGs, which ensured that the effect of procedure order was not driven by a narrow patient group. The association between procedure order and outcomes after off-pump CABG was non-significant in patients with diabetes or low eGFR. Since these subgroups indicated more severe patients, we hypothesise the outcomes were largely and primarily determined by patient conditions so that the effect of procedure order was relatively small in that context. Interesting findings were observed in surgeon-specific subgroup analyses, as the association was only found for less experienced and lower volume surgeons, which indicates that accumulating surgical experience and skills might outweigh the adverse effect of workload.

There are some clinical implications of our study. Surgeons performing multiple surgeries in a day is common in clinical settings. The number of prior procedures can be a new metric of workload, and may have similar impact on outcomes among other clinical specialisations or surgical procedures. Our study suggests that better arrangement on daily schedules can be an actionable and cost-effective approach to balance the workload without major changes on existing infrastructure, for example, prioritising more difficult and complex procedures first, while also taking patient characteristics and surgeon experience into account.

Our study has several potential limitations. First, as an observational study, unmeasured confounders may affect the results. However, we have made every effort

to achieve sufficient adjustment using patient and surgeon characteristics, and tried several methods to reduce confounding. Though we still lack information on receiving teams in the wards and ICUs, the personnel distribution among different wards and ICUs is similar and they provide equal care to the patients regardless of the procedure order. There are no data to distinguish between on-pump arrest and on-pump beating CABG, but hardly any on-pump CABG was performed with a beating heart in our hospital. Second, as a retrospective analysis, we cannot estimate surgeons' extra workload other than recorded procedures, such as working on medical records, or measure their subjective fatigue by standard scales. Moreover, we did not choose the operative time to represent the workload because primary surgeons usually perform key surgical steps only, and the operative time does not really capture their exact operating time. Third, we chose a composite endpoint as our primary outcome that has both advantages and disadvantages.<sup>35</sup> Though without formal validation, the composite is widely used in related fields and we carefully interpreted the composite and individual endpoints. Last, we took CABG (representing technically complicated surgeries) and in-hospital outcomes as a specific scenario to study this question, and the study was limited to a single hospital, which may limit the generalisation of study findings. Further research in different clinical settings is required to unravel the mechanism for the observed findings, and longer term effects need to be taken into account.

In conclusion, procedure order of the surgeon in a day impacts patient outcomes after off-pump CABG, as undergoing a non-first procedure was associated with higher rates of in-hospital events, but outcomes of on-pump CABG did not appear to be affected by procedure order. Further research is needed to unravel the underlying mechanism and put forward quality-improving measures, such as improvement in the daily workflow patterns.

#### Author affiliations

<sup>1</sup>National Clinical Research Center of Cardiovascular Diseases, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, People's Republic of China

<sup>2</sup>Department of Cardiovascular Surgery, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, People's Republic of China

<sup>3</sup>National Health Commission Key Laboratory of Cardiovascular Regenerative Medicine, Fuwai Central-China Hospital, Central-China Branch of National Center for Cardiovascular Diseases, Zhengzhou, People's Republic of China

**Acknowledgements** We appreciate the multiple contributions made by the research team at the Fuwai Hospital, National Center for Cardiovascular Diseases, particularly in data collection and auditing by Ying Zhang and Qiongqiong Yang. We appreciate Cheng Sun for sharing his idea, and Jianyu Qu for his support in data cleaning.

**Collaborators** Chinese Cardiac Surgery Registry Collaborative Group.



**Contributors** Conception and design: ZZ, DZ, DG. Acquisition, analysis and interpretation of data: all authors. Drafting the manuscript: DZ, DG. Revising the manuscript for important intellectual content: all authors. Statistical analysis: DZ, SC. Obtained funding: ZZ. Administrative, technical, and material support: ZZ. Supervision: ZZ. Guarantor: ZZ.

**Funding** This study was funded by the National Key Research and Development Program (grant number: 2016YFC1302000), Beijing Municipal Commission of Science and Technology Project (grant number: D171100002917001), and the Graduate Innovation Fund of Peking Union Medical College (grant number: 2019-1002-30).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** The local institutional review board at Fuwai Hospital approved the study (approval number: 2017-880) and waived the requirement to obtain informed consent.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** No data are available.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

#### ORCID iD

Zhe Zheng <http://orcid.org/0000-0002-9162-6492>

#### REFERENCES

- Liu JH, Etzioni DA, O'Connell JB, *et al.* The increasing workload of general surgery. *Arch Surg* 2004;139:423–8.
- GBD 2016 Healthcare Access and Quality Collaborators. Measuring performance on the healthcare access and quality index for 195 countries and territories and selected subnational locations: a systematic analysis from the global burden of disease study 2016. *Lancet* 2018;391:2236–71.
- Vervoort D, Meuris B, Meyns B, *et al.* Global cardiac surgery: access to cardiac surgical care around the world. *J Thorac Cardiovasc Surg* 2020;159:987–96.
- Barger LK, Ayas NT, Cade BE, *et al.* Impact of extended-duration shifts on medical errors, adverse events, and attentional failures. *PLoS Med* 2006;3:e487.
- Whelehan DF, Alexander M, Ridgway PF. Would you allow a sleepy surgeon operate on you? A narrative review. *Sleep Med Rev* 2020;53:101341.
- Elliott DJ, Young RS, Brice J, *et al.* Effect of hospitalist workload on the quality and efficiency of care. *JAMA Intern Med* 2014;174:786–93.
- Bagrodia A, Rachakonda V, Delafuente K, *et al.* Surgeon fatigue: impact of case order on perioperative parameters and patient outcomes. *J Urol* 2012;188:1291–6.
- Silber JH, Bellini LM, Shea JA, *et al.* Patient safety outcomes under flexible and standard resident duty-hour rules. *N Engl J Med* 2019;380:905–14.
- Nuckols TK, Bhattacharya J, Wolman DM, *et al.* Cost implications of reduced work hours and workloads for resident physicians. *N Engl J Med* 2009;360:2202–15.
- Alexander JH, Smith PK. Coronary-Artery bypass grafting. *N Engl J Med* 2016;374:1954–64.
- Head SJ, Börgermann J, Osnabrugge RLJ, *et al.* Coronary artery bypass grafting: Part 2--optimizing outcomes and future prospects. *Eur Heart J* 2013;34:2873–86.
- Blackstone EH, Sabik JF. Changing the discussion about on-pump versus off-pump CABG. *N Engl J Med* 2017;377:692–3.
- Lazar HL. Should off-pump coronary artery bypass grafting be abandoned? *Circulation* 2013;128:406–13.
- Alexander JH. Clinical-outcome trials in cardiac surgery--have we primed the pump? *N Engl J Med* 2013;368:1247–8.
- Gu D, Zhang X, Diao X, *et al.* Surgeon-Specific quality monitoring system for coronary artery bypass grafting. *Ann Thorac Surg* 2019;107:705–10.
- Rao C, Zhang H, Gao H, *et al.* The Chinese cardiac surgery registry: design and data audit. *Ann Thorac Surg* 2016;101:1514–20.
- Hu S, Zheng Z, Yuan X, *et al.* Coronary artery bypass graft: contemporary heart surgery center performance in China. *Circ Cardiovasc Qual Outcomes* 2012;5:214–21.
- Yuan X, Zhang H, Zheng Z, *et al.* Trends in mortality and major complications for patients undergoing coronary artery bypass grafting among urban teaching hospitals in China: 2004 to 2013. *Eur Heart J Qual Care Clin Outcomes* 2017;3:312–8.
- Hicks KA, Tcheng JE, Bozkurt B, *et al.* 2014 ACC/AHA key data elements and definitions for cardiovascular endpoint events in clinical trials: a report of the American College of Cardiology/American heart association Task force on clinical data standards (writing Committee to develop cardiovascular endpoints data standards). *Circulation* 2015;132:302–61.
- Wang Y, Eldridge N, Metersky ML, *et al.* National trends in patient safety for four common conditions, 2005-2011. *N Engl J Med* 2014;370:341–51.
- Hu Z, Chen S, Du J, *et al.* An in-hospital mortality risk model for patients undergoing coronary artery bypass grafting in China. *Ann Thorac Surg* 2020;109:1234–42.
- Lee A, Iskander JM, Gupta N, *et al.* Queue position in the endoscopic schedule impacts effectiveness of colonoscopy. *Am J Gastroenterol* 2011;106:1457–65.
- Gruskay J, Kepler C, Smith J, *et al.* Is surgical case order associated with increased infection rate after spine surgery? *Spine* 2012;37:1170–4.
- Li X, Zhang Q, Dong J, *et al.* Impact of surgical case order on peri-operative outcomes for total joint arthroplasty. *Int Orthop* 2018;42:2289–94.
- Steuer J, Hörte L-G, Lindahl B, *et al.* Impact of perioperative myocardial injury on early and long-term outcome after coronary artery bypass grafting. *Eur Heart J* 2002;23:1219–27.
- Domanski MJ, Mahaffey K, Hasselblad V, *et al.* Association of myocardial enzyme elevation and survival following coronary artery bypass graft surgery. *JAMA* 2011;305:585–91.
- Gaudino M, Angiolillo DJ, Di Franco A, *et al.* Stroke after coronary artery bypass grafting and percutaneous coronary intervention: incidence, pathogenesis, and outcomes. *J Am Heart Assoc* 2019;8:e013032.
- Govindarajan A, Urbach DR, Kumar M, *et al.* Outcomes of daytime procedures performed by attending surgeons after night work. *N Engl J Med* 2015;373:845–53.
- Rothschild JM, Keohane CA, Rogers S, *et al.* Risks of complications by attending physicians after performing nighttime procedures. *JAMA* 2009;302:1565–72.

- 30 Althoff FC, Wachtendorf LJ, Rostin P, *et al.* Effects of night surgery on postoperative mortality and morbidity: a multicentre cohort study. *BMJ Qual Saf* 2021;30:678–88.
- 31 Kork F, Spies C, Conrad T, *et al.* Associations of postoperative mortality with the time of day, week and year. *Anaesthesia* 2018;73:711–8.
- 32 Nemeth S, Schnell S, Argenziano M, *et al.* Daytime variation does not impact outcome of cardiac surgery: results from a diverse, multi-institutional cardiac surgery network. *J Thorac Cardiovasc Surg* 2021;162:56–67.
- 33 Sessler DI, Kurz A, Saager L, *et al.* Operation timing and 30-day mortality after elective general surgery. *Anesth Analg* 2011;113:1423–8.
- 34 Ronaldson A, Kidd T, Poole L, *et al.* Diurnal cortisol rhythm is associated with adverse cardiac events and mortality in coronary artery bypass patients. *J Clin Endocrinol Metab* 2015;100:3676–82.
- 35 Montori VM, Permyer-Miralda G, Ferreira-González I, *et al.* Validity of composite end points in clinical trials. *BMJ* 2005;330:594–6.

## SUPPLEMENTAL MATERIAL

### Supplemental Methods:

Method 1. Description of coronary artery bypass grafting (CABG) procedure

Method 2. Definition of the outcomes

### Supplemental Tables

Table 1. Completeness of key variables (Created by the authors)

Table 2. Covariates in the models (Created by the authors)

Table 3. Surgeon characteristics in the CABG cohorts, by cardiopulmonary-bypass and procedure order (Created by the authors)

Table 4. Patient demographic and clinical characteristics in propensity score matched cohorts (Created by the authors)

Table 5. Outcome difference associated with non-first versus first procedure groups in propensity score weighted and matched cohorts (Created by the authors)

Table 6. Outcome difference associated with non-first versus first procedure in patients performed by the same surgeon-assistant pairs (Created by the authors)

Table 7. Outcome difference associated with non-first versus first procedure in patients undergoing CABG after 11:00 am (Created by the authors)

### Supplemental Figures

Figure 1. Distribution of number of adverse events (Created by the authors)

Figure 2. Association between later procedure order and adverse events composite compared to the first procedure in the on-pump and off-pump CABG cohorts (Created by the authors)

## **Method 1. Description of Coronary Artery Bypass Grafting (CABG) Procedure**

### **Surgeon qualification**

In Fuwai Hospital, all surgeons should have specialized in congenital or valve heart surgery for more than 3 years before undertaking any CABG procedures. Specially for off-pump CABG, the surgeon needs to perform at least 100 on-pump CABG procedures first, so that he can get the qualification to carry out the off-pump procedure. Once qualified, the choice of off-pump CABG or on-pump CABG for a specific patient was generally decided by individual surgeons.

### **Standard surgical process**

All procedures were performed with standard bypass techniques. Anesthesia was managed by inhalation of isoflurane with the addition of fentanyl or sufentanil, and propofol was administered continuously until the end of the procedure if necessary. Surgical revascularization was performed using standard bypass techniques. For on-pump CABG, a standard cardiopulmonary bypass was established, and moderate systemic hypothermia (28°C to 32°C) and perfusion with antegrade intermittent cold crystalloid cardioplegia were used. Heparin was given to achieve activated clotting times of 480 seconds or above before institution of cardiopulmonary bypass. For off-pump CABG, stabilization devices were used to provide a motionless anastomosis site, and heparin was administered before the start of the first distal anastomosis to achieve an activated clotting time of 300 to 350 seconds. On-pump CABG involved aortic cross-clamping and cardioplegic arrest, while off-pump CABG was performed with a partial occlusion clamp. Whenever possible, complete revascularization was attempted, and the internal thoracic artery was used preferentially for revascularization

of the left anterior descending artery. The remaining vessels were to be bypassed either using another arterial conduit or the saphenous vein in the configuration decided by the surgeon. During reperfusion, the bypass grafting was completed with proximal anastomoses to the ascending aorta. The decision to switch to cardiopulmonary bypass during the procedure was based on significant hemodynamic instability or ventricular arrhythmia. After separation from cardiopulmonary bypass or on completion of all anastomoses, protamine was given to reverse the effects of heparin. Postoperatively, starting within the first 24 hours, aspirin therapy (100 mg/d) is recommended and should be continued indefinitely.

### **Role of the surgical team**

The primary surgeon was in charge of the intricate core process, including the distal and proximal anastomoses for coronary revascularization, and cardiopulmonary bypass (CPB) establishment for on-pump CABG. The surgical assistants were typically responsible for incision on the chest wall, vein and artery conduit harvesting, assisting the primary surgeon with revascularization, closing chest and suturing incision. Other assisting team members included anesthesiologists, nurses, perfusionists, and cardiac intensivists, and they usually stay in operating room or intensive care unit for all surgeries through the day.

## Method 2. Definition of the outcomes

**In-hospital mortality** was defined as death before discharge from any cause.

**Myocardial infarction (MI)** was defined as an absolute rise in cardiac troponin to over 70 times the 99th percentile upper reference limit, or an increase in CK-MB to over 10 times the 99th percentile upper reference limit.

**Stroke** was defined as brain, spinal cord, or retinal cell death attributable to focal arterial ischemia, based on:

- (1) pathological, neuroimaging, or other objective evidence of cerebral, spinal cord, or retinal focal ischemic injury in a defined vascular distribution; or
- (2) clinical evidence of cerebral, spinal cord, or retinal focal ischemic injury in a defined vascular distribution with symptoms persisting >24 h or until death, and other etiologies were excluded.

**Acute kidney injury (AKI)** was defined as (1) an increase in the serum creatinine level over 3 times the latest preoperative level or a serum creatinine level reaching 4.0 mg/dL with a minimum increase of 0.5 mg/dL relative to the last preoperative value and/or (2) a new postoperative requirement for dialysis.

**Reoperation** was defined as re-exploration for mediastinal bleeding with or without tamponade either in the intensive care unit or after returning to the operating room.

**Table 1. Completeness of key variables (Created by the authors)**

Variable	Completeness	Variable	Completeness
Age	100.00%	Myocardial infarction	100.00%
Angina	100.00%	NYHA score	96.32%
Aortic calcification	99.96%	Number of grafts	100.00%
Aortic regurgitation	100.00%	Number of lesions	99.85%
Aortic stenosis	100.00%	Number of other arterial grafts	100.00%
Body mass index	99.73%	Number of vein grafts	100.00%
CCS angina grade	100.00%	Operative time	99.95%
Diabetes	100.00%	Peripheral artery disease	100.00%
EuroScore	99.96%	Preoperative LVEF	99.99%
Gender	100.00%	Prior cardiac surgery	100.00%
Glomerular filtration rate	99.94%	Prior COPD	100.00%
Hyperlipidemia	100.00%	Prior myocardial infarction	100.00%
Hypertension	100.00%	Prior PCI	100.00%
Internal mammary artery	100.00%	Prior stroke	100.00%
Left main stem disease	99.83%	Smoking	100.00%
Mitral regurgitation	100.00%	Tricuspid stenosis	100.00%
Mitral stenosis	100.00%	Tricuspid regurgitation	100.00%

CCS indicates Canadian Cardiovascular Society; COPD, chronic obstructive pulmonary disease; EuroScore, European System for Cardiac Operative Risk Evaluation; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; PCI, percutaneous coronary intervention

**Table 2. Covariates in the models (Created by the authors)**

Dependent variable	Statistical method	Covariates
AEC	Mixed-effects model with Poisson link function	<ul style="list-style-type: none"> <li>• Patient-level: Age, Angina, Aortic calcification, Body mass index, Gender, Glomerular filtration rate, Left main stem disease, NHYA III/IV, Number of lesions, Preoperative LVEF, Prior cardiac surgery, Prior COPD, Prior myocardial infarction, Prior PCI, Prior stroke;</li> <li>• Surgeon-level: years in practice and CABG volume before the index procedure;</li> <li>• Time (year of the surgery)</li> </ul>
Death	Mixed-effects model with Logit link function	<ul style="list-style-type: none"> <li>• Patient-level: Age, Gender, NHYA III/IV, Preoperative LVEF, Prior cardiac surgery. Because of the low event rate and limited sample size, we chose the variables with greater effect size and of clinical importance;</li> <li>• Surgeon-level: years in practice and CABG volume before the index procedure;</li> <li>• Time (year of the surgery)</li> </ul>
MI	Mixed-effects model with Logit link function	The same as covariates of AEC in the primary analysis
Stroke	Mixed-effects model with Logit link function	The same as covariates of AEC in the primary analysis
AKI	Mixed-effects model with Logit link function	The same as covariates of AEC in the primary analysis
Reoperation	Mixed-effects model with Logit link function	The same as covariates of AEC in the primary analysis, except for excluding the prior COPD. Because 0 case had prior COPD in patients with reoperation among off-pump CABG and the model couldn't converge without removing it.
Non-first procedure group	Logistic regression to calculate propensity score	All the variables listed in above online supplemental table 1 and the time variable (year of the surgery)

AEC indicates adverse events composite; AKI, acute kidney injury; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; LVEF, left ventricular ejection fraction; MI, myocardial infarction; NHYA, New York Heart Association; PCI, percutaneous coronary intervention.



**Table 3. Surgeon characteristics in the CABG cohorts, by cardiopulmonary-bypass and procedure order (Created by the authors)**

	Overall	First procedure	Non-first procedure	P value‡
<b>All patients</b>				
Number of cases	21866	16759	5107	
Number of surgeons *	47	47	47	
Years in practice, median (IQR), y	24 (19, 28)	23 (19, 28)	24 (21, 29)	<0.001
CABG volume, median (IQR)	460 (252, 789)	424 (231, 731)	595 (348, 953)	<0.001
Number of surgeon-assistant pairs †	316	316	316	
Number of cases treated by pairs	19785	14838	4947	
<b>On-pump cohort</b>				
Number of cases	10109	8480	1629	
Number of surgeons *	47	47	47	
Years in practice, median (IQR), y	21 (17, 26)	21 (17, 26)	21 (16, 26)	0.09
CABG volume, median (IQR)	306 (172, 501)	297 (170, 487)	357 (190, 565)	<b>&lt;0.001</b>
Number of surgeon-assistant pairs †	285	277	250	
Number of cases treated by pairs	8600	7063	1537	
<b>Off-pump cohort</b>				
Number of cases	11757	8279	3478	
Number of surgeons *	41	40	32	
Years in practice, median (IQR), y	25 (22, 30)	25 (21, 30)	26 (23, 30)	<b>&lt;0.001</b>
CABG volume, median (IQR)	661 (388, 1006)	617 (359, 972)	757 (464, 1070)	<b>&lt;0.001</b>
Number of surgeon-assistant pairs †	243	237	181	
Number of cases treated by pairs	11185	7775	3410	

CABG indicates coronary artery bypass grafting; IQR, interquartile range.

\* Number of surgeons indicates the number of surgeons who had experience performing both first and non-first CABG procedures.

† Number of surgeon-assistant pair indicates the number of surgeon and assistant pairs who had ever performed CABG as both first and non-first CABG procedures, as well as had performed as least 5 CABG cases.

‡ The Kruskal-Wallis test was used to compare the surgeon characteristics between first and non-first procedure group in specific CABG cohorts. Significant differences (<0.05) are in bold. The years in practice and CABG volumes were calculated at patient level.

**Table 4. Patient demographic and clinical characteristics in propensity score matched cohorts (Created by the authors)**

	Overall	Matched on-pump CABG cohort (n=3084)			Matched off-pump CABG cohort (n=6882)		
		First procedure (n=1542)	Non-first procedure (n=1542)	Absolute standardized difference <sup>‡</sup>	First procedure (n=3441)	Non-first procedure (n=3441)	Absolute standardized difference <sup>‡</sup>
Age, mean (SD), y	61.1 (8.7)	60.3 (8.7)	60.5 (8.5)	0.0178	61.7 (8.9)	61.2 (8.5)	0.0587
Female, No. (%)	2319 (23.3)	346 (22.4)	365 (23.7)	0.0293	807 (23.5)	801 (23.3)	0.0041
BMI, mean (SD), kg/m <sup>2</sup> *	25.7 (3.1)	25.7 (3.1)	25.8 (3.1)	0.0155	25.7 (3)	25.6 (3.1)	0.0143
Smoking, No. (%)	5569 (55.9)	823 (53.4)	824 (53.4)	0.0013	1957 (56.9)	1965 (57.1)	0.0047
Diabetes, No. (%)	2219 (22.3)	344 (22.3)	335 (21.7)	0.0141	786 (22.8)	754 (21.9)	0.0223
Hypertension, No. (%)	6393 (64.1)	1000 (64.9)	1016 (65.9)	0.0218	2215 (64.4)	2162 (62.8)	0.032
Hyperlipidemia, No. (%)	7409 (74.3)	1126 (73.0)	1116 (72.4)	0.0146	2592 (75.3)	2575 (74.8)	0.0114
Prior COPD, No. (%)	73 (0.7)	7 (0.5)	9 (0.6)	0.0181	29 (0.8)	28 (0.8)	0.0032
Prior MI, No. (%)	3693 (37.1)	557 (36.1)	594 (38.5)	0.0496	1296 (37.7)	1246 (36.2)	0.0301
Prior stroke, No. (%)	782 (7.8)	121 (7.8)	112 (7.3)	0.0221	269 (7.8)	280 (8.1)	0.0118
Prior PCI, No. (%)	533 (5.3)	70 (4.5)	85 (5.5)	0.0445	198 (5.8)	180 (5.2)	0.023
Prior cardiac surgery, No. (%)	363 (3.6)	45 (2.9)	56 (3.6)	0.0401	126 (3.7)	136 (4.0)	0.0152
Angina, No. (%)	8591 (86.2)	1271 (82.4)	1281 (83.1)	0.0172	3023 (87.9)	3016 (87.6)	0.0062
NYHA III/IV, No. (%)	3391 (34.0)	474 (30.7)	450 (29.2)	0.034	1242 (36.1)	1225 (35.6)	0.0103
eGFR, median (IQR), mL/min/1.73 m <sup>2</sup> †	94 (23.9)	94.1 (23.6)	93.8 (23.6)	0.012	93.9 (24.1)	94.2 (23.9)	0.0113
PAD, No. (%)	1366 (13.7)	137 (8.9)	135 (8.8)	0.0046	535 (15.5)	559 (16.2)	0.0191
LM disease, No. (%)	1784 (17.9)	262 (17.0)	258 (16.7)	0.0064	620 (18.0)	644 (18.7)	0.0186
Three-vessel disease, No. (%)	6174 (62.0)	1092 (70.8)	1052 (68.2)	0.0564	2072 (60.2)	1958 (56.9)	0.0673
Preoperative LVEF, mean (SD)	61.2 (7.4)	60.8 (8.0)	60.7 (7.7)	0.018	61.3 (7.3)	61.5 (6.9)	0.0282
Aortic calcification, No. (%)	941 (9.4)	87 (5.6)	97 (6.3)	0.0275	394 (11.5)	363 (10.5)	0.0287
EuroSCORE>3, No. (%)	2384 (23.9)	308 (20.0)	285 (18.5)	0.0379	919 (26.7)	872 (25.3)	0.0311

BMI indicates body mass index; CABG, coronary artery bypass grafting; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; EuroSCORE, European System for Cardiac Operative Risk Evaluation; IQR, interquartile range; LVEF, left ventricular ejection fraction; LM, left main stem; MI, myocardial infarction; NYHA, New York Heart Association; PAD, peripheral artery disease; PCI, percutaneous coronary intervention; SMD, standardized mean difference; SD, standard deviation.

\*  $\text{BMI (kg/m}^2\text{)} = \text{weight (kg)} / (\text{height [m]})^2$ .

†  $\text{eGFR (mL/min/1.73 m}^2\text{)} = 186 \times (\text{serum creatinine level [mg/dL]})^{-1.154} \times (\text{age [y]})^{-0.203}$ .

‡ Absolute standardized difference was calculated to measure the difference between first and non-first procedure groups in specific CABG cohorts. The standardized difference being less than 0.1 shows the matched groups were well-balanced.

**Table 5. Outcome difference associated with non-first versus first procedure groups in propensity score weighted and matched cohorts**

(Created by the authors)

	IPTW method*		1:1 propensity matching method†			
	RR <sub>adj</sub> /OR <sub>adj</sub> ‡ (95% CI)	P value	First procedure	Non-first procedure	RR <sub>adj</sub> /OR <sub>adj</sub> ‡ (95% CI)	P value
<b>On-pump CABG cohort</b>						
<b>Total patients</b>			1542	1542		
AEC, total number	0.92 (0.81-1.03)	0.15	221	192	0.87 (0.71-1.06)	0.17
one event, n (%)			184 (11.9)	155 (10.1)		
two events, n (%)			14 (0.9)	14 (0.9)		
three events, n (%)			3 (0.2)	3 (0.2)		
four events, n (%)			/	/		
Death, n (%)	1.47 (0.85-2.54)	0.17	4 (0.3)	7 (0.45)	1.75 (0.51-5.98)	0.37
MI, n (%)	0.84 (0.69-1.03)	0.09	60 (3.9)	66 (4.3)	1.12 (0.77-1.63)	0.56
Stroke, n (%)	0.63 (0.42-0.96)	<b>0.03</b>	26 (1.7)	11 (0.7)	0.42 (0.21-0.86)	<b>0.02</b>
AKI, n (%)	0.98 (0.83-1.16)	0.81	120 (7.8)	94 (6.1)	0.77 (0.58-1.02)	0.06
Reoperation, n (%)	0.73 (0.50-1.05)	0.09	11 (0.7)	14 (0.9)	1.27 (0.58-2.80)	0.55
<b>Off-pump CABG cohort</b>						
<b>Total patients</b>			3441	3441		
AEC*, total number	1.28 (1.14-1.45)	<b>&lt;0.001</b>	291	384	1.32 (1.13-1.54)	<b>&lt;0.001</b>
one event, n (%)			263 (7.6)	315 (9.2)		
two events, n (%)			14 (0.4)	27 (0.8)		
three events, n (%)			/	5 (0.15)		
four events, n (%)			/	/		
Death, n (%)	2.04 (1.04-4.01)	<b>0.04</b>	5 (0.15)	12 (0.35)	2.40 (0.85-6.81)	0.10
MI, n (%)	1.46 (1.18-1.82)	<b>0.001</b>	66 (1.9)	112 (3.3)	1.74 (1.27-2.38)	<b>&lt;0.001</b>

Stroke, n (%)	1.69 (1.15-2.47)	<b>0.007</b>	22 (0.6)	39 (1.1)	1.77 (1.05-2.99)	<b>0.03</b>
AKI, n (%)	1.20 (1.01-1.43)	<b>0.04</b>	157 (4.6)	175 (5.1)	1.12 (0.90-1.39)	0.31
Reoperation, n (%)	0.94 (0.69-1.28)	0.68	41 (1.2)	46 (1.3)	1.12 (0.73-1.72)	0.59

AEC indicates adverse events composite, which is defined as the number of any adverse events occurred, including death, myocardial infarction, stroke, acute kidney injury or reoperation; AKI indicates acute kidney injury; CABG, coronary artery bypass grafting; CI, confidence interval; EuroSCORE, European System for Cardiac Operative Risk Evaluation; MI, myocardial infarction; OR<sub>adj</sub>, adjusted odds ratio; RR<sub>adj</sub>, adjusted risk ratio.

\* The propensity score is calculated by establishing a multivariable logistic regression model with all patient characteristics (online supplemental table 1) included to estimate the propensity for patients to undergo a non-first procedure. The inverse probability of treatment weighting (IPTW) is used based on the inverse of the propensity score and the mixed-effects model analyses repeated.

† The 1:1 propensity score matching method is used to form matched CABG cohorts by matching first vs non-first patients treated by the same surgeon with similar propensity score using the greedy nearest neighbor matching algorithm without replacement, with a caliper width at 0.01 according to the absolute value of the difference between propensity scores.

‡ For the adverse events composite, the adjusted rate ratio (non-first vs first) is estimated from the model with a Poisson link function; For every individual event (binary variable, i.e., death, MI, stroke, AKI and reoperation), the adjusted odds ratio (non-first vs first) for each outcome is estimated from the model with a logit link function. Significant differences (<0.05) are in bold.

**Table 6. Outcome difference associated with non-first versus first procedure in patients performed by the same surgeon-assistant pairs (Created by the authors)**

Outcome	On-pump CABG cohort (n=8600)				Off-pump CABG cohort (n=11185)			
	First procedure (n=7063)	Non-first procedure (n=1537)	RR <sub>adj</sub> */OR <sub>adj</sub> † (95% CI)	P value	First procedure (n=7775)	Non-first procedure (n=3410)	RR <sub>adj</sub> */OR <sub>adj</sub> † (95% CI)	P value
AEC*, total number	982	193	0.89 (0.76-1.05)	0.17	720	385	1.29 (1.14-1.47)	<b>&lt;0.001</b>
one event, n (%)	817 (11.6)	153 (10.0)			621 (8.0)	314 (9.2)		
two events, n (%)	62 (0.9)	17 (1.1)			43 (0.6)	28 (0.8)		
three events, n (%)	11 (0.2)	2 (0.1)			3 (0.0)	5 (0.1)		
four events, n (%)	2 (0.0)	0 (0.0)			1 (0.0)	0 (0.0)		
Death, n (%)	25 (0.4)	8 (0.5)	1.51 (0.98-2.33)	0.06	16 (0.2)	12 (0.4)	2.31 (1.49-3.57)	<b>&lt;0.001</b>
MI, n (%)	311 (4.4)	68 (4.4)	0.95 (0.74-1.22)	0.70	194 (2.5)	112 (3.3)	1.40 (1.12-1.75)	<b>0.003</b>
Stroke, n (%)	87 (1.2)	10 (0.7)	0.54 (0.30-0.97)	<b>0.04</b>	53 (0.7)	39 (1.1)	1.82 (1.30-2.57)	<b>0.001</b>
AKI, n (%)	469 (6.6)	91 (5.9)	0.91 (0.69-1.20)	0.52	343 (4.4)	176 (5.2)	1.23 (1.00-1.52)	0.05
Reoperation, n (%)	90 (1.3)	16 (1.0)	0.73 (0.46-1.16)	0.18	114 (1.5)	46 (1.3)	0.98 (0.69-1.39)	0.93

AEC, adverse event composite; AKI indicates acute kidney injury; CABG, coronary artery bypass grafting; CI, confidence interval; MI, myocardial infarction; OR<sub>adj</sub>, adjusted odds ratio; RR<sub>adj</sub>, adjusted risk ratio.

\*For the adverse events composite (AEC, count variable, the “n (%)” presents the number of patients with given number of events and the percentage of those patients among specific procedure group. A mixed-effects model with a Poisson link function and surgeon-specific random intercepts is fitted. The risk-adjusted rate ratio (non-first vs first) is estimated, and when it achieves a statistically significant level of 0.05, the P value is in bold.

†For every individual event (binary variable, i.e., death, MI, stroke, AKI and reoperation), a mixed-effects model with a logit link function and surgeon-specific random intercepts is fitted. The risk-adjusted odds ratio (non-first vs first) for each outcome is estimated, and when it achieves a statistically significant level of 0.05, the P value is in bold.


**Table 7. Outcome difference associated with non-first versus first procedure in patients undergoing CABG after 11:00 am (Created by the authors)**

Outcome	On-pump CABG cohort (n=4604)				Off-pump CABG cohort (n=4992)			
	First procedure (n=3061)	Non-first procedure (n=1543)	RR <sub>adj</sub> */OR <sub>adj</sub> † (95% CI)	P value	First procedure (n=1619)	Non-first procedure (n=3303)	RR <sub>adj</sub> */OR <sub>adj</sub> † (95% CI)	P value
AEC*, total number	406	191	0.91 (0.74-1.12)	0.39	149	303	1.30 (1.04-1.64)	<b>0.02</b>
one event, n (%)	342 (11.2)	150 (9.7)			133 (8.2)	303 (9.2)		
two events, n (%)	26 (0.8)	16 (1.0)			8 (0.5)	28 (0.8)		
three events, n (%)	4 (0.1)	3 (0.2)			0 (0.0)	5 (0.2)		
Death, n (%)	9 (0.3)	9 (0.6)	1.06 (0.54-2.07)	0.87	4 (0.2)	12 (0.4)	2.04 (0.74-5.66)	0.17
MI, n (%)	131 (4.3)	66 (4.3)	0.93 (0.64-1.34)	0.69	39 (2.4)	107 (3.2)	1.59 (1.05-2.40)	<b>0.03</b>
Stroke, n (%)	39 (1.3)	10 (0.6)	0.50 (0.25-1.01)	0.05	11 (0.7)	39 (1.2)	1.58 (0.78-3.19)	0.20
AKI, n (%)	190 (6.2)	88 (5.7)	0.91 (0.66-1.27)	0.59	74 (4.6)	171 (5.2)	1.09 (0.75-1.57)	0.65
Reoperation, n (%)	37 (1.2)	18 (1.2)	0.86 (0.45-1.64)	0.65	21 (1.3)	45 (1.4)	1.10 (0.61-1.98)	0.76

AEC, adverse event composite; AKI indicates acute kidney injury; CABG, coronary artery bypass grafting; CI, confidence interval; MI, myocardial infarction; OR<sub>adj</sub>, adjusted odds ratio; RR<sub>adj</sub>, adjusted risk ratio.


\*For the adverse events composite (AEC, count variable, the “n (%)” presents the number of patients with given number of events and the percentage of those patients among specific procedure group. A mixed-effects model with a Poisson link function and surgeon-specific random intercepts is fitted. The risk-adjusted rate ratio (non-first vs first) is estimated, and when it achieves a statistically significant level of 0.05, the P value is in bold.

†For every individual event (binary variable, i.e., death, MI, stroke, AKI and reoperation), a mixed-effects model with a logit link function and surgeon-specific random intercepts is fitted. The risk-adjusted odds ratio (non-first vs first) for each outcome is estimated, and when it achieves a statistically significant level of 0.05, the P value is in bold.



**Figure 1. Distribution of number of adverse events (Created by the authors)**  
CABG, coronary artery bypass grafting.





**Figure 2. Association between later procedure order and adverse events composite compared to the first procedure in the on-pump and off-pump CABG cohorts (Created by the authors)**

CABG, coronary artery bypass grafting; CI, confidence interval; RR, rate ratio.