

Appendix Table 1. Characteristics of clients with or without AE (all variables tested)

Characteristic	Adverse events			
	No	Yes	Total	p-value <sup>1</sup>
Sex: Proportion of Female (%) <sup>2</sup>	59.87	54.32	59.50	0.325
Age (years)	71.20	75.93	71.52	0.016
Client was at private home/apt before home care admission (%)	54.78	55.56	54.83	0.892
At the time of index admission, living alone (%)	37.44	33.33	37.17	0.460
Number of key informal helpers:				
None	83	3	86	
1-2	868	76	944	
3 or more	41	2	43	
Total	992	81	1073	0.318
Length of stay (days)	347.31	465.60	355.30	0.089
Difficulty with communication at intake of index admission (%)	12.34	21.25	12.95	0.220
Difficulty with communication later on during index admission (%)	14.46	23.75	15.10	0.025
Difficulty with communication at intake or later on during index admission (%)	15.35	24.05	15.94	0.041
Discharge alive from index admission (%)	88.74	82.72	88.33	0.103
Significant procedure performed before admission (%)	29.94	23.46	29.50	0.217
Cognitive patterns problems (out of 3) <sup>4</sup>	.65	1.12	.68	0.004
Communication problems (out of 4)	.71	.96	.72	0.017
Mood & Behaviour problems (out of 2)	.31	.49	.32	0.022
IADL score = (1 independent, 4 dependent)	2.35	2.82	2.39	0.000
IADL (% independent)	43.9	27.4	42.6	0.000
ADL (1 independent, 4 dependent)	1.63	1.86	1.65	0.015
ADL (% independent)	68.5	59.3	67.8	0.011
Client functional status has deteriorated during admission (%)	24.88	56.79	27.11	0.000
Pain (%)	55.57	75.38	57.01	0.002
History of falls (%)	30.00	43.84	31.09	0.014
Notation of environmental home hazard (%)	16.25	24.64	16.86	0.073
Psychotropic medications (%)	26.39	32.05	26.82	0.278
Medications: Number of prescriptions & over-the-counter meds:				
None	33	2	35	
1-3	154	9	163	
4-9	532	36	568	
>9	267	34	301	
Total	986	81	1067	0.015

Appendix Table 1. Characteristics of clients with or without AE (all variables tested) (cont.)

Characteristic	Adverse events			
	No	Yes	Total	p-value <sup>2</sup>
Number of comorbid conditions (out of 32)	3.94	5.56	4.05	0.000
Cardiac and vascular disease (out of 7)	1.37	2.01	1.41	0.000
Cerebrovascular accident (stroke)	.11	.17	.12	0.097
Transient ischemic attack	.08	.15	.08	0.032
Congestive heart failure	.09	.15	.09	0.079
Coronary artery disease	.21	.40	.22	0.000
Hypertension	.50	.67	.51	0.004
Peripheral vascular disease	.08	.10	.09	0.645
Other cardiac/vascular disorder	.29	.38	.30	0.093
Neurological (out of 6)	.19	.31	.20	0.068
Alzheimer	.05	.07	.05	0.345
Other dementia	.07	.17	.08	0.002
Acute brain injury	.01	.01	.01	0.953
Hemiplegia/hemiparesis	.04	.05	.04	0.782
Quadraplegia/quadraparesis	.01	0.00	.01	0.475
Multiple sclerosis	.01	0.00	.01	0.370
Senses (out of 2)	.17	.16	.17	0.856
Cataract	.13	.12	.13	0.965
Glaucoma	.04	.04	.04	0.773
Psychiatric	.12	.22	.13	0.100
Infections (out of 5)	.25	.53	.27	0.001
HIV	.00	0.00	.00	0.703
Pneumonia	.07	.14	.07	0.018
Tuberculosis	.00	0.00	.00	0.590
Urinary tract infection	.06	.20	.07	0.000
Other Infection	.11	.20	.12	0.026
Hematologic (out of 2)	.12	.11	.12	0.878
Chronic Anemia	.10	.10	.10	0.926
Other hematological disease/condition	.02	.01	.02	0.580
Cancer	.27	.25	.27	0.690
Diabetes	.26	.38	.26	0.012
Parkinson	.02	.04	.02	0.224
Musculoskeletal (out of 3)	.41	.51	.42	0.166
Arthritis	.25	.31	.26	0.259
Hip fracture	.05	.06	.05	0.728
Other fracture	.10	.14	.11	0.364
Respiratory (out of 2)	.18	.33	.19	0.020
Emphysema/Chronic obstructive lung disease	.12	.25	.13	0.001
Asthma	.06	.09	.06	0.338
Other comorbid conditions not noted above	.59	.70	.60	0.045

- 
1. p-value indicates significant difference between cases with and without adverse event, based on Chi-square test for categorical variables and t-test for continuous variables.  
None of the p-values correct for multiple comparisons.
  2. Indicates the percentage of clients with this characteristic.
  3. Indicates the average number of problems or conditions in this category.